

DEPARTMENT OF THE INTERIOR
UNITED STATES GEOLOGICAL SURVEY
CHARLES D. WALCOTT, DIRECTOR

GEOLOGIC ATLAS


OF THE

UNITED STATES

JOPLIN DISTRICT FOLIO

MISSOURI - KANSAS

INDEX MAP



SCALE 40 MILES = 1 INCH


JOPLIN DISTRICT FOLIO


OTHER PUBLISHED FOLIOS

CONTENTS

DESCRIPTIVE TEXT
TOPOGRAPHIC MAP
AREAL GEOLOGY MAP

ECONOMIC GEOLOGY MAP
MINE MAPS
ILLUSTRATION SHEET

WASHINGTON, D. C.

ENGRAVED AND PRINTED BY THE U. S. GEOLOGICAL SURVEY

GEORGE W. STOSE, EDITOR OF GEOLOGIC MAPS S. J. KUBEL, CHIEF ENGRAVER

1907

GEOLOGIC AND TOPOGRAPHIC ATLAS OF UNITED STATES.

The Geological Survey is making a geologic map of the United States, which is being issued in parts, called folios. Each folio includes a topographic map and geologic maps of a small area of country, together with explanatory and descriptive texts.

THE TOPOGRAPHIC MAP.

The features represented on the topographic map are of three distinct kinds: (1) inequalities of surface, called *relief*, as plains, plateaus, valleys, hills, and mountains; (2) distribution of water, called *drainage*, as streams, lakes, and swamps; (3) the works of man, called *culture*, as roads, railroads, boundaries, villages, and cities.

Relief.—All elevations are measured from mean sea level. The heights of many points are accurately determined, and those which are most important are given on the map in figures. It is desirable, however, to give the elevation of all parts of the area mapped, to delineate the outline or form of all slopes, and to indicate their grade or steepness. This is done by lines each of which is drawn through points of equal elevation above mean sea level, the altitudinal interval represented by the space between lines being the same throughout each map. These lines are called *contours*, and the uniform altitudinal space between each two contours is called the *contour interval*. Contours and elevations are printed in brown.

The manner in which contours express elevation, form, and grade is shown in the following sketch and corresponding contour map (fig. 1).




FIG. 1.—Ideal view and corresponding contour map.

The sketch represents a river valley between two hills. In the foreground is the sea, with a bay which is partly closed by a hooked sand bar. On each side of the valley is a terrace. From the terrace on the right a hill rises gradually, while from that on the left the ground ascends steeply, forming a precipice. Contrasted with this precipice is the gentle slope from its top toward the left. In the map each of these features is indicated, directly beneath its position in the sketch, by contours. The following explanation may make clearer the manner in which contours delineate elevation, form, and grade:

1. A contour indicates a certain height above sea level. In this illustration the contour interval is 50 feet; therefore the contours are drawn at 50, 100, 150, and 200 feet, and so on, above mean sea level. Along the contour at 250 feet lie all points of the surface that are 250 feet above sea; along the contour at 200 feet, all points that are 200 feet above sea; and so on. In the space between any two contours are found elevations above the lower and below the higher contour. Thus the contour at 150 feet falls just below the edge of the terrace, while that at 200 feet lies above the terrace; therefore all points on the terrace are shown to be more than 150 but less than 200 feet above sea. The summit of the higher hill is stated to be 670 feet above sea; accordingly the contour at 650 feet surrounds it. In this illustration all the contours are numbered, and those for 250 and 500 feet are accentuated by being made heavier. Usually it is not desirable to number all the contours, and then the accentuating and numbering of certain of them—say every fifth one—suffice, for the heights of others may be ascertained by counting up or down from a numbered contour.

2. Contours define the forms of slopes. Since contours are continuous horizontal lines, they wind smoothly about smooth surfaces, recede into all recumbent angles of ravines, and project in passing about prominences. These relations of contour curves and angles to forms of the landscape can be traced in the map and sketch.

3. Contours show the approximate grade of any slope. The altitudinal space between two contours is the same, whether they lie along a cliff or on a gentle slope; but to rise a given height on a gentle slope one must go farther than on a steep slope, and therefore contours are far apart on gentle slopes and near together on steep ones.

For a flat or gently undulating country a small contour interval is used; for a steep or mountainous country a large interval is necessary. The smallest interval used on the atlas sheets of the Geological Survey is 5 feet. This is serviceable for regions like the Mississippi delta and the Dismal Swamp. In mapping great mountain masses, like those in Colorado, the interval may be 250 feet. For intermediate relief contour intervals of 10, 20, 25, 50, and 100 feet are used.

Drainage.—Watercourses are indicated by blue lines. If a stream flows the entire year the line is drawn unbroken, but if the channel is dry a part of the year the line is broken or dotted. Where a stream sinks and reappears at the surface, the supposed underground course is shown by a broken blue line. Lakes, marshes, and other bodies of water are also shown in blue, by appropriate conventional signs.

Culture.—The works of man, such as roads, railroads, and towns, together with boundaries of townships, counties, and States, are printed in black.

Scales.—The area of the United States (excluding Alaska and island possessions) is about 3,025,000 square miles. A map representing this area, drawn to the scale of 1 mile to the inch, would cover 3,025,000 square inches of paper, and to accommodate the map the paper would need to measure about 240 by 180 feet. Each square mile of ground surface would be represented by a square inch of map surface, and one linear mile on the ground would be represented by a linear inch on the map. This relation between distance in nature and corresponding distance on the map is called the *scale* of the map. In this case it is "1 mile to an inch." The scale may be expressed also by a fraction, of which the numerator is a length on the map and the denominator the corresponding length in nature expressed in the same unit. Thus, as there are 63,360 inches in a mile, the scale "1 mile to an inch" is expressed by $\frac{1}{63,360}$.

Three scales are used on the atlas sheets of the Geological Survey; the smallest is $\frac{1}{250,000}$, the intermediate $\frac{1}{125,000}$, and the largest $\frac{1}{62,500}$. These correspond approximately to 4 miles, 2 miles, and 1 mile on the ground to an inch on the map. On the scale $\frac{1}{250,000}$ a square inch of map surface represents about 1 square mile of earth surface; on the scale $\frac{1}{125,000}$, about 4 square miles; and on the scale $\frac{1}{62,500}$, about 16 square miles. At the bottom of each atlas sheet the scale is expressed in three ways—by a graduated line representing miles and parts of miles in English inches, by a similar line indicating distance in the metric system, and by a fraction.

Atlas sheets and quadrangles.—The map is being published in atlas sheets of convenient size, which represent areas bounded by parallels and meridians. These areas are called *quadrangles*. Each sheet on the scale of $\frac{1}{250,000}$ contains one square degree—i. e., a degree of latitude by a degree of longitude; each sheet on the scale of $\frac{1}{125,000}$ contains one-fourth of a square degree; each sheet on the scale of $\frac{1}{62,500}$ contains one-sixteenth of a square degree. The areas of the corresponding quadrangles are about 4000, 1000, and 250 square miles.

The atlas sheets, being only parts of one map of the United States, disregard political boundary lines, such as those of States, counties, and townships. To each sheet, and to the quadrangle it represents, is given the name of some well-known town or natural feature within its limits, and at the sides and corners of each sheet the names of adjacent sheets, if published, are printed.

Uses of the topographic map.—On the topographic map are delineated the relief, drainage, and culture of the quadrangle represented. It should portray

to the observer every characteristic feature of the landscape. It should guide the traveler; serve the investor or owner who desires to ascertain the position and surroundings of property; save the engineer preliminary surveys in locating roads, railways, and irrigation reservoirs and ditches; provide educational material for schools and homes; and be useful as a map for local reference.

THE GEOLOGIC MAPS.

The maps representing the geology show, by colors and conventional signs printed on the topographic base map, the distribution of rock masses on the surface of the land, and the structure sections show their underground relations, as far as known and in such detail as the scale permits.

KINDS OF ROCKS.

Rocks are of many kinds. On the geologic map they are distinguished as igneous, sedimentary, and metamorphic.

Igneous rocks.—These are rocks which have cooled and consolidated from a state of fusion. Through rocks of all ages molten material has from time to time been forced upward in fissures or channels of various shapes and sizes, to or nearly to the surface. Rocks formed by the consolidation of the molten mass within these channels—that is, below the surface—are called *intrusive*. When the rock occupies a fissure with approximately parallel walls the mass is called a *dike*; when it fills a large and irregular conduit the mass is termed a *stock*. When the conduits for molten magmas traverse stratified rocks they often send off branches parallel to the bedding planes; the rock masses filling such fissures are called *sills* or *sheets* when comparatively thin, and *laccoliths* when occupying larger chambers produced by the force propelling the magmas upward. Within rock inclosures molten material cools slowly, with the result that intrusive rocks are generally of crystalline texture. When the channels reach the surface the molten material poured out through them is called *lava*, and lavas often build up volcanic mountains. Igneous rocks thus formed upon the surface are called *extrusive*. Lavas cool rapidly in the air, and acquire a glassy or, more often, a partially crystalline condition in their outer parts, but are more fully crystalline in their inner portions. The outer parts of lava flows are usually more or less porous. Explosive action often accompanies volcanic eruptions, causing ejections of dust, ash, and larger fragments. These materials, when consolidated, constitute breccias, agglomerates, and tuffs. Volcanic ejecta may fall in bodies of water or may be carried into lakes or seas and form sedimentary rocks.

Sedimentary rocks.—These rocks are composed of the materials of older rocks which have been broken up and the fragments of which have been carried to a different place and deposited. The chief agent of transportation of rock debris is water in motion, including rain, streams, and the water of lakes and of the sea. The materials are in large part carried as solid particles, and the deposits are then said to be mechanical. Such are gravel, sand, and clay, which are later consolidated into conglomerate, sandstone, and shale. In smaller portions the materials are carried in solution, and the deposits are then called organic if formed with the aid of life, or chemical if formed without the aid of life. The more important rocks of chemical and organic origin are limestone, chert, gypsum, salt, iron ore, peat, lignite, and coal. Any one of the deposits may be separately formed, or the different materials may be intermingled in many ways, producing a great variety of rocks.

Another transporting agent is air in motion, or wind; and a third is ice in motion, or glaciers. The most characteristic of the wind-borne or eolian deposits is loess, a fine-grained earth; the most characteristic of glacial deposits is till, a heterogeneous mixture of boulders and pebbles with clay or sand. Sedimentary rocks are usually made up of layers or beds which can be easily separated. These layers are called *strata*. Rocks deposited in layers are said to be stratified.

The surface of the earth is not fixed, as it seems to be; it very slowly rises or sinks, with reference to the sea, over wide expanses; and as it rises or

subsides the shore lines of the ocean are changed. As a result of the rising of the surface, marine sedimentary rocks may become part of the land, and extensive land areas are in fact occupied by such rocks.

Rocks exposed at the surface of the land are acted upon by air, water, ice, animals, and plants. They are gradually broken into fragments, and the more soluble parts are leached out, leaving the less soluble as a *residual* layer. Water washes residual material down the slopes, and it is eventually carried by rivers to the ocean or other bodies of standing water. Usually its journey is not continuous, but it is temporarily built into river bars and flood plains, where it is called *alluvium*. Alluvial deposits, glacial deposits (collectively known as *drift*), and eolian deposits belong to the *surficial* class, and the residual layer is commonly included with them. Their upper parts, occupied by the roots of plants, constitute soils and subsoils, the soils being usually distinguished by a notable admixture of organic matter.

Metamorphic rocks.—In the course of time, and by a variety of processes, rocks may become greatly changed in composition and in texture. When the newly acquired characteristics are more pronounced than the old ones such rocks are called *metamorphic*. In the process of metamorphism the substances of which a rock is composed may enter into new combinations, certain substances may be lost, or new substances may be added. There is often a complete gradation from the primary to the metamorphic form within a single rock mass. Such changes transform sandstone into quartzite, limestone into marble, and modify other rocks in various ways.

From time to time in geologic history igneous and sedimentary rocks have been deeply buried and later have been raised to the surface. In this process, through the agencies of pressure, movement, and chemical action, their original structure may be entirely lost and new structures appear. Often there is developed a system of division planes along which the rocks split easily, and these planes may cross the strata at any angle. This structure is called *cleavage*. Sometimes crystals of mica or other foliaceous minerals are developed with their laminae approximately parallel; in such cases the structure is said to be schistose, or characterized by *schistosity*.

As a rule, the oldest rocks are most altered and the younger formations have escaped metamorphism, but to this rule there are important exceptions.

FORMATIONS.

For purposes of geologic mapping rocks of all the kinds above described are divided into *formations*. A sedimentary formation contains between its upper and lower limits either rocks of uniform character or rocks more or less uniformly varied in character, as, for example, a rapid alternation of shale and limestone. When the passage from one kind of rocks to another is gradual it is sometimes necessary to separate two contiguous formations by an arbitrary line, and in some cases the distinction depends almost entirely on the contained fossils. An igneous formation is constituted of one or more bodies either containing the same kind of igneous rock or having the same mode of occurrence. A metamorphic formation may consist of rock of uniform character or of several rocks having common characteristics.

When for scientific or economic reasons it is desirable to recognize and map one or more specially developed parts of a varied formation, such parts are called *members*, or by some other appropriate term, as *lentils*.

AGES OF ROCKS.

Geologic time.—The time during which the rocks were made is divided into several *periods*. Smaller time divisions are called *epochs*, and still smaller ones *stages*. The age of a rock is expressed by naming the time interval in which it was formed, when known.

The sedimentary formations deposited during a period are grouped together into a *system*. The principal divisions of a system are called *series*. Any aggregate of formations less than a series is called a *group*.

(Continued on third page of cover.)

As sedimentary deposits or strata accumulate the younger rest on those that are older, and the relative ages of the deposits may be determined by observing their positions. This relationship holds except in regions of intense disturbance; in such regions sometimes the beds have been reversed, and it is often difficult to determine their relative ages from their positions; then *fossils*, or the remains and imprints of plants and animals, indicate which of two or more formations is the oldest.

Stratified rocks often contain the remains or imprints of plants and animals which, at the time the strata were deposited, lived in the sea or were washed from the land into lakes or seas, or were buried in surficial deposits on the land. Such rocks are called fossiliferous. By studying fossils it has been found that the life of each period of the earth's history was to a great extent different from that of other periods. Only the simpler kinds of marine life existed when the oldest fossiliferous rocks were deposited. From time to time more complex kinds developed, and as the simpler ones lived on in modified forms life became more varied. But during each period there lived peculiar forms, which did not exist in earlier times and have not existed since; these are *characteristic types*, and they define the age of any bed of rock in which they are found. Other types passed on from period to period, and thus linked the systems together, forming a chain of life from the time of the oldest fossiliferous rocks to the present. When two sedimentary formations are remote from each other and it is impossible to observe their relative positions, the characteristic fossil types found in them may determine which was deposited first. Fossil remains found in the strata of different areas, provinces, and continents afford the most important means for combining local histories into a general earth history.

It is often difficult or impossible to determine the age of an igneous formation, but the relative age of such a formation can sometimes be ascertained by observing whether an associated sedimentary formation of known age is cut by the igneous mass or is deposited upon it.

Similarly, the time at which metamorphic rocks were formed from the original masses is sometimes shown by their relations to adjacent formations of known age; but the age recorded on the map is that of the original masses and not of their metamorphism.

Colors and patterns.—Each formation is shown on the map by a distinctive combination of color and pattern, and is labeled by a special letter symbol.

Symbols, and colors assigned to the rock systems.

System.	Series.	Symbol.	Color for sedimentary rocks.
Cenozoic	Quaternary Recent Pleistocene Pliocene Miocene Oligocene Eocene	Q	Brownish-yellow.
	Tertiary	T	Yellow ocher.
	Cretaceous	K	Olive-green.
	Jurassic	J	Blue-green.
	Triassic	T	Peacock-blue.
Mesozoic	Carboniferous (Permian Pennsylvanian Mississippian)	C	Blue.
	Devonian	D	Blue-gray.
	Silurian	S	Blue-purple.
	Ordovician	O	Red purple.
	Cambrian (Saratoga Acadian Georgian)	C	Brick-red.
	Algonkian	A	Brownish-red.
	Archean	R	Gray-brown.

Patterns composed of parallel straight lines are used to represent sedimentary formations deposited in the sea or in lakes. Patterns of dots and circles represent alluvial, glacial, and eolian formations. Patterns of triangles and rhombs are used for igneous formations. Metamorphic rocks of unknown origin are represented by short dashes irregularly placed; if the rock is schist the dashes may be arranged in wavy lines parallel to the structure

planes. Suitable combination patterns are used for metamorphic formations known to be of sedimentary or of igneous origin.

The patterns of each class are printed in various colors. With the patterns of parallel lines, colors are used to indicate age, a particular color being assigned to each system. The symbols by which formations are labeled consist each of two or more letters. If the age of a formation is known the symbol includes the system symbol, which is a capital letter or monogram; otherwise the symbols are composed of small letters. The names of the systems and recognized series, in proper order (from new to old), with the color and symbol assigned to each system, are given in the preceding table.

SURFACE FORMS.

Hills and valleys and all other surface forms have been produced by geologic processes. For example, most valleys are the result of erosion by the streams that flow through them (see fig. 1), and the alluvial plains bordering many streams were built up by the streams; sea cliffs are made by the eroding action of waves, and sand spits are built up by waves. Topographic forms thus constitute part of the record of the history of the earth.

Some forms are produced in the making of deposits and are inseparably connected with them. The hooked spit, shown in fig. 1, is an illustration. To this class belong beaches, alluvial plains, lava streams, drumlins (smooth oval hills composed of till), and moraines (ridges of drift made at the edges of glaciers). Other forms are produced by erosion, and these are, in origin, independent of the associated material. The sea cliff is an illustration; it may be carved from any rock. To this class belong abandoned river channels, glacial furrows, and peneplains. In the making of a stream terrace an alluvial plain is first built and afterwards partly eroded away. The shaping of a marine or lacustrine plain is usually a double process, hills being worn away (*degraded*) and valleys being filled up (*aggraded*).

All parts of the land surface are subject to the action of air, water, and ice, which slowly wear them down, and streams carry the waste material to the sea. As the process depends on the flow of water to the sea, it can not be carried below sea level, and the sea is therefore called the *base-level* of erosion. When a large tract is for a long time undisturbed by uplift or subsidence it is degraded nearly to base-level, and the even surface thus produced is called a *peneplain*. If the tract is afterwards uplifted the peneplain at the top is a record of the former relation of the tract to sea level.

THE VARIOUS GEOLOGIC SHEETS.

Areal geology map.—This map shows the areas occupied by the various formations. On the margin is a *legend*, which is the key to the map. To ascertain the meaning of any colored pattern and its letter symbol the reader should look for that color, pattern, and symbol in the legend, where he will find the name and description of the formation. If it is desired to find any given formation, its name should be sought in the legend and its color and pattern noted, when the areas on the map corresponding in color and pattern may be traced out.

The legend is also a partial statement of the geologic history. In it the formations are arranged in columnar form, grouped primarily according to origin—sedimentary, igneous, and crystalline of unknown origin—and within each group they are placed in the order of age, so far as known, the youngest at the top.

Economic geology map.—This map represents the distribution of useful minerals and rocks, showing their relations to the topographic features and to the geologic formations. The formations which appear on the areal geology map are usually shown on this map by fainter color patterns. The areal geology, thus printed, affords a subdued background upon which the areas of productive formations may be emphasized by strong colors. A mine symbol is printed at each mine or quarry, accompanied by the name of the principal mineral mined or stone quarried. For regions where there are important mining industries or where artesian basins exist special maps are prepared, to show these additional economic features.

Structure-section sheet.—This sheet exhibits the relations of the formations beneath the surface. In cliffs, canyons, shafts, and other natural and artificial cuttings, the relations of different beds to one another may be seen. Any cutting which exhibits those relations is called a *section*, and the same term is applied to a diagram representing the relations. The arrangement of rocks in the earth is the earth's *structure*, and a section exhibiting this arrangement is called a *structure section*.

The geologist is not limited, however, to the natural and artificial cuttings for his information concerning the earth's structure. Knowing the manner of formation of rocks, and having traced out the relations among the beds on the surface, he can infer their relative positions after they pass beneath the surface, and can draw sections representing the structure of the earth to a considerable depth. Such a section exhibits what would be seen in the side of a cutting many miles long and several thousand feet deep. This is illustrated in the following figure:




Fig. 2.—Sketch showing a vertical section at the front and a landscape beyond.

The figure represents a landscape which is cut off sharply in the foreground on a vertical plane, so as to show the underground relations of the rocks. The kinds of rock are indicated by appropriate symbols of lines, dots, and dashes. These symbols admit of much variation, but the following are generally used in sections to represent the commoner kinds of rock:




Fig. 3.—Symbols used in sections to represent different kinds of rocks.

The plateau in fig. 2 presents toward the lower land an escarpment, or front, which is made up of sandstones, forming the cliffs, and shales, constituting the slopes, as shown at the extreme left of the section. The broad belt of lower land is traversed by several ridges, which are seen in the section to correspond to the outcrops of a bed of sandstone that rises to the surface. The upturned edges of this bed form the ridges, and the intermediate valleys follow the outcrops of limestone and calcareous shale.

Where the edges of the strata appear at the surface their thickness can be measured and the angles at which they dip below the surface can be observed. Thus their positions underground can be inferred. The direction that the intersection of a bed with a horizontal plane will take is called the *strike*. The inclination of the bed to the horizontal plane, measured at right angles to the strike, is called the *dip*.

Strata are frequently curved in troughs and arches, such as are seen in fig. 2. The arches are called *anticlines* and the troughs *synclines*. But the sandstones, shales, and limestones were deposited beneath the sea in nearly flat sheets; that they are now bent and folded is proof that forces have from time to time caused the earth's surface to wrinkle along certain zones. In places the strata are broken across and the parts have slipped past each other. Such breaks are termed *faults*. Two kinds of faults are shown in fig. 4.

On the right of the sketch, fig. 2, the section is composed of schists which are traversed by masses of igneous rock. The schists are much contorted and their arrangement underground can not be




Fig. 4.—Ideal sections of strata, showing (a) normal faults and (b) a thrust fault.

inferred. Hence that portion of the section delineates what is probably true but is not known by observation or well-founded inference.

The section in fig. 2 shows three sets of formations, distinguished by their underground relations. The uppermost of these, seen at the left of the section, is a set of sandstones and shales, which lie in a horizontal position. These sedimentary strata are now high above the sea, forming a plateau, and their change of elevation shows that a portion of the earth's mass has been raised from a lower to a higher level. The strata of this set are parallel, a relation which is called *conformable*.

The second set of formations consists of strata which form arches and troughs. These strata were once continuous, but the crests of the arches have been removed by degradation. The beds, like those of the first set, are conformable.

The horizontal strata of the plateau rest upon the upturned, eroded edges of the beds of the second set at the left of the section. The overlying deposits are, from their positions, evidently younger than the underlying formations, and the bending and degradation of the older strata must have occurred between the deposition of the older beds and the accumulation of the younger. When younger rocks thus rest upon an eroded surface of older rocks the relation between the two is an *unconformable*, and their surface of contact is an *unconformity*.

The third set of formations consists of crystalline schists and igneous rocks. At some period of their history the schists were plicated by pressure and traversed by eruptions of molten rock. But the pressure and intrusion of igneous rocks have not affected the overlying strata of the second set. Thus it is evident that a considerable interval elapsed between the formation of the schists and the beginning of deposition of the strata of the second set. During this interval the schists suffered metamorphism; they were the scene of eruptive activity; and they were deeply eroded. The contact between the second and third sets is another unconformity; it marks a time interval between two periods of rock formation.

The section and landscape in fig. 2 are ideal, but they illustrate relations which actually occur. The sections on the structure-section sheet are related to the maps as the section in the figure is related to the landscape. The profile of the surface in the section corresponds to the actual slopes of the ground along the section line, and the depth from the surface of any mineral-producing or water-bearing stratum which appears in the section may be measured by using the scale of the map.

Columnar section sheet.—This sheet contains a concise description of the sedimentary formations which occur in the quadrangle. It presents a summary of the facts relating to the character of the rocks, the thickness of the formations, and the order of accumulation of successive deposits.

The rocks are briefly described, and their characters are indicated in the columnar diagram. The thicknesses of formations are given in figures which state the least and greatest measurements, and the average thickness of each is shown in the column, which is drawn to a scale—usually 1000 feet to 1 inch. The order of accumulation of the sediments is shown in the columnar arrangement—the oldest formation at the bottom, the youngest at the top.

The intervals of time which correspond to events of uplift and degradation and constitute interruptions of deposition are indicated graphically and by the word "unconformity."

CHARLES D. WALCOTT,
Director.

Revised January, 1904.

DESCRIPTION OF THE JOPLIN DISTRICT.¹

By W. S. Tangier Smith and C. E. Siebenthal.²

INTRODUCTION.

The Joplin district lies between 94° 15' and 94° 45' west longitude and 37° and 37° 15' north latitude, thus including two 15-minute quadrangles. The average length from east to west is about 27.6 miles and the width from north to south is 17.2 miles. The total area is about 476 square miles.

The Joplin district was surveyed during the field season of 1901 and parts of the field seasons of 1902 and 1903. Some time has unavoidably elapsed between the field investigation and the preparation of the folio, and in the meantime many minor changes have, of course, taken place in the district. These, however, are of such a nature that the omission of any mention of them will not, it is believed, materially affect the value of the text.

Unless otherwise noted, the determinations of fossils and the correlation of formations are given on the authority of Dr. George H. Girty.

Literature.—The number of published papers dealing with the Ozark region, especially with the ore deposits of the Joplin district, is considerable. Bibliographies, including the earlier works relating to the region, are given in the following papers, that of Keyes bringing the record down to 1896:

SAMPSON, F. A., Bibliography of the geology of Missouri: Bull. Missouri Geol. Survey, No. 2, 1890.
WINSLOW, A., Lead and zinc deposits: Missouri Geol. Survey, vol. 7, 1894, pp. 743-753.
KEYES, C. R., Bibliography of Missouri geology: Missouri Geol. Survey, vol. 10, 1896, pp. 221-232.

The most important publications bearing on the Joplin district are the following:

SCHMIDT, A., and LEONHARD, A., Lead and zinc regions of southwestern Missouri: Missouri Geol. Survey, vol. 1, 1874, pp. 381-392.
WINSLOW, A., Lead and zinc deposits: Missouri Geol. Survey, vol. 8, 7, 1894.
JENNEY, W. P., Lead and zinc deposits of the Mississippi Valley: Trans. Am. Inst. Min. Eng., vol. 22, 1894, pp. 171-225, 623-648.
BAIN, H. F., VAN HISE, C. R., and ADAMS, G. I., Preliminary report on the lead and zinc deposits of the Ozark region: Twenty-second Ann. Rept. U. S. Geol. Survey, pt. 2, 1901, pp. 33-228.
SMITH, W. S., and TANGIER, Lead and zinc deposits of the Joplin district, Missouri-Kansas: Bull. U. S. Geol. Survey No. 218, 1903, pp. 197-204.
BUCKLEY, E. R., and BUEHLER, H. A., Geology of the Granby area: Missouri Bureau Geology and Mines, 2d ser., vol. 4, 1906.
HAWORTH, ERASMUS, CRANE, W. R., ROGERS, A. F., and OTHERS, Special report on lead and zinc: Univ. Geol. Survey Kansas, vol. 7, 1904.

General relationships.—The lead and zinc deposits of the Mississippi Valley occur in three groups—(1) those of the Ozark region; (2) those of the upper Mississippi Valley, including the deposits of southern Wisconsin, eastern Iowa, and northern Illinois; and (3) those of outlying districts, including chiefly southern Arkansas and the Kentucky-Illinois districts. The deposits of the Ozark region are by far the most important.

The Ozark region contains four districts: the southeastern Missouri district, essentially lead producing; the central Missouri district, characterized by small ore bodies yielding both lead and zinc; the Missouri-Kansas or southwestern Missouri district, the main zinc-producing area; and the northern Arkansas district, producing chiefly zinc, with a minor proportion of lead. These four districts are arbitrarily chosen so as to include the chief ore-producing centers. If all the scattered deposits known in the region were included a continuous zone would be outlined, beginning with the southeastern Missouri district on the east, circling

¹The Joplin district folio, which was published in 1907, is reprinted without revision. The district has not been reexamined in order to bring the statement of mining development up to date.

²The survey of the Joplin district was made by Messrs. Smith and Siebenthal jointly. Mr. Smith, who was in charge of the work, devoted his studies chiefly to the mines and ore deposits. The description of the geology, geology, and nonmetalliciferous deposits is largely by Mr. Siebenthal. The description of the ore deposits is by Mr. Smith; his sections on the forms and relations of ore bodies, however, have been rewritten by Mr. Siebenthal in the light of later studies, particularly with reference to the interpretation of structural features. All petrographic work was done by Mr. Smith.

to the northwest, then to the south not far from the margin of the Ozark uplift, and finally eastward along the northern slopes of the Boston Mountains.

The ore deposits of the several districts of the Mississippi Valley are distinguished from one another by certain differences both in the form and structure of the ore bodies and in the mode of occurrence and character of the ore. The Missouri-Kansas district is characterized by the absence of well-defined fissures and by the common occurrence of the ores in large bodies of slight horizontal extent known as "runs," or in comparatively thin tabular bodies of great horizontal extent, known as "blanket veins" or "sheet ground." The ores consist of lead and zinc sulphides, occurring in the runs mainly as a cement to Mississippian chert breccias and in the sheet ground mainly as horizontal layers between only slightly disturbed beds of Mississippian chert.

The Joplin subdistrict as considered herein comprises a part of the Missouri-Kansas district lying near its western margin, partly in Kansas and partly in Missouri. In local usage the name is somewhat loosely applied, generally to about the area here indicated, but at times to a much more extensive region, in some cases even including the whole of the southwestern Missouri and northern Arkansas districts and portions of the central Missouri district.

Topography of the Ozark uplift.—The Joplin district is an integral part of the Ozark region, which has been described in an earlier report (Twenty-second Ann. Rept. U. S. Geol. Survey, pt. 2, 1901, pp. 69-75), so that only an outline of its physical features will be presented here, the reader being referred to that report for details.

Topographically, the Ozark uplift is a low asymmetric dome with rudely elliptical outline lying in southern Missouri and northern Arkansas. It is approximately bounded on the north and northeast by Missouri and Mississippi rivers, on the west by Spring and Neosho rivers, on the south by Arkansas River, and on the southeast by Black River and some of its tributaries. To the north and west it merges into the Prairie Plains; to the east and southeast it passes into the low-lying Gulf Plains; on the south it is separated from the Ouachita Mountains by the Arkansas River valley. The uplift as a whole is a table-land bounded by

the harder formations, with scattered outliers rising above the general level. Over much of the region, especially on the northern and western slopes, these broad uplands are cut by moderately shallow, open valleys, whose depth does not exceed a few hundred feet. Toward the south and east the dissection is greater, the forms are more rugged, and the plateau character is somewhat less evident, especially near the east end of the Boston Mountains. The drainage of the uplift is radial, there being many important streams which flow outward to the marginal rivers. Of these Spring River is the largest within the Joplin district.

While it is probable that the drainage of the Ozark region is in part consequent, the streams following the normal surface and marginal slopes of the dome, it seems to be in part also subsequent, many of the stream courses having been determined rather by the underlying rock structure and texture. Except along the short southern slope of the uplift, all the important streams of the region have developed meanders.

Geology of the Ozark uplift.—The Ozark uplift is structurally, as well as topographically, a broad, low dome of elliptical outline. The structural crest lies north of the Boston Mountains, which form the topographic culmination. This crest follows a curved line from the St. Francis Mountains to Cedar Gap. It is not marked by any conspicuous topographic feature, but falls within the great central upland. From the crest the rocks slope gently outward, with dips slightly greater than the topographic slopes. Near the edge of the uplift the dip commonly increases, and along the southern border in particular there is a pronounced monoclinical structure.

The rocks of the region are in the main of sedimentary origin. In the St. Francis Mountains certain pre-Cambrian peaks of granite and porphyry project from beneath the later beds, while in Camden County there is a dike of granitic granite thought to be of post-Carboniferous age (Missouri Geol. Survey, vol. 7, 1894, pp. 432-434; Water-Sup. and Irr. Paper No. 110, U. S. Geol. Survey, 1905, pp. 113-125). The Cambrian and Cambro-Ordovician dolomites, limestones, and sandstones which underlie the central portion of the Ozarks are distributed in concentric bands around the St. Francis Mountains; these mountains, how-

ever, lie in an eccentric position as regards the whole uplift, the result being that each formation occupies a belt that is narrow east of the mountains and wider on the west.

In Camden County, where the granite dike already mentioned occurs, there is a second but subordinate dome around which circle the formations of the Cambrian and lower Ordovician.

Outside of the area of these older sediments Devonian, Mississippian, and Pennsylvanian rocks outcrop in turn, except on the southeast, where they are covered by the overlapping Tertiary of the

upland lying somewhat below the crest of the Boston Mountains.

The erosion forms throughout most of this area are those characteristic of an early mature topography in a region of nearly flat-bedded rocks of varying degrees of hardness, in which the dip differs but slightly from the general surface slope; and the entire northern slope of the uplift consists of a succession of broad plateaus separated by more or less ragged escarpments which mark the margins of

the harder formations, with scattered outliers rising above the general level. Over much of the region, especially on the northern and western slopes, these broad uplands are cut by moderately shallow, open valleys, whose depth does not exceed a few hundred feet. Toward the south and east the dissection is greater, the forms are more rugged, and the plateau character is somewhat less evident, especially near the east end of the Boston Mountains. The drainage of the uplift is radial, there being many important streams which flow outward to the marginal rivers. Of these Spring River is the largest within the Joplin district.

While it is probable that the drainage of the Ozark region is in part consequent, the streams following the normal surface and marginal slopes of the dome, it seems to be in part also subsequent, many of the stream courses having been determined rather by the underlying rock structure and texture. Except along the short southern slope of the uplift, all the important streams of the region have developed meanders.

Geology of the Ozark uplift.—The Ozark uplift is structurally, as well as topographically, a broad, low dome of elliptical outline. The structural crest lies north of the Boston Mountains, which form the topographic culmination. This crest follows a curved line from the St. Francis Mountains to Cedar Gap. It is not marked by any conspicuous topographic feature, but falls within the great central upland. From the crest the rocks slope gently outward, with dips slightly greater than the topographic slopes. Near the edge of the uplift the dip commonly increases, and along the southern border in particular there is a pronounced monoclinical structure.

The rocks of the region are in the main of sedimentary origin. In the St. Francis Mountains certain pre-Cambrian peaks of granite and porphyry project from beneath the later beds, while in Camden County there is a dike of granitic granite thought to be of post-Carboniferous age (Missouri Geol. Survey, vol. 7, 1894, pp. 432-434; Water-Sup. and Irr. Paper No. 110, U. S. Geol. Survey, 1905, pp. 113-125). The Cambrian and Cambro-Ordovician dolomites, limestones, and sandstones which underlie the central portion of the Ozarks are distributed in concentric bands around the St. Francis Mountains; these mountains, how-

ever, lie in an eccentric position as regards the whole uplift, the result being that each formation occupies a belt that is narrow east of the mountains and wider on the west.

In Camden County, where the granite dike already mentioned occurs, there is a second but subordinate dome around which circle the formations of the Cambrian and lower Ordovician.

Outside of the area of these older sediments Devonian, Mississippian, and Pennsylvanian rocks outcrop in turn, except on the southeast, where they are covered by the overlapping Tertiary of the

TOPOGRAPHY.

GENERAL STATEMENT.

The Joplin district lies on the northwestern slope of the Ozark uplift, not far from its margin, and is included in the more plateau-like, less dissected portion of the region. Throughout the district the physical features are closely related to the geology. The rocks are wholly sedimentary and those outcropping at the surface belong (with few and relatively unimportant exceptions) to two formations—the Boone and the Cherokee. The Boone, composed of limestone and interbedded chert, is the more resistant of the two and covers the larger part of the district. The Cherokee, stratigraphically higher than the Boone, consists of shale and sandstone and is confined largely to the western and northwestern portions, though small patches are abundant all over the district and there are a few larger isolated areas. These scattered outcrops are outliers of the main area on the west, having been left as remnants in the erosion of this formation, which formerly covered the entire district. Both these formations have been important factors in the development of the topography of the district.

UPLAND PLAINS.

The upland surface of the district is on the whole almost flat, with a very low general slope to the northwest, having an average fall of 5 feet to the mile between the southeast and northwest corners. Seen from favorable points the surface presents nearly everywhere an apparently horizontal, unbroken sky line. The general slope of the upland is due to the structural slope of the underlying rocks, modified by orographic movement and peneplaned.

The height of the upland surface of the district above sea level is not great, and the difference in elevation between the highest and lowest points in the district amounts to little more than 400 feet. The highest point of the upland, about 1210 feet above tide, is near the southeast corner, while the lowest point, about 775 feet in altitude, is on Spring River where it leaves the district near the southwest corner.

The upland surface is broken here and there by low hills rising somewhat above the general level. Those about a mile south of Webb City, and the broad, flat-topped, meslike Timbered Hills southwest of Crestline, are excellent examples. They are hills of circumscription and remnants of the once continuous Cherokee formation.

DRAINAGE.

STREAMS.

The important streams of the district are Spring River and its two main tributaries, Center Creek and Shoal Creek. These streams are perennial and carry



FIG. 1.—General south-north section through Springfield and Sedalia. Showing the flat-topped dome of Cambrian and Ordovician limestones overlain by thin Devonian shale and Carboniferous sediments.

long, low northern and western slopes, with a surface inclination generally imperceptible to the eye, and for much of its extent by an abrupt southern slope, facing the open valley of Arkansas River (figs. 1 and 2). Near and parallel to its southern margin the uplift culminates topographically in the Boston Mountains, a long, narrow plateau rising to an elevation of 2000 feet. The plains which surround the uplift have a general altitude of 500 to 750 feet. The central portion of the Ozarks is an upland lying somewhat below the crest of the Boston Mountains.



FIG. 2.—General west-east section through Joplin and the St. Francis Mountains. Showing the flat-topped dome of Cambrian and Ordovician limestones and pre-Cambrian crystalline rocks overlain by thin Devonian shale and Carboniferous sediments.

The erosion forms throughout most of this area are those characteristic of an early mature topography in a region of nearly flat-bedded rocks of varying degrees of hardness, in which the dip differs but slightly from the general surface slope; and the entire northern slope of the uplift consists of a succession of broad plateaus separated by more or less ragged escarpments which mark the margins of

Gulf Plains, and with the further exception that, owing to unconformity around most of the uplift, the continuity of the Devonian outcrop is here and there interrupted.

The Mississippian (Lower Carboniferous) deposits outcrop only in a narrow belt, except on the southwest, where they form the surface rock over a great area approximately 80 miles wide from northwest

abundant water. They rise outside the district, but in their course across it they receive important additions.

Nearly all the numerous valleys tributary to these three principal streams are comparatively short, and even the longer ones, with a few exceptions, do not exceed 4 or 5 miles in length. Most of these valleys carry water only after rains, but there are a number of perennial spring-fed tributaries. In addition to these, a few valleys, normally dry except after rains, carry small streams of water pumped from the mines.

The general direction of the more important streams of the district east of the Kansas-Missouri State line, including Turkey and Short creeks, is a little north of west; the numerous tributaries to these streams flow in the main at right angles to them. The direction of the major drainage lines, except that part of Spring River which flows southwestward, has doubtless been determined by the configuration of the general surface slope of this part of the Ozark uplift. In no case, with the possible exception of Shoal Creek, do these courses appear to have been determined by lines of weakness in the underlying rocks, such as folds; nor have they apparently been influenced by the many small shale areas found over the district. Shoal Creek, as can be seen by an inspection of the structure contours on the economic geology map, takes a course corresponding very closely to a structural depression. Inasmuch as other structural features fully as pronounced seem to have exerted no control on the drainage, it may well be assumed that the alignment of Shoal Creek is simply coincidental.

While the general westerly course of that part of Spring River lying in Kansas has doubtless been inherited from earlier conditions in which it was determined by the original slopes of the margin of the Ozark uplift, its immediate position appears to have been fixed by the line of contact between the Cherokee and Boone formations. Where a soft formation overlies a harder one, both dipping at a moderate angle, the general tendency of a stream flowing parallel to the strike of the rocks, unless checked in some way, is to follow constantly the contact between the two formations, working down the slope as the edge of the softer rocks is gradually eroded. This, as pointed out by Adams (Trans. Kansas Acad. Sci., vol. 16, 1889, p. 56), appears to have been the case with Spring River. It has followed the edge of the Cherokee shale to its present position, where comparatively recent and more active cutting has caused it to become entrenched in the harder Boone formation close to the line of contact.

The minor drainage lines have for the most part been determined by the physiographic factors commonly effective in such a region as this. In some instances, however, according to Bain (Twenty-second Ann. Rept. U. S. Geol. Survey, pt. 2, 1902, p. 221), the stream courses follow zones of fracturing, which may or may not be accompanied by faulted masses of Cherokee shale lying below the general level of the upland. It may be that the softness of the shale, rather than the fracturing, is the factor determining the direction of the drainage in all cases where this rock appears; and where the stream follows fracturing not accompanied by shale, that rock may have been present originally and aided in the location of the stream, which subsequently removed it through erosion. Shale areas not connected with zones of fracturing, but filling pre-Cherokee erosion hollows, may also have been effective in directing the flow of some streams. Examples of valleys following bands of shale are found east and west of Neck; northeast and southwest of Carterville; in Joplin Creek and its southerly fork, east of Joplin; in the vicinity of Klondike; and west of Galena.

VALLEYS.

The dissection of the district is moderate. Important streams are not numerous, and those present are separated by broad, flat-topped divides. In the area underlain by the Boone formation these divides are cut by many small, shallow, and open valleys formed by the headwaters of tributary streams. In their lower courses these tributary valleys are deeper and less open, and they are here and there bordered by low cliffs of limestone and chert. Such tributary valleys lie along all the more important stream courses of the district.

They are deeper adjacent to the deeper main valleys, and are an especially pronounced feature along Shoal Creek, where, near their mouths, they range in depth from about 80 to about 150 feet. In the Cherokee formation the dissection is much less than in the Boone, and the valleys are on the whole much shallower, more open, and less numerous. Even the larger valleys in this formation, as those of Shawnee and Cow creeks, are open and very shallow.

The largest stream valleys of the district are comparatively broad and flat bottomed, and here and there, as along Shoal Creek and the lower reaches of Spring River, are rudely terraced. While these valley bottoms are generally covered with alluvium, those of Shoal Creek and some of its tributaries are locally floored with bare rock, the upper surface of the Grand Falls chert.

At Baxter Springs, just before leaving the district, Spring River enters a narrower, deeper valley which seems younger than the valley above that point.

The valley slopes consisting of the Cherokee shale and sandstone are very gentle, but those formed of the Boone rocks are more abrupt. Cliffs are common in the Boone valleys, especially where the walls are being undercut by the stream meanders. Since the upland surface rises toward the south, while the general drainage of the district is to the southwest, the larger valleys are deeper in their southern parts. The valley of Shoal Creek is the deepest in the district. West of Grand Falls the bluff hills bordering this stream reach a maximum height of about 150 feet.

Terraces.—The terraces of the district are of two varieties—alluvial flats and rock shelves. The former are confined to the valley of Spring River; the latter are found mainly along Shoal Creek.

The best example of the alluvial terrace lies on the east side of Spring River, extending southward from the village of Lowell for 3 miles, nearly to the limit of the area mapped. It has an elevation of about 15 feet above the stream and a width of half a mile. The terrace front descends abruptly to the river bottom, while the surface rises very gently to the bordering hills. At the south end it loses its terrace character and becomes an alluvial slope similar to those on the south side of Shoal Creek, described in the next section. Another terrace lies on the same side of Spring River just northeast of Lowell and similarly develops into an alluvial slope at its southeast end. A third well-developed terrace, about 160 acres in extent, lies east of Spring River and south of Short Creek.

In many places on each side of Spring River from Baxter Springs to Waco the upland plain slopes so gently toward the river that it is quite impossible to distinguish the limits of the present flood plain except by noting the height of the high water in the flooded stream. This is true in the vicinity of Vark and likewise west and north of Smithfield. On the flat west and southwest of the old Boston Mills the terrace or second bottom lies at an elevation of 10 feet or so above the alluvial flood plain and is limited on the north by another flat or third bottom 15 to 30 feet above it, corresponding to the lower country about and east of Eldon.

No second bottom is distinguishable in the great bend south of Messer post-office nor in the bend west and north of Smithfield, but the alluvial plain, as already noted, passes gradually into the upland. These conditions continue upstream to a point east of Waco, where a well-developed terrace is exhibited on each side of the river. The absence of the terraces in this interval is due to the fact that they have been removed by the erosion that has lowered the river to its present level.

Along Shoal Creek in a number of places the creek bottom is bordered by a terrace 20 to 40 feet or more in height, the front of which is a sheer wall of massive chert. These rock-shelf terraces are not true stream terraces, inasmuch as they do not represent graded sections of the stream valley which have been abandoned by the deeper cutting of the stream; moreover, they do not lie at any uniform elevation above the present grade of Shoal Creek. On the contrary, they are but gentle swells in the more resistant Grand Falls chert, which have been etched into relief and cut through by the stream in the process of lowering its bed. At Grand Falls, the type locality, Shoal Creek is even

now attacking one of the more resistant of these bosses of chert. Good examples of these rock shelves are found about Grand Falls and along the stream as far as Gregg's bridge, 2 miles below; also from Redings Mill to a point below the mouth of Silver Creek. These shelves constitute the principal exposures of the Grand Falls chert and are discussed further in the description of that member.

Alluvial slopes.—On the south side of Shoal Creek, about 2 miles southeast of Lowell, the land slopes gently from the creek bank to the foot of the hill a quarter of a mile south, rising 35 to 40 feet in that distance. Just at this point a valley about 300 yards long debouches from the south, forming a well-marked little alluvial fan. This suggests that the slopes are aggradation plains built up of outwash from the hills by the coalescing of alluvial fans, and are thus alluvial slopes. This suggestion is borne out by the fact, shown by well sections, that the slope down to the level of the stream is made up of gravel and wash material.

In the description of the terraces on Spring River reference has already been made to some of these slopes. A characteristic one lies on the south side of Shoal Creek due south of Galena, and, as can be noted on the map, others occur on each side of the creek. By far the largest and best developed commences at Gregg's bridge, 2 miles west of Grand Falls, and stretches westward along the south side of Shoal Creek valley a distance of 2 miles, to a point within a mile of the State line. This slope is over half a mile in width. The edge adjacent to the hills is 55 to 60 feet higher than the banks of the creek. A little west of the middle of the slope a valley perhaps a quarter of a mile in length, at the foot of which is a very plain alluvial fan, comes down from the southern hills.

Various wells at the southern margin of this slope penetrate from 30 to 40 feet into it, the material without exception being rock fragments, gravel, sand, and clay—typical wash material. This shows that the valley throughout its width has been eroded to about the level of the present stream, and has been refilled in part by inwash from the sides. Only the shorter side valleys exhibit alluvial fans, the reason being that the larger tributaries, having cut nearer to grade, experience no great change in slope on reaching the main valley, and therefore drop no great amount of material at that point.

DESCRIPTIVE GEOLOGY.

STRATIGRAPHY.

The rocks outcropping in the Joplin district comprise, in ascending order, the Boone, Carterville, and Cherokee formations, belonging to the Carboniferous system, and various unconsolidated gravels, soils, and alluvium formed at much later periods. Each of the consolidated formations is limited above and below by unconformities, that between the Boone and the Cherokee being particularly of the kind herein described as "solution unconformities."

UNDERLYING FORMATIONS NOT OUTCROPPING IN THE DISTRICT.

In addition to the outcropping rocks, various older formations are penetrated in shafts and drill holes. The best section of these older beds is furnished by the Harrington well at Carthage, a record of which is given by Winslow (Missouri Geol. Survey, vol. 7, 1894, p. 405). This hole, 2005 feet in depth, penetrates 1750 feet of stratified deposits and goes 255 feet into the crystalline rocks. On the basis of stratigraphic studies farther east, Ulrich correlates the section afforded by the well as follows:

Correlation of Harrington well section, Carthage.

	Feet.
MISSISSIPPIAN (LOWER CARBONIFEROUS):	
Burlington and Keokuk limestones.....	355
Chouteau limestone.....	15
Hannibal shale.....	15
Louisiana limestone.....	65
ORDOVICIAN:	
Joseph limestone.....	80
St. Peter sandstone.....	50
Cambro-Ordovician and Cambrian limestones and sandstones.....	1185
Crystalline rocks (porphyry).....	205
	2005

A study of all the available records of other deep wells in the district shows the practical impossibility of carrying this correlation throughout the

district, owing to the imperfection of the records and to the variability of the formations themselves.

The Devonian black shale (Chattanooga formation) has not been identified in the district. The Mystic shaft in the bottom of Gordon Hollow, a quarter of a mile above the mouth, starting at the base of the Grand Falls chert, gives the following section:

Section of Mystic shaft, Gordon Hollow.

	Feet.
Gravel and clay.....	30
Chert and limestone.....	100
Shale.....	35
	165

The shale is a greenish-blue, arenaceous, friable shale, carrying spindles and irregular lenses of soft white chert, now and then with flinty centers. Weathered out from the shale and lying on the surface of the dump are numerous crystals of calcite, one-half inch or so in size, together with more or less sphaerite and galena in crystals. Judging from its stratigraphic position and its appearance, this shale is taken to be the representative in this region of the Hannibal shale.

Some of the deep wells penetrate shale below the Grand Falls chert, but its variable thickness when so found, its variable distance below the chert, and its utter absence in many of the wells, especially in the Freeman Foundry well, complete samples of which were carefully examined, together with the local occurrence of Cherokee shale in cave galleries at great depths, all conspire to throw much doubt on any correlation of such shale, as well as on the correlation of the limestones and cherts immediately above and below it. Shale at about the same horizon in some other wells, notably the Harrington well and the Missouri Zinc Fields well at Webb City, and in a drill hole at the Lincoln mine near Klondike may with some certainty be classed as Hannibal. The shale in the well of the United Zinc Companies at Chitwood may with less probability be called Hannibal, and it is almost certain that the shale in the well of the Boston-Duenweg Company at Duenweg, separated from the broken ground above by only 38 feet of limestone and chert, is not Kinderhook but Cherokee shale occupying a cave gallery, as indeed may be also the shale in the well of the United Zinc Companies. In the correlation of the Harrington well section, already given, the Devonian shale does not appear; but that formation outcrops in irregular patches 30 or 40 miles to the south and east. In the light of present knowledge, and particularly in the absence of lithologic description of the shales struck in the various wells, it is not possible to affirm that these shales are Hannibal and not Devonian. In any event, whether they be correlated with the Hannibal or the Chattanooga shale, the formation is evidently not persistent over the district. Beneath this questionable horizon the great body of magnesian limestones, sandstones, and cherts is undoubtedly of Cambrian and Ordovician age.

ROCKS THAT OUTCROP.

CARBONIFEROUS ROCKS. BOONE FORMATION.

Character.—The Boone formation is limited above by the unconformable Cherokee shale and below by the nonpersistent Hannibal shale. It consists of limestones, cherts, and dolomites. The proportion of limestone is rather larger than that of chert, and the dolomite constitutes a very small part of the formation as a whole. Two subdivisions of the formation are held to have sufficient importance areally and lithologically to justify their classification as members. They are the Grand Falls chert and the Short Creek oolite, the latter lying 100 feet above the former. These horizons divide the Boone into three limestone-chert terranes. The uppermost division is prominent in the vicinity of Carthage, where the lower part of it is extensively quarried. The middle division is also largely made up of crystalline limestone, which is quarried for building stone and burned for lime at the Joplin lime kiln, at the mouth of Opossum Hollow. The lowest division, below the Grand Falls chert, is made up of chert and fine-grained dark-blue flaggy limestone. It outcrops in but few places, one of which is in the bluffs of Shoal Creek southwest of Grand Falls, where the bottom of the chert rises above drainage. The Mystic shaft, in the bottom

of Gordon Hollow near its mouth, went through 100 feet of this division and into the Hannibal shale. This limestone crops out in the bank of Shoal Creek near the Kansas-Missouri line and is struck in shafts 1½ miles south of east, near the mouth of Lee Hollow, where it is reported to be 130 feet thick.

At these places there is usually an intersecting network of veins of crystallized dolomite or pink spar. Its position in the section is inconstant, being here above, there below the Short Creek oolite. Owing to the nature of the outcrop, the dolomite can not be traced laterally for long distances, and in all probability it does

in most of the mines and at all depths which have been reached by mining. Chert approaching cotton flint has been noted in the cuttings from the Freeman Foundry well, Joplin, at intervals to a depth of over 300 feet.

Many of the cherts, both fresh and weathered, are somewhat calcareous, the calcite, in minute or microscopic grains, being either distributed through the rock or concentrated in small areas, many of which are of irregular shape, which mottle or streak the light gray of the chert with darker gray. In places the calcite is limited largely to the megascopic fossils of the rock, and doubtless to the leaching of the calcite in these fossiliferous rocks some of the cavities in such chert are due. Some of the calcite found in the chert is undoubtedly original, but in other instances it is as unquestionably secondary, having either replaced the chert metasomatically or filled cavities formed in the chert by solution.

Microscopically the cherts of the Joplin district consist mainly of a microgranular to cryptocrystalline aggregate of silica in the form of both quartz and chalcedony. The grain is fairly uniform in some specimens, both fresh and weathered, but as a rule it shows more or less variation, which appears to be due largely, if not entirely, to recrystallization. To a considerable extent such recrystallization is selective, occurring in the fossils or in certain parts of the fossils contained in the rock. It is often found also bordering the minute cavities common in many of the more or less weathered cherts. The recrystallized portion varies from a rock with the grain of the normal chert to a fine-granular quartz aggregate. These recrystallized portions, both coarser and finer, consist, like the original rock, of quartz and chalcedony, the coarser grained parts, however, being more commonly made up of quartz alone, though in some instances they consist almost wholly of chalcedony aggregates. Many of the cherts also contain minute opaque grains, for the most part iron sulphide.

As shown both by the microscope and by chemical analyses, the greatly weathered cotton flints do not differ essentially in chemical composition from the unaltered rock, the differences between the two being chiefly physical. Cotton flint, as seen under the microscope, contains many minute cavities and shows on the whole a greater amount of recrystallization. Some of the cotton flints are uniformly granular, though most of them show the variation in grain which usually characterizes recrystallization. Some of the slides of cotton flint examined were almost free from cavities and not noticeably different from unaltered chert. Some of cavities are molds of fossils, but many of them are due to general solution of the rock. In most of the cotton flint examined no calcite was observed, and where found it occurred only in scattered grains.

Distribution and thickness.—The Boone is the surface formation over much the larger part of the district, occupying, with the exception of a few large and numerous very small outliers of Cherokee, all the country east of Spring River and some territory west of the river and adjacent to it. The best exposures are found along Shoal Creek, where some of them reach into the portion of the formation lying below the Grand Falls member. Exposures of the rocks lying above the Grand Falls chert are found along Turkey Creek, Center Creek, Spring River, and the various tributaries of those streams.

The thickness of the Boone is difficult to estimate accurately. The base of the formation is usually taken at the first shale or shaly limestone which occurs below the Grand Falls member. But many of the drill records fail to note any shale whatever. This may be explained in several ways. The shale may really be absent in these localities, since it is known to have an intermittent outcrop farther east. On the other hand, it may be so calcareous as to be confused with limestone, or the driller may have failed to recognize it owing to its thinness. In the deep well on the Missouri Zinc Fields land at Webb City it is but 12 inches thick. When it is not especially watched for, such a thickness might very well be overlooked in working with a churn drill at a depth of 350 feet, the more so when surface shale has been previously penetrated.

Estimated on this basis, the Boone formation in the Carthage well is 355 feet thick, and this is the maximum thickness of the formation penetrated

anywhere in this district. The greatest known thickness of the limestone and chert series overlying the Grand Falls chert member and including the Short Creek oolite is 208 feet, found in a well on the land of E. B. Allen, 2 miles southeast of Joplin, in the NE¼ NW¼ sec. 13, T. 27 N., R. 33 W. At the Freeman Foundry well in Joplin the thickness is 207 feet. These figures may be taken to represent the maximum thickness of the formation above the Grand Falls chert now remaining in the district. The normal thickness of the Grand Falls chert ranges from 50 to 60 feet, but in some districts it is recorded as low as 15 feet and in others as high as 120 feet. The excessive thickness is to be explained by the fact that, owing to the preponderance of chert in the beds between the Grand Falls member and the chert stratum some distance below it, the whole series has been recorded as Grand Falls. The thickness of the alternating bluish limestones and cherts which underlie the Grand Falls member is variable, ranging apparently from 25 to 150 feet.

Generalized section of Boone formation.

	Feet.
Upper limestone (containing Carthage quarry beds).....	100
Short Creek oolite limestone member.....	2 to 8
Middle limestone (containing Joplin linekin quarry beds).....	100
Grand Falls chert member.....	50 to 60
Lower limestone.....	25 to 150

Relations.—At the contact of the Boone with the Cherokee above there is a double unconformity, or rather an unconformity which is a result of two distinct cycles of erosion and solution—the post-Boone-pre-Carterville and the post-Carterville-pre-Cherokee (see columnar section fig. 3). Elsewhere the Boone outcrop has suffered in addition the Tertiary penneplation and the post-Lafayette erosion. This solution unconformity is complicated to a certain extent by minor faulting. It is discussed somewhat more in detail in the description of the Carterville and Cherokee formations.

The greatest variation in thickness and character and the greatest difficulty in correlation are found in that portion of the Boone below the Grand Falls chert. This variability is but natural in beds overlying a nonpersistent formation, the irregularities of which are probably due in part to erosion. The Hannibal shale, which presumably limits the formation below, comes to the surface nowhere in the district and, except in drill holes, is penetrated at but one place—in the Mystic shaft, a section of which has been given. The difficult correlation of shales struck in drilling below the Grand Falls member and the consequent uncertain recognition of the lower limit of the Boone have already been mentioned.

Name and correlation.—The Boone formation derives its name from its typical development in Boone County, Ark. (Ann. Rept. Geol. Survey Arkansas, 1890, vol. 1, p. 129), where it displays many of the characteristics which it exhibits in the Joplin district. Jenney (op. cit., p. 178) proposed the name Cherokee for the limestones of the district, but that term has since been generally used for the Pennsylvanian shale of the region, and such usage has the preference. Elsewhere these limestones have generally been called Burlington.

The formation as a whole belongs to the Osage group of the Mississippian series. It includes portions of the Keokuk and the Burlington, the division line coming probably below the Grand Falls chert. A fauna with Keokuk affinities is described from the Grand Falls chert in the discussion of that member. No paleontologic collections are available from the blue limestones below the Grand Falls chert, or from the Hannibal shale.

A collection obtained near Grand Falls, about 20 feet above the chert, yielded the following species:

Fossils collected near Grand Falls.

Cyclopora fungia.	Spirifer keokuk.
Fistulipora sp.	Spirifer tenuicosta.
Stenopora sp.	Syringothyris 1 sp.
Leioclema punctatum.	Spiriferina sp.
Rhombopora sp.	Eumetria marcyi.
Cystodictya cf. C. lineata.	Aviculopecten amplius.
Polypora sp.	Myalina 1 sp.
Chonetes aff. C. illinoensis.	Conocardium sp.
Productus punctatus.	Pleurotomaria sp.
	Griffithides buto.

A collection from the southwest corner of the SE¼ sec. 34, T. 28 N., R. 31 W., near the mouth of Jones Creek, 10 feet below the Short Creek




FIG. 3.—Generalized section for the Joplin district. Scale: 1 inch = 100 feet.

The stratigraphic constitution of the Boone as a whole is variable from place to place. Records of borings in the neighborhood of Grand Falls indicate much variability in the rocks below the Grand Falls chert; locally a second horizon of massive cherts occurs a short distance below that member, being separated from it by bedded limestone and chert. Where there is a local decrease in the amount of limestone in this interval there may be 100 feet or more of massive cherts, including the Grand Falls member and the cherts of the lower horizon, together with the intervening beds.

The limestone of the Boone formation is a fossiliferous granular crystalline rock which varies from fine-grained bluish limestone to light-gray, coarsely crystalline limestone or white marble, such as is quarried in the Carthage district. The individual strata range from sheets a few inches thick to massive beds 20 to 30 feet thick, with no indication of a parting on the weathered surface. These massive beds scale off under the action of frost, leaving an unbroken vertical or inclined, perhaps overhanging, face. In favorable localities the action of frost reaches to a depth of 3 feet and results in peripheral joints parallel to the exposed surface and cutting across the bedding planes. The massive beds traced horizontally are usually found to split up into smaller beds, perhaps with the intercalation of sheets and lenses of chert along the bedding planes. The occurrence of stylolites, or "crowfeet," is common.

The crystalline limestones of the Boone are nearly pure carbonate of lime and very soluble in meteoric water, and in these rocks underground drainage systems and caves have existed in the past and are found at present. Quarries in this part of the formation exhibit the usual number of crevices and enlarged joints filled with red clay and boulders of residual chert. The upper surface of the limestone is likewise covered with a mantle of red clay and residual chert. In many places where the formation at the surface originally consisted of thin-bedded limestone with thin intercalated chert beds the limestone has been carried away by solution, leaving the chert in broken and wavy but clearly perceptible beds, separated by sheets of red clay. The sinuosity of the chert beds corresponds to the irregular decomposition surface of the solid limestone beneath, and in places the chert is sharply tucked downward over a clay joint, as if sucked into it. Here and there the sides of clay seams and the walls of sink holes exhibit a fluted columnar style of weathering.

The bedded dolomite is confined to the eastern part of the Joplin area. It is buff in color, massive, finely granular, and nonfossiliferous. In appearance of outcrop and texture it very closely resembles a fine-grained sandstone. In places the bedding seems to be lost and the rock has a massive

not so extend. The indications are that it is a result of local dolomitization. As pointed out by Broadhead (Missouri Geol. Survey, vol. 1, 1874, p. 81), referring to localities several miles farther east, the alteration has usually been so complete as to entirely obliterate any traces of fossils. The eastern portion of the district is unusually free from deformation and dislocation, so that here the dolomitization does not seem to be associated with faulting.

The chert of the Boone formation occurs in beds or in lenses and nodules in the limestone. Where in beds, the chert is either intercalated with the limestone or occurs massive with more or less limestone interbedded or in lenses. Locally there is little or no interbedded limestone.

The chert varies considerably in physical characters, both in different beds and in the same bed under different conditions, such as above and below ground-water level. It is in general, when fresh, a light-gray (in places dark-gray) to nearly white, highly siliceous, compact rock, and as a rule its luster is dull, or nearly so. Locally it is light to moderately dark bluish gray in color, with a waxy luster, this type being popularly known as "live flint." Most of it is rather brittle, some beds more so than others. "Butcher-knife" and "hog-chawed" flints are extremely brittle, the former breaking, like glass, into thin or very angular fragments with sharp edges. The chert in general has a more or less pronounced conchoidal or sub-conchoidal fracture, and the fracture surfaces are usually smooth. Here and there in the mines, adjacent to some disturbance, as, for instance, along the Cornfield "bar," the chert, especially that of the sheet ground, is affected by a system of close-set, parallel fractures more or less nearly at right angles to the bedding, which in many places completely obscure the stratification and break up the heavy strata into highly inclined sheets. This type will be hereinafter referred to as "sheeted chert."

On weathering the rock often whitens, becoming dull and chalky, as in "cotton flint," sometimes being stained more or less with iron oxide to a buff color; it grows gradually less brittle and tougher and finally more or less incoherent; its fracture, instead of being conchoidal, becomes more or less rough and uneven, like that of a moderately fine-grained rock; it also becomes gradually softer, so that, as cotton flint, it may be readily grooved with a knife, though the groove still shows the color of the steel. Much of the chert, even where considerably weathered, appears to be compact, though in many instances it contains numerous and usually minute cavities due to solution. Molds of fossils are often found in partially weathered chert, and are locally abundant. The weathering of the cherts is not confined to the narrow zone above ground-water level, but more or less altered chert is found

oolite and consequently about 70 feet higher than the beds from which the collection was made near Grand Falls, afforded the following species:

<i>Fossils collected near mouth of Jones Creek.</i>	
Zaphrentis centralis?	Productus setiger.
Zaphrentis carinatus.	Productus alternatus.
Stenopora sp.	Productus mesialis.
Cystodictya lineata.	Productus wortheni.
Rhipidomella dubia.	Spirifer tenuimarginatus.
Derbya keokuk.	Syringothyris sp.
Chonetes aff. C. illinoisensis.	Composita aff. C. humilis.
Productus levicosta.	Rhynchopora aff. R. pustulosa.
Productus magnus.	Griffithides bufo.

The following species were collected at the Carthage quarries, in beds ranging from about 20 to about 60 feet above the Short Creek oolite:

<i>Fossils from Carthage quarries.</i>	
Monilopora sp.	Productus levicosta.
Fenestella tenax.	Spirifer keokuk.
Hemitypa proutana.	Spirifer rostellatus.
Polyopora halliana.	Spirifer tenuimarginatus.
Archimedes n. sp.	Spirifer neglectus.
Archimedes n. sp. aff. A. owenani.	Spirifer subserififormis.
Archimedes owenani?	Spirifer lateralis.
Meekopora 2 n. sp.	Reticularia pseudolineata.
Stenopora sp.?	Spiriferina? sp.
Cystodictya cf. C. lineata.	Composita aff. C. humilis.
Leiodoma punctatum.	Eumetria mareyi.
Rhipidomella dubia.	Dielasma gorbyi.
Chonetes aff. C. illinoisensis.	Myalina keokuk.
Productus magnus.	Aviculopecten amplus.
Productus setiger.	Aviculopecten sp.
Productus n. sp.	Fish tooth.
Productus punctatus.	

The highest recognizable horizon in the Boone is in the vicinity of Waco. At the bridge over Spring River, 2 miles east of Waco, limestone approximately 100 feet above the Short Creek oolite gave the following forms:

<i>Fossils collected near Waco.</i>	
Stenopora sp.	Syringothyris texta.
Productus magnus.	Composita aff. C. humilis.
Productus sp.	Canarozochia sp.

These faunas, representing the Boone very closely from the upper part of the Grand Falls chert to the top of the formation as exposed in the Joplin district, are essentially the same and are evidently of Keokuk age.

GRAND FALLS CHERT MEMBER.

Character.—The Grand Falls member of the Boone lies about 100 feet below the Short Creek oolite and is made up almost wholly of heavy beds of solid chert. As a rule, where there is locally a heavy development of chert in the Boone more or less limestone is interstratified with it; but in the typical development of the Grand Falls member there is a comparatively small amount of limestone in thin beds and lenses, and locally even this is replaced by jasperoid or has been removed to a greater or less extent by solution. As all occurrences of chert are more or less localized, and heavy developments especially so, it happens that the Grand Falls member can not be traced persistently to every place where, from the stratigraphic position, it would naturally be expected to appear. For instance, in the section of the west bluff of Shoal Creek half a mile west of Grand Falls, the type locality, a 15-foot bed of chert and limestone appears at the proper horizon of the Grand Falls. The chert is not massive and heavy bedded, the limestone occurs in irregular patches, and altogether, but for its position, this bed would not be taken to represent the Grand Falls chert.

The typical chert of this member is of the splintery, fresh, unaltered-appearing type known as "live" or "butcher-knife" flint. Much of it has been thoroughly crushed in place and recemented with a darker, bluish siliceous cement, the original bedding remaining practically undisturbed. Breccias of this sort occur both in the mines, especially those operating in the sheet ground, and at the surface, having been noted at several points along Shoal Creek. Where the chert has been subjected to rather sharp deformation, instead of suffering simple brecciation it loses its bedded character and becomes gnarled and knotted in structure, weathering with a very rough surface. A typical example of the transformation of the bedded brecciated chert into the massive gnarled and twisted chert is seen at Gregg's bridge, 1½ miles west of Grand Falls. Isolated outcrops of massive brecciated and gnarled cherts occur in different localities, but as there is no way of correlating them with the Grand Falls member they have not

been distinguished on the geologic map from the Boone in general.

Here and there the chert contains small, scattered, dark-colored, though still siliceous spots or areas which are rounded, oval, or irregular in shape, some of the rounded spots being half an inch or more in diameter. Most of these areas are zoned, usually with a dark-gray, nearly black, jasperoid-like nucleus with sharp boundaries, surrounded in many instances by a lighter gray, sharply marked zone that is as a rule nearly the color of the main rock. This in turn is bordered by a zone of nearly white chert, approaching cotton flint, this outer zone being commonly narrow, with an indefinite outer margin. While such spotting is not limited to any particular horizon, it appears to be most abundant in the chert of the sheet ground, forming one of its marked characteristics. As shown by the microscope, the spotting is due to recrystallization of the chert and is in places accompanied by more or less solution, both in the nucleus and in the outer zone.

The distinctive characteristics of the Grand Falls chert are, therefore, heavy bedding, "live" splintery texture, fine brecciation and cementation, and spotting.

Thickness and distribution.—With one exception, the recognized outcrops of the Grand Falls are confined to Shoal Creek or its tributaries. The exception is a small patch of the typical chert which forms the surface over a portion of the Turkey Creek bottom just north of the mouth of Leadville Hollow. Fresh exposures here show the characteristic white spotting of the blue flint, the brecciation, and the splintery texture. The thickness, as revealed by drill holes over the bottom, ranges from 30 to 35 feet, though probably some has been eroded from the surface. The identification of this chert with the Grand Falls member is corroborated by the stratigraphic position of the Short Creek oolite in the neighboring hills. The outcrop has the rough hummocky appearance usual where this formation is exposed as a rock floor.

The cause of the hummocks is not clear. They are probably due to irregular weathering, but possibly to irregularities in the silicification of the upper surface of the formation. An instance of the latter kind can be seen along the Grand Falls branch of the Missouri Pacific Railway on the west bluff of the small creek flowing southward from Joplin, about half a mile above its mouth. Here the bedded chert crops out 25 feet in thickness, but in one place it extends 15 feet higher. As there is no break in the rocks below, this seems a clear case of local silicification.

In ascending Shoal Creek the first outcrop of undoubted Grand Falls is encountered near the middle of the south side of sec. 27, 2 miles southwest of Galena. The chert forms a bluff about half a mile long, facing the east, reaching almost across the creek bottom, and varying from 5 to 23 feet in height. The chert is exposed back from the top of the bluff for 100 yards, beyond which it is covered by the alluvium of the valley. A section of the bluff at the point where Shoal Creek swings against it is as follows:

Section on Shoal Creek 2 miles southwest of Galena.

	Fe.	In.
Massive chert, showing local brecciation in horizontal sheets and recementation.	12	0
Breccia	0	3 to 6
Massive chert	3	0
Breccia	0	3 to 6
Massive chert	2	0
Weathered jasperoid	0	3
Massive chert, shattered but not brecciated	6	0
	24	0

The uppermost chert is fossiliferous at the top, yielding the brachiopod fauna hereinafter described. The first breccia band has a level definite upper limit, probably slickensided, but the bottom is very irregular, in places reaching down from 8 to 12 inches into the flint below, the whole space being filled with finely shattered chert in horizontal sheets, cemented with jasperoid. This brown jasperoid cement here and there contains cavities where crystals of blende have been leached out and other smaller cavities which may have held dolomite. Though ordinarily following the bedding planes, in places the brecciation also follows angling fracture planes. It seems reasonably certain that the brecciation along the bedding has been caused by lateral movement along these planes.

The Short Creek oolite is found in the bluff west of Harmon Branch at an elevation of 132 feet above Shoal Creek, but the lower portion of the section, at the horizon of the Grand Falls chert, is concealed, though chert fragments are strewn over the interval. Limestone and chert interstratified outcrop near the mouth of Harmon Branch, although about three-fourths of a mile to the south, up the branch, the chert is characteristically exposed with a thickness of 10 feet in sight.

Half a mile east of the State line Shoal Creek makes an abrupt bend to the north, being deflected by a bluff of Grand Falls chert which extends northward nearly to the north edge of the creek bottom. The chert follows the bluff south-eastward and outcrops in the mouths of two small branches which enter the valley from the south at this point. The chert also crops out in several places up these branches where slight anticlines bring it to the surface.

The chert is next exposed in the vicinity of Gregg's bridge, where it forms a bench or rock shelf 25 or 30 feet in height. Just west of the bridge the chert has been flexed and subjected to some torsion, the axis, along which there has been considerable brecciation and recementation, bearing N. 52° E. Two sets of joints have resulted and cut the rock up into "butcher-knife" flint. One of these systems bears N. 42°-56° W., and the other N. 10°-14° W.

The chert underlies the flat bench adjacent to Shoal Creek between Tanyard and Gordon hollows. As shown on the geologic map, it outcrops in two irregular patches in Tanyard Hollow. At the southern outcrop it rises into a 20-foot bench on the east side of the road. A little farther south a shallow shaft at the level of the road struck the underlying limestone a short distance from the surface. At the northern exposure the chert forms a rock floor with characteristic hummocky appearance. A strongly marked system of jointing bearing N. 62° W. cuts the rock at intervals of 1, 4, and 8 inches. Another less pronounced system bears N. 38° W. and, intersecting the first system, produces rhombic and wedge-shaped blocks. A very faint system bears due east and west. Though more or less covered by soil and alluvium, the chert is the underlying rock from Tanyard Hollow to Grand Falls, where the bench coincides with the creek bottom above the falls. At the falls the drop is 24 feet, and adding 10 feet of chert above the water at the top of the falls gives a thickness of 34 feet of the chert in sight at the type locality.

Northward along the railway from Grand Falls the chert maintains a uniform level for half a mile, then disappears, reappearing in a small roll at a distance of a quarter of a mile, and again rising with a dip of 3° S. at the mouth of the hollow by the waterworks. It crops out on both sides of this hollow for a distance of half a mile to the north, then spreads out over the bottom of the hollow and disappears below drainage. Just south of the mouth of the main hollow from the east, in a 6-foot railway cut, the chert affords one of the best illustrations of close jointing in the district. The principal system bears N. 30° W., and dips 70° NE., and the cracks are one-fourth inch to 3 or 4 inches apart. Less numerous but larger joints strike N. 50° W. and dip 45° to 50° SW. The intersection of these systems produces characteristic "butcher-knife" flint. Along horizontal planes the chert is finely brecciated and cemented with a reddish leached jasperoid.

Southeast of the waterworks the chert crops out along the base of the hills, rising into a bench or rock shelf just below the mouth of Silver Creek, and forming the flat valley of that creek as far up as the road crossing. Still farther southeast, in the vicinity of Redings Mill, the chert forms bluffs on each side of Shoal Creek, with a level flat above reaching back to the hills. The chert here displays the common characteristics of spotting, brecciation, and recementation with undisturbed bedding. Southwest of Redings Mill the Grand Falls member outcrops along Spring Creek for a distance of a mile and a half and the various exposures are eminently characteristic. A shaft in the NW. ¼ NW. ¼ sec. 3, T. 26 N., R. 33 W., shows 32 feet of "live," spotted Grand Falls chert, the top of which is near the bottom of the valley. A quarter of a mile farther south the chert rises 10 feet above the valley in a little roll, dipping under the surface

both to the north and to the south within 200 yards. The chert crops out along the foot of the hill southward from Redings Mill for half a mile, to a point where, dipping 2° S., it disappears beneath the alluvium of the valley.

About 1½ miles south of east of Redings Mill, in the SW. ¼ sec. 36, T. 27 N., R. 33 W., on the north side of the creek, the Grand Falls member rises in a small anticline and shows a thickness of 35 feet in the bluff just east of the house. A well at the foot of the bluff went through 18 feet of chert, making the total thickness 53 feet. The chert outcrops along the hill for a distance of half a mile, dipping beneath the overlying limestone at either end.

The Joplin anticline brings the Grand Falls chert to the surface at several points where its axis is cut by tributaries of Shoal Creek. The principal outcrop is 2 miles due east of Saginaw station, on the Kansas City Southern Railway, and others occur 1 mile west of north and 1½ miles southeast of that one. The chert rises to a height of 30 feet near the house in the middle of the SW. ¼ sec. 28, T. 27 N., R. 32 W., and to about the same height half a mile due northeast, on the north side of the hollow. The chert is bedded, but brecciated and recemented with weathered reddish jasperoid cement.

Owing to the variability of the Grand Falls member from place to place, as already pointed out, the recognition of this formation in drill records, unless corroborated by the harmonious occurrence of the Short Creek oolite in the vicinity, is always accompanied by more or less uncertainty. But in the sheet-ground region its recognition is a matter of ease, and while only the top portion of the member is exploited for ore, its thickness can readily be determined from drill records. The top of the upper ore horizon as a rule is only a few feet from the top of the formation, and these two horizons have been used indiscriminately in drawing the structure contours, which show the elevation above mean tide level of the top of the chert.

The thickness of the Grand Falls member about Duenweg ranges from 35 to 55 feet, about Prosperity it has a maximum of 55 feet, and from Webb City to Oronogo it ranges from 35 to 45 feet. At Carl Junction the thickness is 55 feet and west of Joplin it is about the same. At Carthage the thickness is something less than 80 feet, at Alba it is 60 feet, and at Waco the chert has been penetrated for 25 feet without getting through it.

Name and correlation.—As long ago as 1855 Swallow (First and Second Ann. Repts. Geol. Survey Missouri, 1855, p. 95) described the heavy cherts about Grand Falls, but no name was applied to them until 1893, when Jenney (op. cit., p. 178) alluded to the upper beds of the Mississippian as "Cherokee limestone" and "Seneca chert," meaning by the latter presumably the chert which has herein been called the Grand Falls member. This reference is entirely too indefinite to satisfy the demands of geologic terminology. The chert at Seneca is not to be correlated with the Grand Falls, and moreover the term Seneca has been used for many years as the name of a Devonian formation in New York. Winslow (op. cit., pp. 407-419) described the chert about Grand Falls and referred to other outcrops of the chert as equivalent to the "Grand Falls chert." Thus while he did not name the member, because he considered that it was not persistent but simply a series of chert lenses, for all practical purposes he used this term as a geologic formation name, and the senior author (Smith, op. cit., p. 198) has recently adopted it as such.

A single collection of fossils was obtained from the Grand Falls chert member. This was from the upper part of the bluff on the north bank of Shoal Creek in sec. 27, 2 miles southwest of Galena, a section of which has already been given. The following list comprises the species found at this locality:

Fossils collected on Shoal Creek 2 miles southwest of Galena.

Derbya keokuk?	Chiothyris hirsuta?
Productus mesialis?	Spiriferina aff. S. solidirostris.
Productus setiger?	Canarophoria subtrigona.
Spirifer sp.	Dielasma sp.
Reticularia pseudolineata.	

There is nothing in this fauna which is particularly suggestive of the Burlington, whereas some

of the species, as, for instance, *Camarophoria subtrigona*, are distinctively Keokuk.

SHORT CREEK OOLITE MEMBER.

Character.—The Short Creek member is a thin but very persistent bed of oolitic limestone. Generally it forms a single, massive, homogeneous bed which in the Joplin area ranges from 18 inches to 8 feet in thickness. Exceptionally it divides into two beds which may have slightly different characteristics. The spherules are round, never flattened, though some are concave where they touch, as if pressed into one another or as if they had interfered with one another in the process of concentric growth. One of the most constant characteristics of the Short Creek oolite is the regularity in size of the spherules in a hand specimen, though they vary somewhat in size from place to place. They are uniformly smaller than those of the Carterville formation, averaging perhaps one-fiftieth of an inch in diameter, while the Carterville spherules are about one-twenty-fifth of an inch. The individual spherules in the Short Creek are solid and, though apparently formed by concentric growth, rarely show the center darker than the shell, as is common in those of the Carterville. The Short Creek spherules are embedded in a calcareous matrix which in places is coarsely crystalline. On a freshly broken surface the matrix gives a patchy reflection from cleavage surfaces one-fourth to one-half inch across, in which the embedded spherules appear as dark spots, a variety of poikilitic structure. In most cases the rock has a slightly splintery fracture, and this character is more pronounced the more complete the cementation. The Short Creek has two characteristics methods of weathering. In one it scales off in flakes parallel to the surface, and, being softer than the crystalline limestones above and below it in the bluff, it generally forms a slightly retreating gallery. A somewhat harder variety is found locally on sloping hillsides, where it usually breaks into rhomboid blocks which weather away along the cracks, leaving the blocks firmly fastened together and standing out somewhat in the fashion of irregular paving blocks; or the loose blocks may be strewn over the surface. In a few instances about Lowell and on Killbrev Branch the oolite is silicified into chert, but this was not noticed elsewhere. In the eastern part of the area the limestones above and below are in places altered to dolomite, but the oolite was nowhere observed to be so altered.

Thickness and distribution.—The Short Creek oolite, though thin, is a very persistent bed and outcrops all over the area where it is not carried below drainage by the dip of the rocks and where it has not been removed by erosion. Even where it lies slightly below drainage its presence is usually revealed by some of the numerous prospect shafts. The exposures along Shoal Creek show that south and southeast of Joplin the member lies from 100 to 150 feet above the water level, and that to the west it gradually drops until, in the vicinity of Lowell, it disappears. The thickness of the member increases through the same distance from 2½ feet at the east to 5 feet west of Grand Falls and 8 feet near the mouth of Harmon Branch, from which point it thins again to 2½ feet at Lowell.

Along Short Creek it crops out at Cave Springs with a thickness of 3 feet, which is increased to 8 feet at the type locality west of Empire, thinning to 6 or 7 feet at the westernmost bluff on the south side of Short Creek and to 6½ feet in the east bluff of Spring River near the Boston Mills bridge, which is 1½ miles northwest of Empire. To the west the member is below drainage except at two points along the lower mile of Shawnee Creek. At the ford a mile above the mouth of Shawnee Creek the oolite rises to the surface for a short distance in the bed of the creek. A quarter of a mile above the mouth of the creek, in the bluff forming the west bank, a bed of oolite 5 feet thick rises in a small anticline to a height of 25 feet above the creek, but goes below water level within a quarter of a mile both to the north and to the south.

Along Turkey Creek it outcrops northeast of Villa Heights, and at Midway station, on the electric line, it forms the top portion of the pinnacle known as Castle Rock. It also occurs in the bluff on the north side of the creek a mile to the east. The thickness at Midway is 6 feet. On each side of Joplin Creek north of Joplin it is 5 feet thick,

Joplin.

and at the quarry of the Joplin Lime Company, near the mouth of Opossum Hollow, it is 72 feet above the creek and 4 feet thick. Near the mouth of Leadville Hollow, half a mile west of the quarry, the Grand Falls chert comes to the surface in the valley of Turkey Creek and the Short Creek oolite horizon should be 100 feet above drainage. Half a mile west of the mouth of Leadville Hollow the thickness is reduced to 3 feet and the height above drainage to 50 feet. Within a mile or so below this point the oolite disappears beneath the water level and is not seen again except in shafts until the vicinity of Belleville is reached; at the Davison Cave Spring, half a mile southwest of the village, it has a thickness of 4 feet and an elevation of 45 feet above the creek. At the bridge over Spring River a quarter of a mile north of the mouth of Turkey Creek the oolite is typically developed, fossiliferous, very white, 4 feet thick, and about 35 feet above the river. Toward the west it dips below drainage and in the vicinity of Badger is found at depths of 40 or 50 feet.

The oolite is found all along the course of Center Creek through the district, alternating above and below water level. In the south bluff of the creek 1 mile due southeast of Smithfield the axis of the Joplin anticline raises the oolite 14 feet above the creek and it is here 2½ feet thick. A short distance to the east along the creek it sinks below drainage, and in the bottom near Lehigh and south of Carl Junction it lies at the base of the gravel, about 20 feet below the surface of the ground. It maintains this level nearly to the crossing of the Kansas City Southern Railway, beyond which it sinks deeper. In the vicinity of the mines in the creek bottom south of Oronogo it is about 20 feet below the surface, but it outcrops just at the water level by the bridge a mile southeast of Oronogo. It is exposed again under the railway bridge over the small brook a quarter of a mile west of Lakeside station, where it is 2 feet thick and 10 feet above the water. The spherules here are exceptionally small and the rock is so firmly and thoroughly cemented that only on a weathered surface or by the aid of a hand lens can its oolitic character be made out. The bed dips northward a degree or more and within a short distance goes below drainage. The next outcrop to the east is at the Scot Big Spring, 600 yards south of the Independent Powder Works, where it is 22 inches thick and 8 feet above the water level. From this point the creek bends to the south and the oolite rises higher in the bluffs, reaching an elevation of 30 to 35 feet near the mouth of Grove Creek, where it has a thickness of about 2 feet. At the Scotland Spring on Grove Creek the oolite is 3 feet thick and about 25 feet above the creek. It goes below water at the sharp bend in the creek 1½ miles east of south of Scotland. Along Center Creek above the mouth of Grove Creek the oolite, about 2 feet thick, is found in the bluffs on either side, somewhat higher on the south side, as far as the mouth of Jones Creek, above which it lies at an unknown depth, though in the small drain near the eastern limit of the district, a quarter of a mile east of the American Mill bridge, it was struck at about 10 feet below drainage. The member is exposed at several points in the bluffs of Jones and Jenkins creeks at a height of about 20 feet above the water, but, dipping 4° SW., it goes below water level on the small branch which flows northeastward through the S. ¼ sec. 24, T. 27 N., R. 31 W., about 600 yards above its junction with Jones Creek.

On Spring River, at the eastern edge of the district, the Short Creek oolite crops out in the bluffs on both sides 20 feet above the river, with a thickness of 2½ feet. Within a short distance downstream, however, it drops below water level, reappearing half a mile west of Carthage, in the bank and bed of the small stream near the bridge 200 yards east of Knell's. The bed here is 40 inches thick. At the Gaston Spring, in the bluff of Spring River at the mouth of the branch three-fourths of a mile west of Knell's, it is 30 inches thick and lies but 4 feet above the water. The oolite does not outcrop on Spring River between this point and Badger Camp, but it was struck in a shaft at Alba at a level about 60 feet below that of Spring River.

Value as a datum plane.—The Short Creek oolite, owing to its persistency, its wide distribution, and its lithologic distinctness from the other formations

of the region, is a most valuable stratigraphic datum, not the least advantageous feature of which is the fact that it can be easily recognized and followed by one without paleontologic knowledge. It has been noted at Granby, Neosho, and Seneca, Mo., and in Indian Territory nearly 50 miles southwest of the Joplin area; and Weller, as is shown more fully in the next section, has described a fauna from it at Springfield, Mo., over 50 miles northeast of this district; so that it seems probable that this oolite bed is a persistent feature of the Boone throughout southwestern Missouri.

The oolite has been found by many measurements to lie just about 100 feet above the ore-bearing sheet-ground horizon of the Grand Falls chert, and this horizon is known to lie near the top of that member. This fact has been made use of in drawing the structure contours on the economic geology map, and elevations of the two horizons have been used indiscriminately, the oolite being given the preference, however, as less liable to error in recognition.

Name and correlation.—The member is named from Short Creek, a stream flowing westward between the cities of Galena and Empire, Kans., and the type locality is the north bluff of the creek half a mile south of west of the Empire depot and a hundred yards north of the crossing of the Missouri, Kansas and Texas and the Frisco railways. Here the formation reaches its greatest thickness and here the appearance is characteristic.

A collection made at the type locality yielded the following species:

Fossils collected near Empire.

<i>Rhipidomella dubia</i> .	<i>Dielasma formosum?</i>
<i>Derbya keokuk</i> (small variety).	<i>Camarophoria subneusta</i> .
<i>Productus levicosta</i> .	<i>Avienlopecten</i> sp.
<i>Productus</i> sp.	<i>Orthonychia acutirostris</i> .
<i>Composita</i> aff. <i>C. humilis</i> .	

Fossils were collected near the mouth of Jones Creek, in the southwest corner of the SE ¼ sec. 34, T. 28 N., R. 31 W., as follows:

Fossils collected near the mouth of Jones Creek.

<i>Fenestella</i> sp.	<i>Composita</i> aff. <i>C. humilis</i> .
<i>Rhipidomella dubia</i> .	<i>Eumetria marcyi</i> .
<i>Derbya kaaskasiensis</i> .	<i>Camarophoria subneusta</i> .
<i>Derbya keokuk</i> (small variety).	<i>Pelecyrop</i> indet.

Weller (Am. Jour. Sci., 3d ser., vol. 49, 1895, p. 195) gives a section at Mackey's quarry near Springfield, Mo., as follows:

Section at Mackey's quarry, near Springfield.

Shaly gray limestone.....	6
Gray limestone.....	3
Gray limestone, stylolite band.....	1
White oolitic limestone.....	14
Hard gray limestone.....	2
Limestone and chert.....	1
Coarse crystalline gray limestone.....	5
Chert nodules in limestone.....	4
Coarse crystalline gray limestone.....	6
	36

The shaly limestone at the top of the section yielded an abundant Keokuk fauna, and the following fossils were obtained from the oolite bed, which is without doubt the Short Creek oolite:

Fossils from oolite at Mackey's quarry, near Springfield.

<i>Rhynchonella mutata</i> Hall.	<i>Platyceras acutirostris</i> Hall.
<i>Orthis dubia</i> Hall.	<i>Conocardium meekannum</i> Hall?
<i>Eumetria verneuillana</i> Hall.	<i>Batoceras</i> sp.
<i>Derbya keokuk</i> Hall.	<i>Pleurotomaria</i> sp.

It is interesting to note the similarity between this fauna and that of Spergen Hill, Ind. Both faunas are made up, to a very great extent, of diminutive forms, most of the species occurring at Springfield being also present at Spergen Hill. In both localities the rock is an oolitic limestone, and the similarity of the fauna is probably due to the similarity of the environment during deposition. In Indiana these conditions were present at the beginning of the St. Louis period, but in southwest Missouri this diminutive fauna is followed by a good Keokuk fauna [Weller, op. cit., pp. 195, 196].

Considered by itself, the fauna of the Short Creek oolite might easily be mistaken for a younger fauna; however, not only does a Keokuk fauna occur above it, but most of the species which lend this Genevieve color have at other points in the Joplin district or elsewhere been found in association with Keokuk faunas.

CARTERVILLE FORMATION.

Character.—The Carterville formation occurs in isolated patches and has a most heterogeneous character. It consists of shaly, lumpy, somewhat conglomeratic, and usually oolitic limestones, calcareous shale, light to dark argillaceous shale, arenaceous shale and shaly sandstone, massive undurindated sandstone, massive hard sandstone, and quartzite; in short, the whole category of sedimentary rocks, with the exception of chert and quartz conglomerate. The limestone is in some places shaly, but in others is firm and lumpy, the lumps ranging in diameter from a few inches to several feet. These lumps occur embedded in calcareous or argillaceous shale, and their rounded shape seems due not to water action, but to the circumstances of their deposition or consolidation. Where embedded in a calcareous matrix, the lumps are usually light gray in color, and the spherules are large, up to one twenty-fifth of an inch in diameter, or large and small mixed, some elongated, but most of them spherical, having a white shell with a darker filling. Some of the more solid chunks are rounded and smoothed as if waterworn, and clinging to them is disintegrated oolitic material which breaks away easily, leaving a continuous rounded surface. In places these harder oolitic central chunks are more or less fractured, probably owing to mud cracks in the original deposition, and these interstices are filled with fine-grained calcareous or clayey material. Where the matrix is argillaceous the limestones are very dark and the spherules are smaller and for the most part flattened or elongated. In fact, at every point this oolite presents the most obvious contrast to the even, homogeneous texture and regularity of bedding of the Short Creek oolite. The limestones, particularly the calcareous shales, and here and there the sandstones, contain an abundant, well-preserved, and characteristic fauna. The limestones occur with calcareous and argillaceous shale, and the sandstones with argillaceous shale, but in one or two instances both limestones and sandstones have been found in the same outcrop or deposit. In such cases the sandstone and arenaceous shales occupy lower stratigraphic positions than the limestone, the significance of which will appear later.

Relations.—The Carterville formation outcrops in several localities, but by far the greater number of known occurrences have been discovered in sinking prospect shafts. These deposits are in places of comparatively great thickness, but everywhere of limited areal extent, and occupy depressions in the Boone. The depth of these depressions, as compared with their areal extent, and the fact that they are surrounded by a continuous rim wall of the Boone, show that they are due to solution rather than to erosion, in other words, are sink holes. The Carterville possibly never covered the region in a continuous sheet, but if it once did so it has since been completely moved except for those deposits which were protected by their position in depressions. For this reason the outcrops are as numerous on slopes or in hollows as in other situations, and nowhere occur capping the high hills. The fossiliferous limestone and sandstone of the Carterville are in many places underlain by unfossiliferous black fissile shale, in all respects similar to the Cherokee shale. Occupying such a position, the shale is plainly Carterville. But in numerous instances such shales or sandstones, having identical stratigraphic relations, are not associated with fossiliferous rocks, and their proper reference is a difficult question. The great majority of the shale and sandstone patches scattered over the district are known to be of Cherokee age by fossils or by the occurrence of coal. The law of probabilities being followed, any doubtful case is referred to that formation. An illustration of the difficulties of Carterville stratigraphy may be seen 3 miles south and 1½ miles east of Carthage, on the farm of W. H. Black, in the E. ¼ SW. ¼ sec. 23, T. 28 N., R. 31 W. On this quarter section and extending southward across the road into section 26 is the most extensive outcrop of the Carterville formation in the Joplin area. It covers a space of about 70 acres and is surrounded by a rim of the Boone. This rim in places projects a foot or so above the adjacent shale area, and is easily traced by a limiting line of residual chert bowlders. The Carterville rises 30 feet above the lowest point in this rim and is known in one part of the area to reach a depth of over 200 feet. At

this point, which is near the north end of the outcrop, three small domes of sandstone, each about 200 feet in diameter, display quaquaversal dips of 15° to 20° on their sides. In the trough between the two western domes a shallow shaft has exposed a ledge of oolitic limestone apparently conformable with the sandstone and certainly overlying it. Both the limestone and the sandstone are fossiliferous. A number of drill holes and shafts near by show that the shale reaches a depth of 217 feet at least, and how much more is not known. A list of the fossils from the sandstone is as follows:

Fossils from sandstone on the Black place, near Carthage.

Spirifer increbescens.	Aviculoplectesh sp.
Spiriferina subelliptica?	Myalina near swallowi
Composita subquadrata.	and arkansana.
Eumetria marcyi.	

This list and the lists which follow demonstrate the Chester age of the formation.

Distribution.—The Carterville formation is known in twenty-three occurrences, the locations of which are shown on the accompanying geologic map. Some of the more important and interesting of these are described below.

In the NW. $\frac{1}{4}$ NE. $\frac{1}{4}$ sec. 3, T. 27 N., R. 31 W., fossiliferous Carterville shale and oolite were found in sinking a 50-foot shaft and digging a cellar. The dump around the shaft is made up almost altogether of oolitic rock.

Several patches of Carterville are noted on the east side of Jenkins Creek, the northernmost one of which was discovered in sinking a shaft on the Hofnagle place, in the NW. $\frac{1}{4}$ SE. $\frac{1}{4}$ sec. 1, T. 27 N., R. 31 W. Large boulders of black oolitic limestone with flattened and elongated spherules occur in a soft dark shale, both shale and limestone being fossiliferous and holding the following fauna:

Fossils from Hofnagle place, near Jenkins Creek.

Zaphrentis sp.	Productus adalrensis.
Pentremites sp.	Spirifer increbescens.
Fenestella cf. tenax.	Spiriferina spinosa.
Polypora cf. cestriensis.	Spiriferina subelliptica.
Archimedes swallowanni.	Composita subquadrata.
Archimedes compactus.	Eumetria marcyi.
Archimedes communis.	Pugnax grosveneri.
Rhombopora sp.	Bellerophon sp.
Fistulopora sp.	Dentalium (?) sp.
Stenopora sp.	Phillipsia sp.
Derbya sp.	

In the NW. $\frac{1}{4}$ SE. $\frac{1}{4}$ sec. 18, T. 27 N., R. 31 W., several prospect holes were sunk in a small area of shale and sandstone, and a small amount of ore was taken out. The dump shows large blocks of soft yellow sandstone which are highly fossiliferous. A list of the species is given herewith:

Fossils from sec. 18, T. 27 N., R. 31 W.

Fenestelloid, very abundant but indeterminate.	Composita subquadrata.
Spirifer increbescens.	Eumetria marcyi.
Spiriferina subelliptica.	Myalina near swallowi and arkansana.
	Phillipsia sp.

A characteristic occurrence of the Carterville is seen on the land of Stephen Kibler, in the SW. $\frac{1}{4}$ NW. $\frac{1}{4}$ sec. 16, T. 27 N., R. 32 W., 2 miles southwest of Duenweg. In a 20-foot shaft Cherokee shale and sandstone were encountered on top, with Carterville shale beneath, inclosing large boulders and small lumps of limestone. Both shale and limestone are fossiliferous. This occurrence seems to be a sink-hole deposit, covered later by the Cherokee formation, which spread out over most of the 40-acre tract.

On the Center Creek Mining Company's land, lying between Carterville and Webb City, a number of occurrences of the formation have been disclosed by shafts and cave-ins. They are within a short distance southwest, south, southeast, northeast, and north of the Missouri Pacific station. With the exception of the last mentioned, they show calcareous shale and the usual fossiliferous oolitic limestone with elongated spherules. The occurrence just west of the railway and 800 feet north of the station is more interesting. It is exposed by a cave-in, the west wall of which shows well the characteristic stratigraphic features of the formation. The north wall of the cave-in exhibits the section given in the next column.

While otherwise typical of the stratigraphic constitution of the Carterville, the quartzitic portion of the section is, so far as known, confined to this one locality, and the shale, which here makes

up but a small part of the section, is usually the predominant material.

Section at cave-in between Webb City and Carterville.

Chert boulders, clay, and soil	Feet.
Shale and shaly limestone	1
Oolitic limestone	2
Bluish mottled clay shale	24
Arenaceous shale	1
Massive quartzitic sandstone	6
Bluish clay shale	4
Arenaceous shale and sandstone boulders	4
Quartzite, grains secondarily enlarged	18
	464

Two miles southeast of Joplin, on the Williams land in the N. E. $\frac{1}{4}$ sec. 23, T. 27 N., R. 33 W., Cherokee black shale covers about 40 acres of the bottom. Beneath the shale, near its southern margin, the Carterville is present, as shown in the dump of a shallow shaft. The calcareous shale and limestone yield these fossils:

Fossils from the Williams land, 2 miles southeast of Joplin.

Zaphrentis sp.	Spirifer increbescens.
Stenopora sp.	Spiriferina spinosa.
Productus semireticulatus.	Composita subquadrata.
Productus cestriensis.	

A mile southeast of Central City there is an occurrence of the Carterville on the highest point in this vicinity. A shaft about 60 feet in depth penetrated yellow clay shale with a few 24-inch limestone boulders in the upper half and soft shale weathering into fine flakes, blue to olive-green in color, in the lower half. Embedded in the limestone boulders and clinging to them are blue claystone lumps with typical Carterville fossils. Other shafts 100 yards east and west of this and at the same elevation show nothing but the chert and limestone of the Boone.

A mile north of Central City, in the southwest corner of sec. 6, T. 27 N., R. 33 W., a shaft exhibited the following arrangement of the Carterville, as well as can be judged by an inspection of the dump and the upper part of the shaft:

Section of shaft 1 mile north of Central City.

Soil and shale	Feet.
Limestone boulders, large spherules	5
Fossiliferous oolitic limestone and calcareous shale	20
Solid, fine-grained, crystalline oolitic limestone	10
	40

A mile east of Galena, in the southwest corner of the NW. $\frac{1}{4}$ sec. 11, T. 27 N., R. 34 W., several shafts in the hollow went into the Carterville formation. From a shaft not over 15 feet in depth much elongate oolitic limestone and yellow shale were thrown out, the whole rich in characteristic fossils. A deep shaft near by brought up boulders of the black elongate oolitic limestone and much black shale. A list of the fossils found at this locality is given herewith:

Fossils from shaft 1 mile east of Galena.

Palaeocia obtusa.	Productus adalrensis.
Lophophyllum cf. profundum.	Productus pileiformis.
Pentremites sp.	Spirifer increbescens.
Fenestella cf. tenax.	Reticularia? sp.
Polypora cf. cestriensis.	Spiriferina spinosa.
Archimedes swallowanni.	Spiriferina subelliptica.
Archimedes n. sp.?	Composita subquadrata.
Stenopora sp.	Platyoceras sp. (?)
Derbya sp.	Chilton sp.
Chonetes near fleuvingi.	Phillipsia sp.

Winslow (op. cit., p. 408) notes the occurrence of Chester fossils near the Henry Tucher shaft, in the northwest corner of sec 84, T. 28 N., R. 33 W., on Turkey Creek, $\frac{1}{4}$ miles northwest of Joplin, and lists the following species on the authority of R. R. Rowley:

Fossils collected near Henry Tucher shaft, on Turkey Creek.

Synocladia rectistyla.	Spirifera setigera.
Chonetes sp.?	Chonetes sp.?
Archimedes sp.?	Agassizocorinus sp.?
(form)	

Keyes (Am. Geol., vol. 16, 1895, p. 89) notes the occurrence of *Chonetes*, *Terebratula*, *Rhynchonella*, *Retzia*, and *Phillipsia* at the same locality and gives Ulrich as authority for the species listed at the top of the next column.

Name and correlation.—The formation is named from the village of Carterville, just west of which there are a number of occurrences. One of these, shown in the cave-in 800 feet north of the Missouri Pacific station, gives the best notion of the stratigraphic arrangement and relations of the formation.

Fossils collected near Henry Tucher shaft, on Turkey Creek.

Agassizocorinus daetyliformis.	Meekopora approximata.
Agassizocorinus gibbosus.	Prismopora serrulata.
Spiriferina spinosa.	Prismopora n. sp.
Athyris sublamellosa.	Streblotrypa nicklesi.
Productus parvus.	Streblotrypa subsiphiacea.
Productus setigerus.	Septopora subquadrata.
Stenopora ramosa.	Polypora corticosa.
Anisotrypa solida.	Fenestella cestriensis.
Anisotrypa fistulosa	Fenestella flexuosa.
(Kelsoukia variety).	Archimedes compactus.
Batostomella abrupta.	Archimedes intermedium.
Rhombopora persimilis.	Archimedes invaginatus.

The age of the formation is indubitably Chester, as shown by the fossils which have been listed in the foregoing paragraphs. The formations nearest related, faunally, to the Carterville are the Fayetteville shale and the underlying Batesville sandstone of northern Arkansas. The arenaceous phases of the Carterville, as has been noted, occupy a lower stratigraphic position than the calcareous phases and seem to have closer faunal affinities with the Batesville, while the limestones and calcareous shales seem to be more nearly related to the Fayetteville. Certain outliers of the Batesville in Benton County, northwestern Arkansas, which have been described in the Fayetteville folio (Geologic Atlas U. S., folio 119, 1905, p. 4), betray a very striking resemblance lithologically and stratigraphically to the occurrences of the Carterville in the Joplin district.

CHEROKEE FORMATION.

Character.—The Cherokee consists chiefly of shale which included sandstones and occasional beds of coal. The flatness of the surface, the irregularity of the base of the formation, and the local deformation, preclude a perfect understanding of the stratigraphy without a more detailed study than the importance of the formation warrants. In character, the shale is in general the common drab to black fissile variety which is usually associated with the coals of the Carboniferous. Local variations are the beds of "coal bone" or "coal rash" and the fire clays beneath the coal. The sandstones are in few places persistent for any great distance, in either thickness or quality, and range from an indurated massive rock, through flagstones, to shaly sandstone and arenaceous shales. Cross bedding is very prominent and with the stratigraphic relations indicates erratic deposition along a shifting and encroaching shore line. Differences in the amount of cementation and induration have resulted in all varieties between a firm sandstone and an almost incoherent sand bed. No limestone is included in the Cherokee in the Joplin district, although elsewhere a few thin beds occur higher up in the formation.

Thickness and distribution.—Though the full thickness of the Cherokee ranges from 400 to 500 feet (Univ. Geol. Survey Kansas, vol. 3, 1898, p. 23), the lower portion of it, with which this folio has to deal, does not exceed 150 feet, except in some of the shale patches occupying depressions in the Boone. A section on the land of Frank Betty, in the NW. $\frac{1}{4}$ SW. $\frac{1}{4}$ sec. 19, T. 33 S., R. 25 E., commences on a level with the top of the Timbered Hills, and is as follows:

Section at Betty place, sec. 19, T. 33 S., R. 25 E.

	Feet.	In.
Sandstone	13	0
Shale	6	0
Coal	0	10
Fire clay	3	0
Shale	15	0
White sandstone	5	0
Shale	9	0
Sandstone	18	0
Shale	2	0
Coal	1	2
Fire clay	2	0
Sandstone	8	0
Shale	2	0
Coal	0	11
Fire clay	1	0
Shale	14	0
Fire clay	4	0
Shale	18	0
	112	11

A well a quarter of a mile farther north went about 40 feet lower than the bottom of this section and struck the Mississippian cherts. A log of the well is unobtainable, but it is highly probable that in the lower 40 feet another bed of sandstone and one of coal were encountered, for a bed of each was struck at that elevation in a well half a mile southeast of the Betty well. It is probable also that elsewhere there is a bed of coal above the 4-foot

bed of fire clay appearing in the section at Frank Betty's. This would make a total of five coals in the general section of the Cherokee, with thicknesses of 10, 14, 11, 8, and 6 inches respectively, named in order from the top down.

The sandstone which occurs in the Betty section would all be embraced in the "Columbus sandstone" (Kansas Univ. Quart., vol. 2, 1894, p. 106; Univ. Geol. Survey Kansas, vol. 3, 1898, p. 25). The term Columbus as a formation name is preoccupied, a member of the Cherokee, which is defined, rather loosely, as including the sandstones outcropping along Brush Creek east of Columbus, capping the hills (Timbered Hills) about Tehama, and passing near Baxter Springs, Crestline, and Pleasant View, and which is stated to be about 150 feet above the base of the Cherokee, with shale in the interval. This definition would undoubtedly include all the Pennsylvanian sandstones in the Joplin district and probably all which occur in the Cherokee in the type region. The thickness of shale below the sandstone becomes much less as the margin of the Cherokee is approached. The following is a section on the south slope of the hill 1 mile south of Baxter Springs:

Section of hill 1 mile south of Baxter Springs.

	Feet.	In.
Soil, gravelly	1	3
Gravel, waterworn	0	10
Sandstone, shaly	1	3
Sandstone, heavy, buff	5	0
Shale to base of hill	30	0
Chert and limestone	—	—
	38	4

Not all the sandstone appears in this section, however, for a well on the higher part of the hill shows the sandstone to be 35 feet in thickness. A well half a mile east of Eldon shows 8 feet of sandstone, separated from the chert by 6 feet of shale. At Eldon the sandstone ranges from 25 to 30 feet in thickness.

The same sandstone crops out on the river hills 2 miles east of Eldon and skirts the west slope of Shawnee Creek to a point within a mile of Crestline, where it swings eastward to Messer post-office, thence follows the west bluff of Spring River to the mouth of Cow Creek and the west bluff of Cow Creek from the mouth nearly to Pleasant View, and thence swings southeastward to and beyond Lawton and northeastward to Waco. This bed of sandstone seems to be fairly persistent along its eastern margin, but to the west it becomes erratic in thickness and less persistent. It is probably to be correlated with the heavy bed which shows in the Betty section and is the same bed that is struck in the well near the middle of the north side of the NE. $\frac{1}{4}$ sec. 35, T. 33 S., R. 24 E., south and west of the Timbered Hills, and that caps the hill half a mile still farther south. At the southern foot of this latter hill a shaft 70 feet deep penetrates shale, with three beds of coal, but no sandstone—an illustration of the thickening of the underlying shale toward the west. North of the Timbered Hills the country is almost flat. Coal crops out in almost all the streams and is reached in the wells of the interstream areas. If there is more than one bed, the interval between them must be small. The sandstone described above appears in many of the sections and occurs here above, there below, the coal. Apparently the sandstone lies about on a level with the surface of the country. A section of a drill hole at F. J. Lampton's, half a mile east and a quarter of a mile north of the northwest corner of the district, starting below the sandstone, is as follows:

Section at Lampton place, near northwest corner of Joplin district.

	Feet.	In.
Soil	5	0
Shale	15	0
Coal	0	4
Shale	26	0
Coal	1	0
Shale	23	0
Fire clay	6	0
Shale	14	0
Shale	1	0
Coal rash	23	0
Flinty rock	5	0
Coal	0	4
Shale	15	6
	135	4

This section is a further illustration of the tendency of the underlying shale to thicken toward the west.

East of Spring River the Cherokee occurs as outliers on the Boone and as patches occupying

depressions in it. A series of large outliers occurs between Spring River and Center Creek and extends from Smithfield to Neck. The "Mound," on the State line a mile northwest of Smithfield, is a sandstone remnant, capped by a heavy bed of sandstone, with more massive sandstone at the base. It is probable that these beds correspond to the two main beds in the Timbered Hills and that they are here below their normal level, having been downthrown on the west side of the Joplin anticline. There is an outlier a mile northeast of Smithfield; another and larger one, 4 miles in diameter, north and east of Carl Junction; a third between Carl Junction and Oronogo; and a fourth north of Oronogo and west of Neck. Sandstone is found in the little knolls that dot these outliers, being somewhat more abundant in the western ones.

South of Center Creek there are several similar but smaller outliers—one 2 miles south of Webb City which has a length of 1½ miles, another, just southwest of Chitwood, and a third a little southwest of Lodi.

The remaining areas of Cherokee shown on the map consist largely of shale occupying depressions in the Boone and in places reaching to considerable depths. Owing to their softness they are usually eroded slightly lower than the general level of the country.

Relations.—Throughout the western interior coal field there is a stratigraphic break of erosional unconformity and overlap between the Pennsylvanian and the underlying rocks. In the Joplin district the Cherokee formation lies unconformably, though without stratigraphic discordance, upon the irregular surface of the Boone limestone. As has been already pointed out in the description of the Boone formation, this unconformity, more or less complicated by minor faulting, is the result of two distinct cycles of erosion and solution. It is elsewhere discussed as a solution unconformity because the degradation of the Boone limestone was largely accomplished by underground solution. For this reason there is a general absence in the Cherokee of a basal sandstone or conglomerate, such as would naturally be expected; in its place is a basal breccia of residual chert blocks embedded in shale. In only one or two places some slightly rounded and abraded fragments of chert were observed in a matrix of yellow sandy clay, separating the Cherokee shale from the Boone chert.

In general the relation of the shale boundaries to the topography is such as can be interpreted only on the assumption of an extensive unconformity largely due to solution.

A typical manifestation of the unconformity is seen at many places along the eastern limit of the Cherokee formation where it caps the hills on the west side of Spring River. Tongues of Cherokee shale and sandstone extend down into the eastward-sloping valleys much below the normal level of the upper surface of the Boone formation. Typical occurrences of this sort may be seen in the southeastern part of Baxter Springs; near the Ryan branch, 2 miles west of Galena; and at the Howard coal bank a mile northwest of Badger.

Baxter Springs is situated on a plain formed of the Boone limestones and cherts, which extends southward on a level and is surmounted by the sandstone knoll a section of which has already been given. In the southeastern portion of the city a branching gully has been cut out which leads northward to Spring River. Occupying the two branches of the little valley is a bed of shale at a level more than 40 feet below the surface of the limestone plain. A drill hole shows that the shale has a thickness of 60 feet, its bottom being 10 feet below the bed of Spring River and 100 feet below the normal limestone surface, thus showing solution as well as erosion.

One mile northwest of Eldon the chert comes to the surface over an area of perhaps 80 acres, and down a branch of Brush Creek 50 yards from the western margin of this area a shaft shows 25 feet of shale. In the southeast corner of sec. 19, T. 33 S., R. 25 E., 1½ miles south of Crestline, two drill holes in the space of 1 acre struck the Boone at 40 feet, and a third, less than 100 yards distant, is reported to have gone 125 feet through shale and sandstone without striking chert. A shaft southeast of the crossroads half a mile north of Crestline struck the chert at 90 feet. In a boring 300 yards farther north it was 160 feet to the chert; and three-

Joplin.

fourths of a mile southeast, at the point where the railroad crosses Shawnee Creek, a drill hole went 138 feet in shale. Three-fourths of a mile farther southeast, down the creek, the Boone comes to the surface. Of two borings at nearly the same elevation and about 400 yards apart on the George Rogers place, 1 mile east and half a mile north of the northwest corner of the district, the western one struck the chert at 30 feet and the eastern one went 60 feet without striking it.

The outliers of the Cherokee partake of the same unconformable character as the main body of the formation. The sandstone hill 1½ miles south of Webb City may serve as an example. The Boone forms the flat just north of this hill, the junction with the Cherokee coming at the intersection of the northern margin of the hill and the electric railway line, and likewise at the western margin. But between these two contacts a tongue of shale extends a quarter of a mile in a northwesterly direction from the northwest portion of the hill, and a shaft halfway between the hill and the Frisco Railroad shows 100 feet of shale. A shaft near the road at the southwest extension of the hill goes 105 feet in shale, but a quarter of a mile to the east, at the crossroads just south of the point where the street railway cuts through the hill, there is a shallow prospect hole entirely in the Boone. Two hundred yards southeast of the shaft with 105 feet of shale there is another one showing 175 feet of shale and 4 feet of sandstone, while 200 yards southwest of this shaft the Boone limestone was struck at a depth of 20 feet. The shale forms a forked tongue reaching to Oakland and extending thence eastward to the railroad, bearing such relation to the topography, as may be seen from the map, that unconformity is the only logical conclusion.

The shale patches have already been described as occupying depressions in the Boone, thereby implying unconformity. These depressions are of two kinds, being due either to subterranean solution or to subaerial erosion. Most of the smaller, more or less circular patches seem to belong to the former class, the origin of which is taken up under the heading "Structure." But some of the outcrops of shale are so long, narrow, sinuous, and bifurcating that no explanation other than stream erosion seems adequate. Yet in most cases the channels filled by the shale were not simple erosion channels, but had been subjected with the rest of the district to the processes of underground solution, with the result that the bottoms of the channels were very uneven, descending here and there to relatively great depths, analogous to the "lost rivers" of modern cave regions. Perhaps the best example of this form is seen in the bifurcated patch north of Webb City, which sends one branching arm to the southwest and another long, slender arm directly across the present drainage in a southeasterly direction to and beyond Prosperity Junction on the electric line. It also exhibits the irregularity of bottom in a marked way.

Other examples of this form are seen at Prosperity, extending northward, at Neck, at Carthage, at Oronogo, and near Rise post-office. Just east of Joplin a strip of shale 4 miles long in a northwest-southeast direction and several smaller strips to the south seem to mark channels formed by stream erosion combined with solution. Several good examples are found about Galena and Klondike. One mile northeast of Klondike there is a peculiarly arborescent patch whose branches correspond very closely to hollows in the present topography. Whatever erosion of the shale has taken place here has served merely to uncover and restore more and more of the pre-Pennsylvanian topography. Considerations of this nature led Ball (Jour. Geol., vol. 12, 1904, p. 339) to refer to the topography of Miller County, Mo., 150 miles northeast of the Joplin district, as "resurrected." For limited areas the drainage of the district is unquestionably in part a resurrection of the former drainage system, but a study of the relation of the topography to the areal geology of the Joplin region will show that the present system intersects the former system at angles varying from 45° to 90°. Anything more than minor readjustments of the two systems by erosion of the softer shale could not reasonably be expected when it is remembered that between the pre-Pennsylvanian and the present erosion cycle there have intervened Cherokee deposition, base-

leveling, and an epirogenic movement which shifted the direction of the drainage from northwest to southwest. From the general direction of the old valleys, as well as from the manner of their bifurcation, it can be seen that the streams of the old Boone highland flowed northwestward into the Pennsylvanian sea.

Name and correlation.—The Cherokee shale was named by Haworth and Kirk (Kansas Univ. Quart., vol. 2, 1894, p. 105) from Cherokee County, Kans., the southeastern portion of which is included in the area discussed in this folio. The name Cherokee was preoccupied, for Jenney, as already remarked, had applied the term to the upper limestone of the Mississippian. The usage of the Kansas geologists, however, has so far supplanted the prior usage that it must stand as the accredited one.

The age of the formations of the western interior coal field has not yet been thoroughly studied. Collections have been studied by David White from a shaft in a shale pocket in sec. 31, T. 28 N., R. 33 W., near Belleville, in the Joplin area (Ball, U. S. Geol. Survey No. 98, 1893); from Henry County, Mo. (Mon. U. S. Geol. Survey, vol. 37, 1899); and from the Hartshorne coal beds, Indian Territory (Nineteenth Ann. Rept. U. S. Geol. Survey, pt. 3, 1899, pp. 457-538); and by Fairchild and White, from Washington County and from the Spadra and other coals of Arkansas (op. cit., p. 470).

Mr. White's correlation of these localities is as follows:

The coal measures of Washington County, Ark., are much older than the coal at Spadra, Ark., which is provisionally correlated with the Hartshorne coal of Indian Territory. The Hartshorne coal belongs at the very base of the Allegheny ("lower Coal Measures?"), or more probably below that formation. The Henry County, Mo., plants are regarded as younger than the Hartshorne and probably not far from the middle of the Allegheny ("lower Coal Measures?"). The plants collected near Belleville are rather younger than the Henry County flora. This is interpreted to mean that the sandstones and shales at the eastern margin of the Pennsylvanian area were laid down in a sea which transgressed diagonally from the Indian Territory locality to Henry County, and that the Belleville plants grew at a higher level which was not reached by the sea until after deposits equivalent to the Henry County beds had been laid down to the west.

The increase in thickness of the shale underlying the "Columbus" sandstone and the progressive thinning of that sandstone westward from the eastern margin of the coal measures, as already pointed out, are in strict accord with such an interpretation.

Fossil invertebrates of Pennsylvanian age have been found in two localities in the district. On the Bingo land, at the southeastern edge of Joplin, the dump from a shale shaft gave the following species:

Fossils from the Bingo land, southeast of Joplin.

Lophophyllum profundum.	Hustedia mormoni.
Stenopora sp.	Astartella vera.
Derbya crassa.	Nucula sp.
Spirifer inebescens?	Pleurophorus? sp.
Ambocoelia planiconvexa.	Naticopsis sp.
Composita subtilita.	Pleurotomaria sp.

A miscellaneous dump of limestone boulders and shale from a shaft near the northwest corner of the NE. ¼ SE. ¼ sec. 20, T. 28 N., R. 32 W., on the Burgner land, 1½ miles south of Carterville, yielded the following species from the disintegrated limestone boulders:

Fossils from the Burgner land, 1½ miles south of Carterville.

Zaphrentis sp.	Spirifer cameratus.
Schizophoria? sp.	Squamularia perplexa.
Derbya crassa?	Spiriferina kentuckyensis?
Productus semireticulatus.	Composita subtilita.
Marginifera n. sp?	Euomphalus catilloides.

These faunas are both Pennsylvanian, more refined correlation not being possible at this time. They are thus in substantial agreement with the testimony of the plants as to the age of the shales, which may be assigned to the middle of the Allegheny ("lower Coal Measures").

TERTIARY ROCKS.

LAFAYETTE GRAVELS.

Character.—Gravels occur widespread over the flat upland of the Joplin area, except where removed by erosion. These gravels are of chert, the occur-

rence of other material, except in the northwestern part of the area, being very rare. The pebbles are somewhat rounded by water action, though many of the rounded pieces are doubtless due to the original concretionary shape of the flint nodules. They are ordinarily glazed and stained a yellowish or reddish brown and resemble so much the gravels found along the present streams (contributing doubtless to them) that drillers record them as "creek gravel." In size the pebbles range from 2 or 3 inches (in a few instances more) in diameter down to coarse sand, though the larger sizes are exceptional, and the average diameter is about 1 inch. They are embedded in a matrix of variegated but prevalently red sand which alternates locally with reddish or yellowish ochereous clay.

Thickness and distribution.—Ordinarily, where not disturbed by a rehandling of material, the gravels are found beneath the soil in a bed varying from a few inches to several feet in thickness, but on slopes the soil and gravel are commonly mixed. The bedding of the gravel has the usual characteristics of fluvialite deposition.

Within the Joplin area these deposits are best developed along Spring River or within a few miles of it. They are well exposed all over the ridge lying between Spring River and Center Creek. There is a fine exposure 2 miles due west of Carthage, where a number of old shafts show a thickness of 6 to 18 feet of bedded gravel. This was the maximum thickness noted in the district, except in valleys where there has been reworking and accumulation of material. In the hills 3 miles north of Oronogo the gravel is cemented by iron oxide into a conglomerate which forms slabs of rock several feet in length and a foot or two in thickness. Good exposures of the gravel are also to be seen beside the road 2 miles due north of Carl Junction and near Spring River 2½ miles east of Waco.

The gravel occurs more sparingly over the ridge between Center Creek and Turkey Creek. The best exposures are on the sandstone knoll south of Webb City, where in places it has a thickness of 10 feet. It is well developed about Joplin and Galena and especially good outcrops are shown on the sandstone remnant a mile west of Lodi and on a similar hill a mile south of Chitwood, the latter exposure showing the contact of the gravel with the underlying Cherokee sandstone. Gravel is found over the high flats south of Shoal Creek and to a less extent over the high prairie in the southeastern part of the area, where the excessive thickness of soil, from 10 to 50 feet, buries the gravel so deeply that it is exposed only in excavations or gullies.

Good exposures of the gravel are found on the hills just west of Messer post-office, on the high, narrow ridge between Shawnee Creek and Spring River 3 miles southwest of Messer, and farther south on the hills north of Vark station. The higher points about Baxter Springs, especially the high sandstone knoll south of town, are covered by gravel.

The extreme northwestern part of the district, west of a line drawn from Pleasant View through Crestline to the Timbered Hills, shows no gravel. This area is entirely underlain by the Cherokee formation. On the top of the Timbered Hills the gravel is very scanty, only a few small pebbles of concretionary ironstone being present. Careful search showed but one or two small chert pebbles, which may, indeed, have found their way there adventitiously. At the eastern base of the hills genuine waterworn chert gravel occurs, but not plentifully. Along the roadside a mile northwest of Eldon there is an outcrop of waterworn gravel of local materials, concretionary ironstone and sandstone, but no chert.

Origin and correlation.—There can be no reasonable doubt that the source of the gravels is the chert of the Paleozoic limestones. In the region under consideration the gravels were certainly furnished by the cherts of the Mississippian series, because they are nearest at hand and because Mississippian fossils have been found in the chert of the gravels. That they have not been moved a great distance is true, as pointed out by Haworth (Kansas Univ. Quart., vol. 2, 1894, pp. 136-142); but it is also clear that they are not in place. True residual cherts are found in many places and in no case are they like these gravels, either in uniformity of size or in waterworn character. Some of the

residual blocks are in a measure finely shattered in weathering, yet many large pieces remain, and the smaller broken pieces preserve each sharp edge. Moreover, the binding material of residual cherts is reddish clay, whereas the gravels under consideration are interbedded with and embedded in deep-red sand. It is further to be noticed that many of the best exposed beds of gravel lie upon the Cherokee sandstone outliers, which at the time the gravels were deposited were parts of a continuous sandstone terrane. The height at which the gravels occur upon these outliers is in places 50 feet above the limestone plains below. Such gravels can have been derived only from limestone areas at a higher level than the gravels now are. The gravels on the knoll south of Baxter Springs, for instance, can not have originated at any nearer place than the hilltops across the river, at least 3 miles away, and may have traveled much farther than that. Their occurrence on this high point near Baxter Springs and elsewhere to the east and their absence at similar levels in the neighborhood of the Timbered Hills, as well as the facts cited in the preceding section, show that the limit of their distribution was determined by the limit of the current which bore them, rather than by the nature of the underlying rocks.

They were slowly etched out of their limestone matrix, reduced to the small size they now have, and perhaps partly rounded as well, in the long period of peneplanation following the post-Carboniferous diastrophism. In the latter part of this period, particularly, when the land was near base-level, they must have covered much of the surface with a heavy mantle, and lain deeply in the wide, shallow waterways and drainage basins. When the uplift followed the base-leveling the streams were quickened and the gravels swept down the sides of the dome and, in a southeastward direction, far out onto the Mississippi delta plain.

The gravels were deposited as a result of the chief uplift which followed the main period of base-leveling and which also followed more or less closely the deposition of the lignitic Eocene of southeastern Missouri. The Lafayette gravels of the Atlantic Coastal Plain and their westward continuations in the Mississippi embayment were likewise deposited at this time, as well as can be judged, and otherwise in the facts of their occurrence and distribution display such a similarity to these gravels that there can be no hesitation in correlating them as one formation.

While it is undoubtedly true, as Harris claims (Ann. Rept. Geol. Survey Arkansas, vol. 2, 1892, pp. 61, 62), that similar gravels in Arkansas are embedded in a Cretaceous matrix, this in nowise militates against the assignment of the gravels of southwestern Missouri to the Lafayette when all the facts point to such correlation and none are against it.

Hershey (Am. Geol., vol. 17, 1896, p. 38), correlates the terrace gravels of Spring River valley with the Lafayette, but takes no note of the upland gravels. It seems more in accord with the facts to make the upland gravels equivalent to the Lafayette. To do otherwise is to assign the Lafayette to the slight subsidence at the close of the main period of canyon cutting following the uplift of the peneplain, which is not the succession as traced out at other points in the Mississippi Valley.

Marbut (Univ. Missouri Studies, vol. 1, No. 3, 1902, p. 27) has called the equivalent deposits of southeastern Missouri the Piketon gravels, but until the stratigraphic relations of the gravels in the Joplin district are more thoroughly investigated no title can be assigned to them.

QUATERNARY DEPOSITS.

The deposits of the Joplin area which may be referred to the Quaternary period are the soils, alluvium, and terrace deposits. The soils and terrace deposits have already been discussed to some extent in connection with the Lafayette gravels.

SOILS.

The soils are in the main a thin deposit, but in the southeastern flat upland portion of the district they reach surprising thicknesses. Numerous wells on the prairie here go through 40 to 50 feet of soil and clay before striking rock. The gravel mantles the bed rock, so that this amount of soil

can not be wholly residual, but must be largely transported.

ALLUVIUM.

The alluvium as delineated on the map includes both the terraces and the present flood plains of the streams, the two, indeed, being in places well-nigh indistinguishable except during the prevalence of high water. It occupies the valleys of the larger streams and varies in width from a quarter of a mile in the smaller of these valleys to half a mile along Center Creek and three-fourths of a mile in Spring River valley at Carthage. Lower Spring River valley from the mouth of Cow Creek to Baxter Springs averages $1\frac{1}{2}$ miles in width; below that point it closes in again.

In depth the alluvium varies greatly from place to place. In some places the streams run over rock floors with banks of alluvium from 5 to 10 feet in height, while in others the valleys have at an earlier period been cut some distance below the present level and are now filled with alluvium and gravel, thus indicating differential movement at the time of the Columbia subsidence or the elevation which followed the Columbia. Along Center Creek from Oronogo to Carl Junction the depth to bed rock seems to average about 20 feet. On Spring River near Neck wells show from 30 to 35 feet of gravel and alluvium. On the same stream from Waco to the south side of the district the depth seems to run between 15 and 25 feet. These depths are for wells in the flood plain. Wells in the terraces would show a greater depth.

A good section of the alluvium and basal gravel can be seen in the west bank of Spring River at the railway bridge just east of Varkok and is as follows:

Section on Spring River near Varkok.

	Feet.
Soil, loam.....	1 to $\frac{1}{2}$
Clay, reddish, leached white along cracks and rootlets.....	4 to 5
Clay, light and dark blue.....	6 to 8
Gravel, white chert.....	2 to 4
Massive chert to water level.....	8 to 10

The top of the section is just about the upper limit of modern overflow.

Two miles farther up Spring River, near the crossing of the Missouri, Kansas and Texas Railway, this section is exposed:

Section on Spring River $2\frac{1}{2}$ miles northwest of Galena.

	Feet.
Soil, gray.....	$\frac{1}{2}$
Loam, light orange.....	1
Loam, red clay.....	1
Loam, red, gravelly, gravel increasing toward bottom.....	4 to 5
Gravel, chert.....	4 to 5
Limestone, gray, crystalline.....	10 to 12

The top of this section is likewise slightly above the level of overflow and is not typical of the alluvial bottoms.

TERRACES.

The occurrence and topographic aspect of the terraces have been described. The materials of the terraces have been derived by transportation and rehanding from the soils and gravels of the uplands. These either lie as a veneer upon a rock shelf or constitute the body of the terrace. The upper portion of the typical terrace consists of loam which lies upon a bed of gravel of variable thickness. The soils of the terraces are less arenaceous than those of the present alluvial flood plains.

Hershey, as before noted, places the gravels of the terraces in the Lafayette and assigns the gravels beneath the alluvial bottoms to the Columbia. It has been shown that the upland gravels should preferably be assigned to the Lafayette. The terrace deposits, being later and separated from them by an erosion interval, may, with some degree of probability, be referred to the later period, the Columbia. The white, washed gravels underlying the modern alluvium belong to a still later period.

STRUCTURE.

GENERAL STATEMENT.

The structural processes which have been operative in this district are regional uplift, resulting in the Ozark dome; orogenic deformation, producing the folds upon and bordering the dome; and underground solution, with accompanying brecciation and minor faulting. The latter uplift which involved the region occurred in mid-Ordovician time, but as

the surface rocks are of later age its effects are not now apparent, though it is conceivable that these may have predetermined the lines which were followed by the structural effects of later elevations. The next uplift with which we are here concerned occurred at the close of Boone deposition, when the Ozark highland assumed somewhat its present outline but had at the west end an elevation relative to the underground-water level approximately 200 feet higher than at present. This uplift resulted in a torsion, possibly in part due to differential elevation and in part to the twist which must occur at the curved margin of a quaquaversal dome formed from level rocks. The joint systems resulting from this torsion served as channels for the ensuing underground drainage and predetermined the directions of the zones of ore-bearing breccias which were formed at a later period. A similar elevation occurred also at the close of the Carterville, the effects of which are indistinguishable from those of the post-Boone uplift.

Another differential movement occurred in Pennsylvanian time prior to and during the deposition of the Cherokee formation. The Joplin district was above water, while thousands of feet of strata of Pottsville age were being laid down in Arkansas to the south. By a warping of the west end of the Ozark uplift the area in Arkansas became land, while the sea expanded northeastward until the Joplin area was under water. The last uplift of structural importance in the Joplin district was the one which terminated the period of peneplanation and initiated the period of canyon cutting, the early part of which was contemporaneous with the deposition of the Lafayette gravels.

The joints and fissures produced by the post-Boone elevation, enlarged by underground solution, are largely filled with ore-bearing breccias, cemented by jasperoid, dolomite, and secondary limestone. The joints and fissures resulting from the much later Lafayette elevation are largely open or filled by uncemented crushed chert. They cut across the ore horizons of the sheet ground and therefore are of later age. They are responsible for much of the "butcher-knife" flint of the drillers and probably for some of the sheeted chert in the mines. Possibly the calcite-filled seams that cut the ore in places in the Oronogo region are to be correlated with the fractures of this period. Faults of small displacement, likewise cutting the ore deposits, may also date from this time.

The chief, and probably the only, period of decided orographic movement involving the Joplin district was the one which affected the whole Mississippi Valley at the close of the Carboniferous. The orogenic character of the deformation at this time was much more pronounced to the south of the Ozark region, where it resulted in the Ouachita system of mountains. The deformation of this period is entirely independent of the earlier system of jointing, as is clear from the discordance of the ore belts representing that jointing and the structural axes of the deformation as shown by the deformation contours. It has been shown by Van Hise that the joints caused by folding are arranged in a single system parallel to the axis of folding, whereas the ore belts, as hereinafter shown, display several systems, all of them at angles to the deformation axes.

UNDERGROUND DRAINAGE.

By far the most important process which has operated to bring about the present structural features of the district is underground solution, and any attempt to give an adequate description of those features necessitates a general discussion of the subject of underground drainage.

All regions which are made up of limestone rocks have developed more or less complete systems of underground drainage. The disappearance and reappearance of the water at intervals along the courses of the smaller streams in such regions is a familiar fact. The underground stream in such cases closely parallels the surface drainage. But in many places the underground waterways are entirely independent of the topography. A study of such a system can best be made in a cave region in a district of horizontally bedded rocks, like that about Mammoth Cave in Kentucky and Wyandotte Cave in Indiana. Maps of these cave groups show that there is a more or less regular arrangement of the larger galleries intersecting more or less

nearly at right angles. These galleries owe their origin to the widening by solution of the intersecting systems of joints and crevices in the limestones of the region. The tensions, torsions, and thrusts incident to the elevation and deformation of the land are generally assigned as the causes of these systems of joints. The joints which served to initiate the systems of underground channels of the Joplin district were formed during the elevation of the land at the close of deposition of the Boone formation. As reconstructed from the ore belts, these systems of underground channels are seen to have had various trends, the most pronounced of which was about N. 25° W., followed in order of importance by N. 50° E., N. 5° W., and N. 55° W. These joints are believed to be due to stresses, chiefly torsions, developed by the post-Boone uplift, for the reason that thrust movements are not known to have accompanied that uplift and for the further reason that joints due to the simple tension of uplift would comprise but a single system parallel to the axis or tangential to the margin of the uplift, whereas the joints of this area exhibit the different trends already noted, none of which is tangential to the uplift. It is to be remembered, however, that the uplift as it exists today may not coincide with the one produced by the post-Boone movement.

Cave-region topography.—Such an underground drainage system as has been described manifests itself superficially in a characteristic set of topographic features which, from its typical development in the region of the Karst, on the north coast of the Adriatic Sea, has come to be known as Karst topography. Among these features are sink holes, closed valleys, lost rivers, and cave springs. Sink holes are the commonest feature of a limestone region. They may be hopper-shaped depressions, connecting by a narrow throat with cave galleries below, or they may be gateways to the underground system through which enter streams draining considerable territory. When such a stream has its origin near by in a large spring the depression is a closed valley, or popularly a "gulf," and has been caused by the collapsing of a section of the roof of an underground stream. A large underground stream may be traceable for miles across country by a succession of such "gulfs," at greater or less intervals, and is known as a "lost river." Nearly all the larger springs in such a region emerge from some sort of a subterranean passage, which may increase in size upward until the "cave spring" becomes a stream flowing from the mouth of a cave.

Collapsed caves.—When a cavern collapses, as a result of enlargement beyond the self-supporting strength of the roof, or from any other cause, it is locally called a "natural cave," in contradistinction to the "cave-ins" which occur when the roof of a mine drift gives way. Where caverns are below the ground-water level and full of water the water helps support the weight of the roof. Where the balance is nicely adjusted and the roof is only just self-sustaining, if the water be withdrawn the cave will collapse. In a number of cases in the Joplin district such collapsing has followed the lowering of the ground-water level by pumping in the deeper mines. Without doubt many "natural caves" have in the geologic past resulted from the draining of caverns by elevation of the land and other causes, and have developed into closed valleys.

SOLUTION UNCONFORMITY.

At the time the post-Carterville land was depressed beneath the Cherokee sea it exhibited a typical Karst topography of sink holes, caves, "lost rivers," closed valleys, and here and there valleys of surface erosion. With the incursion of the sea, shale and sandstone filled the sink holes and valleys, and the finer sediments, seeking out the cave galleries, filtered and sifted through the crevices and passages until these too were mostly filled. These sediments, filling in and around the limestone and chert debris in the floors of valleys and caves, formed the matrix of much of the basal breccia, the familiar "mixed," "confused," or "broken" ground. This sort of unconformity, where the degradation interval is largely a period of underground solution, differs from the usual erosional unconformity in two respects—in the substitution of basal breccia for waterworn basal conglomerate and of Karst topography for erosional configuration of the degraded surface.

The fact that so many of the caves and channels were filled at this time and later is no doubt one reason why so few sink holes are now found in the Joplin district. They are much less numerous here than in other limestone districts or even in other parts of the Ozark area. Another reason is found in the protection afforded the limestones by the Cherokee shale until that formation was removed during the Lafayette period. Caverns, many of them lined with large crystals of calcite, are numerous, being struck in mining or revealed by cave springs or by the drill.

GEOLOGIC STRUCTURE CONTOURS.

The geologic structure contours on the economic geology map represent, with close approximation where data are available, the configuration of the upper surface of the Grand Falls chert and its height above sea level in 10-foot intervals. This surface is chosen because it is an easily recognized horizon which persists throughout the district and which, in extensive known territory and probably in some yet unknown, is marked by valuable ore deposits. The depth of the chert at any point is the difference between the elevation of the surface as shown by the topographic contours and the corresponding elevation of the Grand Falls chert as shown by the structure contours. The Short Creek oolite and the Grand Falls chert, the interval between which is practically constant, have been used indiscriminately in preparing the contours. The elevation of these members has been determined from outcrops, mines, and shafts, and from drill records when the identity of the bed could be clearly made out from the log. In those regions which have been most thoroughly prospected and mined data are most plentiful and consequently the contours for such regions are the most accurate. In general their accuracy varies directly as the amount of mining and prospecting, but this is not true of the vicinity of Shoal Creek and other large streams along which there are many outcrops of the Short Creek oolite and the Grand Falls chert. In the area underlain by the Cherokee in the northwestern part of the district the drill holes which reach the Grand Falls chert are too few and their records too untrustworthy to permit extending the structure contours to that region.

FOLDING.

General structure slope.—Structurally, as well as topographically, the culmination of the district is in the southeastern portion, from which the rocks incline gently northward, northwestward, and westward, at a somewhat higher angle than the surface slope. This dip of the rocks carries the Boone formation below drainage a short distance west of Spring River.

A stratum of rock, particularly one of limestone, is rarely deposited in a true plane, but usually displays a series of very low domes and shallow basins with a differential variation from the true plane varying from a few inches to a few feet. This structure is depositional entirely. In the Joplin district a similar structure on a much larger scale is one of the characteristic features, as may be noted on the geologic structure map. The domes and depressions have diameters varying from half a mile up to 2 miles or more, and differential elevations varying from 10 to 30 or 40 feet. These features seem to be on too large a scale to be attributed to the processes of deposition and apparently originated in the general deformation of the district.

Joplin anticline.—The principal orogenic feature of the Joplin district, as shown by the geologic structure contours, is the pronounced anticline which enters the district from the south just east of Shoal Creek and bears northwestward to the vicinity of Waco. The inequalities of elevation arising from this orogenic movement amounted to 150 or 200 feet at least. The anticline can not be recognized from dips or surface features, except in places where streams cut across the axis and expose the Grand Falls chert, though it appears plainly on the geologic structure map. It is, in general, much steeper on the western limb than on the eastern, and possibly along its course between Joplin and Smithfield may break down into a fault. Along Turkey Creek for a mile westward from the mouth of Leadville Hollow the level of the Short Creek oolite declines abruptly, dropping over 100

Joplin.

feet in that distance, though no dips are visible and no evidence of faulting is in sight. In another locality a mile southeast of Smithfield, to judge by drill records, the descent to the west is steep, while between these two localities data bearing on the anticline are entirely wanting.

Cow Creek anticline.—West of the mouth of Cow Creek there is a short and rather sharp anticline in the Cherokee formation. Commencing in the SE. $\frac{1}{4}$ sec. 10, T. 33 S., R. 25 E., it extends west of north to the SE. $\frac{1}{4}$ sec. 33, T. 32 S., R. 25 E., a distance of a little over 2 miles. The eastern limb of this anticline is much the steeper, having dips of 35° to 40°, while those of the western limb vary from slight angles up to 10° or 12°.

FAULTING.

Duenweg fault.—A study of the logs of drill holes on the land of the M. & B. Mining Company, in the NE. $\frac{1}{4}$ SE. $\frac{1}{4}$ sec. 4, T. 27 N., R. 32 W., near Duenweg, indicates that a fault cuts across the property, passing north of both shafts and bearing nearly due east and west. The break divides the sheet ground and the broken ground in this mine and probably reaches half a mile both east and west of the M. & B. property. The maximum throw is very nearly 35 feet and is to the north. No evidence of the fault is visible at the surface.

Portland fault.—On the land of the Portland Mining Company, in the SW. $\frac{1}{4}$ SW. $\frac{1}{4}$ sec. 7, T. 28 N., R. 32 W., just north of Webb City, there is a fault which bears due northeast and reaches from the southwest corner of the section northeastward to the railroad. The displacement is 25 feet and the throw is to the southeast. This fault is likewise determined from drill records, and as there is no evidence at the surface, whether it is normal or reverse can not be determined.

Minor faults.—In addition to the larger faults just described, there are scattered over the district many small faults of a few feet displacement. Some of these occur at the surface, but most of them are to be seen only in the mines. They are both normal and reverse and are associated with more or less fracturing and brecciation. In some of them, particularly those in the sheet ground near the Cornfield "bar," the fault zone is filled with finely crushed uncemented chert, and they seem to be rather recent as compared with the original brecciation of the sheet ground. These faults, when seen in cross section or for short horizontal distances, are not distinguishable in general from the dislocations which may be referred to as "solution faults," due to the settling of large blocks as a result of underground solution. Solution faults may be associated with fracturing, slickensiding, brecciation, or any other ordinary accompaniment of faulting. The slickensided limestone walls, the broken character of the shale at the steeper peripheral contacts of the shale patches, and the local zones of brecciation, which have led to the ascription of deep faulting to the Joplin district, and have introduced such confusion into its structural geology, are no doubt due to such readjustments. Solution faulting is generally normal, but often the rock masses, subsiding under gravity, become wedged between the walls and induce small thrust faults.

When an opening is made in a plain brick wall, as for a window, some of the bricks above the opening, if not supported, will drop out, the edges of those that remain forming a rude arch, the slope of whose sides will vary with the dimensions of the bricks. Similarly, when the roof of a mine or cavern in level rocks begins to fall at any point, the break arches over toward the top, or, in miners' language, "arches out" and ceases to fall. The rough broken edges of the rocks remaining in the arch tend to assume an angle of 45° with the plane of the roof, but this angle will vary with the character of the jointing and the thickness of the strata. When the roof of a cavern gives way bodily, the break may be described as a simple shear, the rigid wall and the settling block constituting the opposing forces. Under these circumstances, in homogenous rock, the rupture will follow the plane of maximum shear, which will have 45° toward the rigid wall, relatively, the upthrown side. The actual angle will diverge from the theoretical for various reasons, but in most cases the inclination will simulate thrust faulting. This effect seems to have been particu-

larly prominent in the formation of runs and circles, and accounts in large part for the downward-diverging, slickensided walls of circles, as well as in part for the common structural arrangement of runs, in which the breccias come into contact with the limestones and dolomites along a highly inclined plane.

Though not distinguishable from ordinary tectonic faults so far as fault phenomena are concerned, solution faults differ radically from them in the restriction of the stresses to the settling block concerned and in their vertical limitation to the zone of underground solution.

BRECCIATION.

The breccias of the district may be divided into three classes—basal breccias, sheet breccias, and zonal breccias.

Basal breccias.—These, the familiar "mixed," "confused," or "broken" ground, constitute by far the largest proportion of the breccias. They correspond to basal conglomerates, save that the fragmentary material has not been rounded or waterworn, except to a slight degree in one or two instances. They were formed by shaly or arenaceous cementation of angular to subangular blocks and fragments of chert and other rocks residually concentrated upon the slopes and bottoms of the valleys resulting from the solution of the Boone limestone in the formation of Karst topography during the elevation preceding the deposition of the Cherokee formation. Further solution of the underlying limestones caused various readjustments and subsidences of the breccias, resulting in the breaking up of the shale cement and the mingling of scattered blocks of sandstone and limestone with the chert blocks of the breccias. Here and there, where not closely cemented by shaly infiltration, the breccias have been more completely cemented by jasperoid and ore. As a result of the residual origin of the fragmentary blocks in the breccias they are generally more or less weathered, the chert being "dead" or even altered to "cotton rock." The fragments may vary from small pieces upward to blocks of very large size, though in general they range from a few inches in diameter up to a foot or so.

In some places where cross sections of the lower portions of shale patches are seen the basal breccias exhibit a regular gradation from the solid bed rock upward into the body of the shale, as noted in the description of the Cornfield "bar." Good exposures of the basal breccias are rare, on account of the soft, undurated character of the matrix. A typical though somewhat small development of the breccia in the south bluff of Short Creek 14 miles west of Galena is shown in fig. 19 on the illustration sheet. The breccia lies in a trough in the limestone, the walls of which rise 15 feet above the bottom of the trough. The boulders consist of various blocks of the chert and limestone which make up the walls. They are embedded in soft, dark shale, from which they readily weather out, making the cone of talus shown in the foreground of the picture.

Sheet breccias.—These occur principally in the Grand Falls chert and especially at the sheet-ground horizon. They have been noted in many of the mines and at various places where the formation outcrops along Shoal Creek. The heavy ledges of "live," splintery chert have been thoroughly and finely crushed in place and recemented by darker bluish chert, the bedding remaining practically undisturbed. Among the characteristics of this sheet brecciation of the ledges are the following: The fragments are of small size, ranging from one-fourth to 1 inch in length; they consist of "live" chert alone; the matrix is chert, not jasperoid, and is not ore bearing; the bedding is continuous; and, as the fragments have been rotated but slightly in the ledge, many of them show conanguinity of parts. Such of these features as can be shown in a hand specimen are illustrated in fig. 18 (illustration sheet).

Between the heavy ledges are sheets of jasperoid enclosing a variable amount of angular chert fragments. The jasperoid has metasomatically replaced the thin lenticular strata of limestone which were originally intercalated with the chert. The brittle chert was brecciated apparently by gentle warping, possibly that incident to the elevation of the land, while the more elastic limestone practically escaped

brecciation. After the recementation of the chert, at the time of ore deposition, the limestone was replaced by jasperoid.

Zonal breccias.—These are the breccias associated with the runs. The sharply angular fragments consist usually of chert, here and there of limestone, rarely of other material, and, as might be expected, have a wide variation in size, ranging from small particles up to blocks as large as the heaviest ledges afford. They are cemented for the most part by jasperoid, but locally the cement is largely of dolomite, calcite, or sphalerite. Like the runs, the length is the greatest dimension of these breccia bodies, and the vertical dimension is usually next in extent, although not uncommonly the width is greater than the height. They naturally come into contact with and grade over into basal breccias.

In many places the cementation of the chert is complete, the chert and jasperoid together forming a dense rock which breaks in any direction across the chert and its matrix. Elsewhere the cementation is less complete and the rock less coherent. Where a compact rock is formed the relative amount of jasperoid varies widely. In some cases this material forms the bulk of the rock, the chert occurring in scattered boulders and fragments which may be several feet apart. As a rule, however, the chert is more abundant, being in many places largely in excess of the jasperoid. The chert occurs in fragments ranging from microscopic size to large blocks or slabs, many of which are merely large unbroken portions of beds of chert. In all cases the fragments have a sharp angular form. Here and there near-by fragments show by their shape and attitude that they were originally parts of one piece. A typical hand specimen of this breccia is shown in fig. 16 (illustration sheet).

Forces of brecciation.—The sheet and zonal brecciation is associated variously with minor faulting, solution readjustment, warping, and horizontal thrust. The processes by which these forces have accomplished the brecciation may be made clearer by the accompanying sketch (fig. 4).




FIG. 4.—Diagrams showing method of brecciation in Boone formation.

A. Depression in the limestone, extending down to the Grand Falls chert member, filled with Cherokee shale.
B. Same after compression. The shale has been horizontally shortened, and crumpled and crushed at its contact with the limestone; the wall rock has been brecciated along the chert-limestone contact and in a vertical zone below the shale pocket.

Many of the areas of "broken" or "confused" ground reach down to the Grand Falls chert, which thus forms the "bed rock" and was the competent stratum in transmitting the lateral-thrust movements of the deformation period. The thrust of the chert on one side of the area of broken ground was balanced by the opposite thrust of the chert on the other side, and movement could result only through brecciation or upward buckling. On the other hand, the thrust of overlying limestones and chert would be but partly met by the resistance of the softer depressed shale, which would be displaced upward by the lateral movement of the limestone, resulting in slickensides and upward flexure of the shale along its contact with the limestone, and simulating normal faulting along this line. This movement of the limestone would result in brecciation along the bedding planes, particularly at the contact with the Grand Falls chert. Slickensides along the bedding planes were noted near the upper limit of the Grand Falls chert on Shoal Creek southwest of Galena. This brecciated contact plane was a most favorable place for solution, which is no doubt the reason that the Grand Falls chert constitutes the "bed rock" in so many regions of "broken ground."

The zonal brecciation of the competent stratum would necessarily start in the uppermost stratum

of the chert, presumably at the joint which controlled solution originally. The broken chert resulting from this brecciation, necessarily occupying more space than in its original unbrecciated condition, would be forced upward, displacing the shale. Then the next lower stratum of chert, no longer confined by the chert above it, would be likewise brecciated, the broken chert pushing upward against the broken chert of the first stratum and further displacing the shale above. Then the third stratum would be subject to brecciation, and in time the fourth, and so on, the brecciation progressing downward until the crushing strength of the chert, added to the friction encountered by the upward movement of the loose chert, plus the weight of the superimposed shale, together balanced the force of the lateral thrust. In places along Shoal Creek where the Grand Falls chert has been subjected to rather sharp deformation it has lost its bedded character and become gnarled, knotted, and brecciated, apparently by some such process as has just been described.

Localization of brecciation.—The sheet brecciation in the Grand Falls chert took place presumably wherever there was warping, but the fracture and breccia zones are largely limited to the areas of underground solution, which, being weakened by jointing and further weakened by underground drainage, more readily yielded to the stresses of deformation. This accommodation to the stresses of the time apparently accounts for the absence of a definite and harmonious system of faulting in the district.

DISCUSSION OF SPECIAL AREAS.

The structure of the Joplin district, more particularly in the mining areas, appears at first sight to be bewilderingly complicated owing to the confusion arising from the solution unconformity and later readjustments of the underground drainage. This will be made plain in the following discussion of certain areas of typical difficulty.

Cornfield "bar."—Through the Cornfield tract of the American Zinc, Lead, and Smelting Company and neighboring tracts, a mile south of Cartersville, runs a line of disturbance which throws much light on the structural features of the district. This region, which has elsewhere (*Economic Geology*, vol. 1, 1905, pp. 119-128) been considered somewhat in detail, will be but briefly described here.

On this tract both the upper broken ground and the sheet ground are mined, the former in many places reaching down almost to the latter, being separated from it by a few feet of barren chert, or here and there cutting into or through it. The "bar" is so called because it is a barren strip, for the most part of more or less broken and confused ground, occurring in otherwise well-developed sheet ground, which has a working face of 10 feet and is reached at a depth of about 170 feet from the surface. The width of the bar varies from 50 to 300 feet, and its length, with its extension beyond the Cornfield tract, is nearly three-fourths of a mile. The relation of the bar to the surficial distribution of the Cherokee shale can be seen on mine map B, which also shows the location of the cross sections given in fig. 5.

As one approaches the bar from either side along a mine drift the first manifestation is a slight dip of the rocks toward it. Next, long cracks make their appearance in the roof parallel to the face of the bar, the dip increases, the fractures come at closer intervals and have in inverse direction and proportion to the dip, until the chert has a sheeted structure. At this point the mining usually stops because of lean ore and bad roof. Beyond, the dip may be reversed and the rocks may resume their normal position as sheet ground, or the trough of the bar may be occupied by a confused mass of limestone and chert boulders mixed with mud or masses of contorted shale. An instance of the latter sort is shown in section A-A (fig. 5), the location of which is indicated on mine map B. The broken and contorted shale, in the form of a closely appressed syncline, is flanked on either side by confused ground consisting of broken shale and chert boulders. This lies unconformably on the chert, which is more or less faulted, jointed, and sheeted, and the dip gradually flattens out in both directions from the center of the bar until the level sheet ground is reached.

In sections B-B and C-C, 300 and 500 feet

respectively from section A-A, the synclinal deformation varies from about 10 feet in the former to about 2 feet in the latter, accompanied in both cases by fracturing, sheeting, and some minor faulting.

In section D-D the central part of the bar is occupied by a mass of red clay containing blocks and boulders of limestone. On either side of the broken area large inclined slabs of limestone lean toward walls of chert, but rest in part on masses of crushed and brecciated chert. As in the other sections, the chert dips toward the center of the bar, but away from the center the dip lessens until the chert resumes its normal level position as sheet ground.




FIG. 5.—Cross sections of Cornfield "bar," on American Zinc, Lead, and Smelting Company's property. Showing the synclinal character of the bar, the shale pocket, and divergent fracturing in the chert at the edges of the fold. Location of section lines is indicated on mine map B. Horizontal scale: 1 inch=50 feet. Vertical scale: 1 inch=25 feet.

At the westernmost cross drift on the Homestead tract, about 450 feet northwest of section D-D, the chert dips 1°-2° S. on the north side of the bar and 4° N. on the south side. There are very few fractures and the bedding is not disturbed except for the gentle syncline.

On the Richland tract, 1500 feet southeast of section A-A, the continuation of the bar consists simply of a strip of barren sheet ground without fractures or dips. Where both the sheet ground and the broken "upper" ground are mined the two horizons are separated by a 4-foot ledge of white chert. Above this ledge shale is found in the cracks of the chert, then more shale with flat chert boulders, then more and more shale with irregular boulders until it becomes typical basal breccia, the shale constituting a third of the mass. In various places on the Richland and Alexandra tracts this mixed ground reaches down into the sheet ground, forming a caving spot in the roof. Depths of 100 feet or more of shale with more or less mixed ground below have been noted at various points in the shale area of this region, both in the bar and outside of it.

After noting the surface distribution of the shale and the linear shape of the bar, a study of section A-A would logically lead to the conclusion that the shale was thrown down here 185 feet from the surface as the result of faulting, probably a double fault, letting down a long, narrow block of shale. The evidence is the most decisive for deep faulting that the district affords. Section D-D likewise shows disturbance which might be interpreted as extensive faulting. But the absence of shale over the bar in this vicinity and the substitution of red clay for shale in the section argue against the presence of extensive faulting. The lack of deformation on the Richland tract, in the eastern part of the bar, and the limited deformation shown in sections B-B and C-C and in the west drift on the Homestead tract prove conclusively that the amount of deformation by both folding and faulting is small—not over 20 or 25 feet and probably not so much.

As has been seen, the unconformity alone suffices to account for the depressed position of the shale in the section. The steeply dipping limestones and cherts in section D-D and the flexed shale in section A-A are strongly suggestive of the synclinal attitude usually assumed by strata which have sunk into cavities due to solution. The general synclinal habit of the bar argues a cavity beneath, since a syncline without attendant anticlines, as we have here, would, if due to lateral thrust, increase the stress rather than relieve it. The dipping cherts, the slickensides, the normal faults, and to some extent the reverse faults, are natural incidents in the development of an underground-drainage tract such as this seems to be. It

is probable that the thrusts of the post-Carboniferous deformation period may have found relief in appressing the syncline, in reverse faulting, in horizontal slickensiding, and in sheet brecciation.

Missouri Pacific Railway cut at Carthage.—In the northwestern portion of the city of Carthage there is a patch of shale 250 yards wide and three-fourths of a mile long in an east-west direction. Near the east end, where the shale reaches a depth of 150 feet, with 30 feet or more of mixed ground below, some very rich ore was found beneath the shale. Near the west end the shale is 225 feet thick, including some basal breccia.

The cut for the Missouri Pacific Railway strikes diagonally across the shale patch, displaying the




FIG. 6.—Missouri Pacific Railway cut at Carthage, Mo. Shows synclinal structure of the shale outcrop, crumpling and brecciation at its northwestern margin, and basal breccia at the bottom of the shaft. The conjectured outline of the shale pocket is suggested.

The shale patch occupies a relatively long area under which there has been extensive solution and removal of the limestone, with the dropping of masses of shale and sandstone, which have taken, in places, the characteristic synclinal shape. What part of the shale in the depression is due to deposition in the trough and what part to subsidence into it is debatable, though the amount of solution

the position of the oolite just to the east and the occurrence of shale shown in the section and in the drifts noted, it is necessary to suppose either downward faulting or solution channels. Another interpretation might be that the beds above the oolite have been faulted down. The displacement need not have been more than 15 or 20 feet. This would account for the upturned edges





FIG. 7.—Missouri, Kansas and Texas Railroad cut one-half mile west of Empire, Kans. The Short Creek oolite in the Boone limestone at each end of the section is practically undisturbed. The middle of the section is occupied by chert, presumably from a higher horizon, filling a depression in the limestone. At the right a smaller depression is filled with Cherokee shale.

necessary is the same in either case. The compression resulting from subsidence into the V-shaped cavity has more or less crushed, crumpled, and brecciated the shale near the north end of the section, and probably the same effects would be seen at the south end also, were that visible.

Empire cut.—The Empire cut begins half a mile south of west of the Frisco station at Empire, just west of the crossing of the Frisco and the Missouri, Kansas and Texas railways. From this point westward for 1500 feet the latter railroad, skirting the north bluff of Short Creek, exposes the section shown in fig. 7. At the east end of the

section the Short Creek oolite is 8 feet thick and lies upon 20 feet of interbedded limestone and chert. About 100 feet farther west the middle portion of the limestone and chert is replaced by irregular boulders of shaly sandstone embedded in black shale and lying unconformably upon the limestone. Higher in the bluff, however, the oolite is uninterrupted, though at the west edge of the shale the oolite is about 6 feet below its normal height, a result probably of settling consequent on solution of limestone beneath. It seems clear that the whole shale and sandstone mass lies in a pocket in the face of the bluff. About the middle of the section solid chert reaches up to 30 feet from the level of Short Creek. The lower half as exposed in the creek bluff is massive, while the upper half exposed in the cut is heavy bedded and dips 20°-30° W., the dip dying out to 4° within 300 feet. The steeply dipping portion is highly fractured, the fractures being nearly at right angles to the stratification. At this point there is a vertical, open, slickensided fracture with a width of 6 inches between its walls. A hundred feet farther west the chert is interrupted for a horizontal distance of about 10 feet by a vertical band of drab to yellow clay shale with embedded boulders of chert. A shaft, perhaps 150 feet to the northwest, went down in heavy limestone, but the drifts show considerable shale. About 250 feet west of the vertical shale band cherty limestone similar to that beneath the oolite at the east end of the section crops out in the bank of the creek, dipping 10° W. At the west end of the section the Short Creek oolite appears immediately over this limestone, dipping uniformly with it. Viewed from the south across the creek the outcrops of oolite with one exception are seen to fall into alignment with a dip of slightly more than 1° W.

The problem here is to explain the occurrence of the massive and bedded cherts at the horizon of the oolite. The first suggestion is that the Grand Falls chert has been thrown up by a double fault. But several facts are opposed to this explanation: (a) Here, as in almost every other such case, the fault postulated is necessarily double. Even though displacement be granted, the faults can not be traced away from this point in any direction. (b) The synclinal shape of the chert and its upturned eastern edge are antagonistic to the theory of an upward-thrust movement; with such a movement the bed should be bent the reverse way. (c) To explain



the position of the oolite just to the east and the occurrence of shale shown in the section and in the drifts noted, it is necessary to suppose either downward faulting or solution channels. Another interpretation might be that the beds above the oolite have been faulted down. The displacement need not have been more than 15 or 20 feet. This would account for the upturned edges



A third interpretation is that solution has allowed an irregular-shaped area of the overlying rocks to settle below the level of the oolite. This accounts for the stratigraphic break and the difficulty of

the stratigraphic break and the difficulty of

tracing it, the upturned edges, and the shale in the drifts, and puts this occurrence in the same category with neighboring sink-hole deposits of sandstone and shale. The slight anticlinal structure and the spreading apart of the walls of the cleft in which the yellow shale occurs are also readily intelligible in the light of irregularities in the floor of the solution chamber. For the reason indicated in the preceding paragraph it is necessary to suppose more or less local silicification to account for the massive cherts. However, the conditions accompanying such structural relations are those most favorable to silicification—extensive water channels, shattering accompanying the settling into the basins, and the presence of shale and sandstone at close range. It is a significant fact that the heavy cherts, where identification with the Grand Falls member is difficult, are prone to occur in this fashion—i. e., the stratigraphic order is broken up, blocks of shale and sandstone are in close proximity or even mingled with the chert, and there is a difference scarcely definable between the chert and the true Grand Falls variety.

Gimlet circle.—In Jackson Hollow, in the NE. $\frac{1}{4}$ SE. $\frac{1}{4}$ sec. 24, T. 27 N., R. 34 W., on the Gimlet lease, between the Gimlet mill and the Columbus mill, there are several "cave-ins" located in a roughly elongated circle 400 feet wide and 600 feet long. These openings show that there is an overhanging rim wall of horizontally bedded bluish cherts, facing inward, against which the broken shale, with included masses of chert, as shown in fig. 8, lies




FIG. 8.—Section of wall of Gimlet cave-in, Jackson Hollow. Confused broken shale, with chert and shale masses on the margin of the cave-in, lying against an overhanging rim wall of horizontal chert.

topsy-turvy. At the west end of the cave-in east of the Gimlet mill the contact face of the chert is 18 feet in height. Below this is a drift 30 feet in height which follows the contact. Near the bottom of the drift, on the outer or chert side, a seam of shale 12 inches thick extends back 10 or 12 feet in a gallery between the level chert strata. Similar seams occur farther east in the same cave-in, and southeast of the Columbus mill and east of the circle several shafts 150 to 200 feet distant from the cave-in go through 35 to 50 feet of solid chert and strike shale beneath. The shale is thin in the center of the circle, but thickens to 15 or 20 feet nearer the rim wall. The broken shale mass extends down the contact as far as mined, about 50 or 60 feet. The ore makes against this shale mass on both sides and extends outward and inward until cut off by dead walls. The chert wall shows no fracturing parallel to the contact face, but is brecciated in horizontal planes without destroying the bedding and is recemented by black jasperoid from which the blende has been leached. The contact face is not slickensided and the shale seems to be mashed into it. All the cave-ins of the circle display similar features, but they are best seen in the one just described.

Circle deposits are not uncommon in the district. The case in hand allows better opportunities for study of the surface relations than any other known to the writer.

ORIGIN OF CIRCLES.

The origin of circles has excited a lively interest from the time of the first geologic exploration of the ore deposits. Meek (Geol. Survey Missouri, 1855, pt. 1, p. 159; pt. 2, pp. 117-119), in describing the High Point lead mine of Moniteau County, Mo., a circle deposit, with brecciated central core and downward-diverging, slickensided walls, ascribes the origin to an upward thrust of the central brecciated mass. Schmidt (Geol. Survey Missouri,

1873-74, pp. 487, 522) attributes circles to the disintegration and solution of a circular mass of limestone, with the precipitation or settling of the overlying strata, resulting in many cases in the synclinal shape of the central portion. Winslow follows Schmidt in attributing the origin to underground solution. Bain, admitting that some of the smaller circles may be so formed, found that there was a close relation between the larger circles and certain definite intersecting fault planes, which, he thought, made it impossible to ascribe their origin to solution. It has also been suggested that circles are due to intersecting parallel pairs of faults in which the opposing lateral thrusts have by simultaneous rupture resulted in a single curved fault rather than in a series of intersecting faults.

The general coincidence of circles with corresponding depressed patches of shale and the downward-diverging walls argue for thrust faulting. The occasional quaquaversal structure of the central portion of the circles and the orientation of slabs in the breccia parallel to the outer wall favor the explanation by normal faulting and upthrust of the center of the circle. The slickensided walls, the brecciation, and the numerous observed displacements are all in harmony with the origin by faulting, either normal or reverse.

But there seems to be an insuperable dynamic objection to circular thrust faulting in the fact that, by itself, it relieves no stress. Even if it be granted that the thrusts from all directions were so nicely balanced as to produce a circular fault, as suggested, there must of necessity be radial faults and breccia zones to take up the surplus which would result when the rocks of an area moved bodily toward a common point. Such radial breccia zones are not found in connection with characteristic circles.

The upward thrust of the central mass, as suggested by Meek, assumes a condition (laccolitic) so unusual in the Mississippi Valley as to require strong corroborative evidence to substantiate it. With a hade varying much from the vertical, such a circular normal fault of any considerable displacement must inevitably result in the radial disruption of the hanging wall and the upward deflection of its strata. These have not occurred.

Less easily understood is the genetic relation of the circles to the linear zones of fracturing and faulting, as suggested by Bain.

The theory of the origin of circles by underground solution, as advanced by Schmidt and followed by Winslow, somewhat amplified to fit more extended developments, offers at all points the most satisfactory explanation of this form of ore body.

In caves circular rooms are not uncommon. Many are due to simple enlargement at the intersection of two solution channels. Circular passages are also more or less common. While the solution channels in a general way follow the joints in the limestone, they do not do so exactly. When two parallel channels intersect two parallel channels, the quadrangular block inclosed will show a tendency to have the angles rounded off and in time develop a more or less circular shape, the passage assuming an annular form.

To these more or less circular openings, collapsed or otherwise, is attributable the form of the circular ore deposits. Where the collapsed opening is a circular room, the attitude taken by the subsiding rocks depends on the amount of debris on the cavern floor: (a) When the cave has been one of pure solution and the floor of the room is clear of debris, the settling may take place evenly over it, leaving the edges of the sunken strata upturned at the rim and more or less shattered and crushed. (b) Some large circular rooms, however, have a central opening connecting with a sink hole at the surface, through which enters debris that piles up in the center. The Jack Johnson mine at Chitwood is reported to have been of this character. The shaft went down through solid rock into the center of a circular room almost completely filled with ore-bearing basal breccia. When the rocks above settle onto a central cone of debris breccia or onto a remnant of the original bedded rock they assume an anticlinal or dome shape, with the outer edges brecciated, and the rim wall, in some instances slickensided, diverges downward, as has been explained in the discussion of "solution faults." The Gimlet circle in Jackson Hollow which has just been described seems to be of this class. The thin shale

at the center of the circle, thickening toward the rim, the shale breccia in contact with the outer wall, and, so far as may be determined, the quaquaversal structure of the shale area all point to this origin. (c) As annular passages are narrower than circular rooms, their collapse and the resultant sinking of the superincumbent strata produce an annular syncline.

HISTORICAL GEOLOGY.

Pre-Carboniferous land areas.—Through the Cambrian and early Ordovician periods the Ozark region, with the exception of the comparatively small crystalline area to the east, was, barring brief intervals, under continuous sedimentation. From early Ordovician until early Carboniferous time the region was for the most part above water. The irregularity of the Silurian deposits at the contact of the Ordovician and Carboniferous on the southeastern edge of the uplift, and their entire absence on the western border, indicate that the land area of that period corresponded approximately to the present area of Cambrian and Ordovician rocks, perhaps restricted somewhat on the north and south and extending an unknown farther distance to the west. The Devonian outcrop follows very closely that of the Silurian rocks except that it continues across the west end of the uplift, where the Silurian is wanting, indicating probably a restriction of the land in that direction. The patchy, nonpersistent nature of the Chattanooga shale along the southern and western sides of the uplift shows that the shore of the Devonian sea was not far inland from the present outcrop of this formation.

Boone deposition.—In the type locality on the southern border of the uplift the Boone formation was the first member of the Mississippian to be deposited, but on the western border the Boone was preceded by the Choutou and the Hannibal, whose irregular and intermittent outcrops indicate a continuation into the early Mississippian of the condition of gentle oscillation of the land near sea level which was characteristic of the border of the uplift during the Silurian and Devonian time.

With the initiation of the Boone there was a uniform subsidence of the Ozark area, conditions became settled, and 250 to 350 feet of limestones and cherts were then deposited all along the southern and western edges of the area, with correlative formations to the north and east. The formation presents its full development everywhere at the outcrop, so that its former extent landward upon the uplift, while problematical, certainly was extensive. Chert boulders with impressions of indeterminate Mississippian invertebrate remains have been found in caves in the Ordovician far up on the dome, within 20 miles of the crystalline nucleus. It is not to be understood, however, that cave deposits such as these or such as those of the Carterville imply complete submergence or the deposition of a continuous sheet of sediments, but they unquestionably prove that the subsidence was such that the waters of the Boone sea reached to that distance.

Post-Boone emergence.—During the period between the close of Boone deposition and the beginning of the Carterville the land at the west end of the uplift stood at an elevation above the underground-water level approximately 200 feet higher than now, as is shown by cave deposits of shale at that depth below the present drainage, though probably, owing to the proximity of the Pennsylvania sea to the northwest, the actual altitude above sea level was considerably less than at present. Caves are formed at or near the level of underground drainage, which is practically that of the master streams of surface drainage. Numerous pockets of shale in the limestones of the Boone formation in the Joplin area lie in such fashion that they could have been formed only as deposits in cave galleries, and many others resulted apparently from the roof giving way over caves and allowing the overlying shale and sandstone to drop into the openings. The force which elevated the Boone further manifested itself in the formation of an intersecting system of jointing. The surface waters, taking advantage of these openings, penetrated the rocks and developed an underground drainage system, while at the surface a simplified Karst topography was formed.

Carterville deposition.—This old land surface, in Chester time, was depressed beneath the sea and the heterogeneous deposits of the Carterville formation

were laid down upon it. The deposits of this age are comparatively thick in a few depressions, but altogether absent over the remainder of the district. Since all the known occurrences of the formation yield rich faunas there must have been marine connection between them, but the connecting sediments, being thin, have since been eroded.

Post-Carterville emergence.—After the deposition of the Carterville the land resumed the elevation it had during the post-Boone emergence, and those portions of the Carterville which were not protected by their position in depressions were carried away. At the same time the subterranean drainage system was carried to a much higher degree of development than during the preceding emergence, and the surface was wrought into a Karst topography of great detail.

Judged by its effects, the post-Carterville emergence was of much longer duration than the post-Boone emergence, and with this the faunal record is in entire agreement. The post-Boone emergence persisted during the whole of the St. Louis, while the post-Carterville lasted from the latter part of the Chester into the Pennsylvanian. The paleobotanic record is more definite, and from this point of view the period lasted from the Chester well on into the middle of the Allegheny ("lower Coal Measures"), during which interval thousands of feet of shales and sandstones were deposited to the south in the Arkansas Valley.

Cherokee deposition.—In middle Allegheny or "lower Coal Measures" time the sea encroached from the west, covered the Joplin district, and extended far eastward. Over the western part of the district, at present covered by the Cherokee formation, shale was first laid down; then, over this, sandstone; and with the widening of the sea and the oscillation of the shore line sandstones and shales were indiscriminately deposited, with here and there thin beds of coal. The surficial depressions were all filled up and the subterranean openings largely so, resulting in the formation of much of the basal breccia, which was mineralized at a later date. The Cherokee in a continuous mantle covered all the district and, transgressing the Boone, reached far up toward the center of the dome, though now almost altogether removed, remaining only in protecting depressions or as a few scattered outliers.

Post-Carboniferous deformation.—About the close of the Allegheny or middle of the "lower Coal Measures," an uplift raised the region of southwestern Missouri above water, in which position it has since remained. At the close of the Carboniferous period the region shared in the general diastrophic and orogenic movements of the time and suffered various forms of deformation that produced inequalities of surface from 150 to 200 feet in magnitude, which have been described on a preceding page.

Cretaceous-Tertiary penneplain.—After the close of the Carboniferous the region was subjected for a long period to the processes of subaerial and subterranean erosion, which eventually left it in the condition of a penneplain. If the present topography of the region be generalized by omitting the recent erosion and canyon cutting, and taking only the flat-topped uplands, the old penneplain may be practically reproduced, and this has been done in fig. 9. This surface east of Spring River corresponds approximately to the surface of the Boone




FIG. 9.—Sketch map of the Cretaceous-Tertiary penneplain in the Joplin district.

Shown by contours on the restored surface of the penneplain. Except that the orographic features have been entirely effaced—in other words, reduced to a base plane. West of Spring River the Timbered Hills stand as a monadnock. The gorge on Spring River below Baxter Springs and the lower country to the west show that at the time of the formation of this plain the drainage passed westward north of Baxter Springs and then southwestward to the

Neosho. The elevation of the divide between Neosho and Spring rivers in this direction is about 850 feet above tide. A description has already been given, under the heading "Tertiary rocks," of the manner in which the chert gravels were being shaped and accumulated until they mantled the low-lying surface to the east.

Post-Eocene (Lafayette) elevation.—More or less closely following the deposition of the lignitic Eocene of southeastern Missouri came the uplift of the Ozark region nearly to its present elevation, which quickened the sluggish streams of the penplain and swept the accumulated gravel down the sides of the uplift and out onto the Mississippi delta plain. The rejuvenated streams attacked the penplain with renewed energy, cutting canyons such as that of Shoal Creek, 150 to 200 feet in depth, and removing the Cherokee mantle from the greater part of the district, leaving only outliers standing here and there. At some time within this period was established the new direct southerly course of Spring River below Baxter Springs.

Post-Lafayette deposition.—When the Lafayette erosion had reached an early mature stage there occurred a slight subsidence which resulted in the deposition of sand and gravel in the valley of Spring River and, to a minor extent, in the valleys of its larger tributaries. These sediments now exist in the form of terraces fringing the valleys of the streams. Their reference to the period of deposition of the Columbia formation of the Atlantic Coastal Plain has already been suggested.

Recent elevation.—After the deposition of the terrace gravels the streams were again rejuvenated by a slight elevation. At the present time most of the terrace material has been removed from the valley of Spring River. The interterrace width of the valley varies from half a mile to a mile, the whole, with the exception of the river channel, which averages 200 yards in width, being occupied by the alluvial flood plain.

ECONOMIC GEOLOGY.

ORE DEPOSITS.

GENERAL CHARACTER.

The ore deposits of the Joplin district occur in large but very irregular bodies of chert and limestone, which are usually brecciated and cemented by or impregnated with dolomite, jasperoid, calcite, or sphalerite, and which carry important amounts of sphalerite, galena, and iron sulphide. They contain unimportant amounts of chalcocopyrite, greenockite, barite, and other minerals, and on weathering give rise to a wide variety of oxides, carbonates, sulphates, and silicates. They occur in the Boone formation, associated with certain forms of fracturing and brecciation, and have two general forms—the breccias proper and the blanket-vein or sheet-ground deposits. In the latter the brecciation is relatively unimportant, the ore occurring in thin sheets intercalated between nearly horizontal beds of chert.

COMPOSITION OF THE ORE BODIES.

MINERALS OF THE ORE DEPOSITS.

From the standpoint of relative abundance and economic importance the minerals of the district may be classed as follows: (1) Those valuable minerals which occur in sufficient amount to render their extraction profitable and which therefore constitute the ore minerals of the district; (2) those associated minerals which, though of common occurrence, are economically unimportant either in themselves or because they are not found in sufficient abundance to constitute ores; (3) those minerals which are found but rarely and in small amounts.

The first class includes both lead and zinc minerals. The lead-ore minerals comprise galena, in the belt of cementation below ground-water level, and a limited amount of cerussite, in the belt of weathering above that level. The zinc ores consist chiefly of sphalerite, in the belt of cementation, with a minor proportion of smithsonite and calamine above ground-water level.

In the second division may be included marcasite, pyrite and chalcocopyrite, calcite and dolomite, and quartz, all occurring within the belt of cementation; also, as products of weathering, greenockite, limonite, gypsum, and epsomite.

To the third division belong wurtzite, millerite,

and barite, all original minerals; also pyromorphite and anglesite, due to weathering of galena; hydrozincite and goslarite, from the weathering of sphalerite; and various other rare and unimportant minerals.

Sphalerite.—Zinc sulphide (ZnS) is the principal ore mineral of the district. It is popularly known as "jack," a number of varieties being recognized. These depend chiefly on differences in color or transparency and include among others "black," "rosin," "steel," and "ruby jack." The mineral occurs in crystals and crystal aggregates lining cavities; in crystals and grains disseminated in various gangues, especially in jasperoid; and in massive granular form in seams and as a cement for chert breccia. It crystallizes in more or less distorted, isometric-tetrahedral forms, most of the crystals with similar distortion, in two unsymmetrical parts, suggesting hemimorphic types. The tetrahedral faces are usually small, minutely pitted, and rough, the other faces being lustrous and commonly united to produce low subconical forms (fig. 20, illustration sheet). Many of the crystals are twinned according to the spinel law, and repeated twinning is not uncommon. The crystals vary in size from those which are microscopic to those 3 inches or more in diameter.

The luster of the mineral is usually resinous or resinous-vitreous, though in some cases adamantine. Some of the sphalerite crystals are iridescent with a thin film of what is perhaps chalcocopyrite or iron sulphide. The color of the mineral is in most places brownish yellow ("rosin jack"), ranging from this to a very dark brown, nearly black ("black jack"). Most of it transmits light readily only on the thin edges of fragments; however, some crystals as much as half an inch in diameter are semi-transparent. The most transparent variety is the "ruby jack," which characteristically occurs in small subtransparent crystals, usually of a rich brownish-red or dark-amber color by transmitted light.

Sphalerite as seen in thin section under the microscope occurs in two colors—deep yellow and gray with a tinge of yellow in places. The two colors generally occur together, usually as a mottling without any well-defined arrangement, though in many specimens imperfectly zoned, the yellow in narrow bands along the margin of or within the crystal or grain.

Theoretically pure sphalerite contains 67 per cent of zinc and 33 per cent of sulphur, but as found in nature it contains impurities, generally slight and consisting chiefly of iron and cadmium. The sphalerite of the Joplin district is remarkably pure, containing very small amounts of cadmium and iron. The milled sphalerite averages about 57 per cent zinc, with about 1 per cent of iron, but varies considerably in different parts of the district, partly on account of variations in the abundance of marcasite or other deleterious minerals associated with the ores, and partly on account of imperfect milling.

Smithsonite.—Zinc carbonate (ZnCO₃), next to sphalerite, is probably the most abundant zinc ore in the district. Much of the ore mined and sold as "silicate" is smithsonite, although most of the mines yielding the carbonate have also more or less associated zinc silicate. At a few mines the carbonate has been recognized and mined as such. Although both the carbonate and the silicate are important ores of zinc, they do not occur in large amount, as compared with sphalerite, and they are generally limited to a narrow vertical interval in the belt of weathering. They are not usually found in paying quantity below a depth of 50 feet, and on account of their solubility in surface waters they are seldom found at less depths than 15 or 20 feet.

Smithsonite occurs crystallized, either in individual crystals or crystal aggregates, or in massive, fine-granular condition. The crystals are uniformly small, averaging about 1 mm. in diameter. They occur in rhombohedral form, with more or less curved faces, some of them so rounded as to be almost spherical, and also in elongated, club-shaped form. They are subtranslucent or translucent, some nearly colorless, many with a characteristic greenish-gray color, and a few brown. The luster is usually vitreous to pearly, in some cases velvety. The crystals and aggregates, many of the latter having botryoidal surfaces, are found coating other minerals, especially sphalerite, or lining cavities. Smithsonite is most abundant, however, in the

massive granular condition, mainly as a metasomatic replacement of limestone. Here and there it occurs in small amount as a replacement of other substances—jasperoid, less commonly other minerals, and rarely chert. The form and structure of the more massive smithsonite are varied, depending in part on the substance replaced and the mode of replacement. A lamellar or platy form, with openings between the layers lined with crystal aggregates of smithsonite, is one of the common varieties where limestone has been replaced, though much of the ore replacing limestone is considerably less open and more compact than this. Locally hollow shells are formed as a result of the partial replacement of limestone. A cellular structure, produced in the process of weathering of the ores, by solution and the partial replacement of jasperoid, calcite, or galena by the zinc carbonate, is not uncommon. The massive carbonate is usually gray in color, and much of it resembles limestone in general appearance. It is commonly compact, though here and there rather porous.

Much of the smithsonite as sold is impure, as a result of admixture with rock and other substances, especially with limestone, which has been only partially replaced by the zinc carbonate.

Calamine.—There is some confusion in literature regarding the use of the term calamine, but in the United States it is generally applied to a basic silicate of zinc having the composition H₂Zn₂SiO₄. It is commonly known as "silicate." It is an important but, as already noted, minor ore of zinc. It is found in the zone of oxidation as a product of the weathering of sphalerite, usually associated with more or less smithsonite and occurring under somewhat similar conditions. When pure, calamine contains 54.2 per cent of metallic zinc, but the ore as marketed falls considerably short of this, being more or less contaminated with gangue material.

It occurs in both crystal and crystalline aggregates. The crystals are invariably small, orthorhombic-hemimorphic in form, and many of them tabular parallel to the brachypinacoid. Many of the aggregates are hemispherical or globular, locally in scattered sheaf-like forms. The mineral also occurs in druses or in botryoidal aggregates, coating surfaces, especially the walls of cavities. The small globular or botryoidal aggregates are commonly known as "buckshot silicate." The crystals are usually colorless, and generally show a vitreous luster; in places they are stained yellow by cadmium sulphide or red by clay. Crystal aggregates are abundant on chert and jasperoid, and not uncommon on sphalerite, calcite, and other associated minerals of the ore deposits.

The massive mineral is usually grayish, much of it with more or less of a yellow tinge, and in many places somewhat stained with clay. It is as a rule composed of small, fine-grained, spherulitic radial aggregates. In the massive form the mineral is found chiefly filling cavities in and metasomatically replacing jasperoid, and therefore much of it occurs cementing chert breccia, associated in many places with more or less unreplaced jasperoid (fig. 15, illustration sheet). Less commonly it replaces limestone; also, to a limited extent, galena or calcite, after which latter it here and there forms pseudomorphs. Rarely, calamine is found replacing chert.

In connection with calamine may be mentioned the tallo clays, which from their chemical composition appear to be merely residual clays containing more or less basic zinc silicate that has not crystallized. The amount of zinc is variable, the oxide in analyses ranging from 2 to 56 per cent, but in most cases being between 20 and 40 per cent. The chief remaining constituents are silica and water.

In color these clays are light reddish to brownish, locally yellowish, or light to dark gray. As seen in the mines, they are soft and perfectly plastic and have a peculiar, characteristic feel. They show a tendency to crumble on drying. In dry specimens they do not differ essentially in general character from surface clays. They are fine grained and rather soft—they can be scratched with the finger nail, though not easily—and have a conchoidal fracture.

Goslarite.—The hydrous zinc sulphate (ZnSO₄·7H₂O) is the first product of the weathering of sphalerite, but is seldom found on account of its ready solubility in water. It has been noted here

and there on the walls of old drifts, as small translucent stalactites and stalagmites, as white opaque globular or botryoidal forms, and as an efflorescent growth of small white crystals.

Greenockite.—Cadmium sulphide (CdS) is not uncommon near the lower limit of the zone of weathering, where it occurs as a product of secondary enrichment. It is most abundant on the surface of sphalerite crystals or on fracture surfaces within them, but is also found on galena, calcite, chert, and calamine. It is present rarely as a pigment in smithsonite crystals. As a coating it forms very thin, dull films, easily rubbed off with the finger, and with a color ranging from grass-green through yellowish green to citron-yellow.

Galena.—Lead sulphide (PbS) is the most important lead ore of the district and is commonly known as "lead." It is characterized by its lead-gray color and metallic luster, its perfect cleavage, and its high specific gravity. It occurs in crystals and crystal aggregates on the walls of cavities; in crystals, less commonly in grains, disseminated in jasperoid, selvage, limestone, or shale; and massive, in seams in fractured rocks, especially chert; also as a cement to chert breccia, at many places in association with other minerals. Its crystal form is generally a combination of the cube and the octahedron, the octahedral faces most commonly merely replacing the solid angles of the cube (fig. 22). This form characterizes the mineral both where crystallizing freely, as in cavities (pockets), and where crystallizing against resistance, as disseminations. Rarely the form is chiefly octahedral, with only very slight development of the cubic faces.

The crystals vary considerably in size, though as a rule they are medium sized or small, a diameter of 2 inches or more being unusual. Skeleton forms of galena occur locally, though they are rare. Intricate skeleton intergrowths in stalactitic form of fine-grained galena and fine-grained radial sphalerite are common at the Combination mine, east of Joplin, forming what is called "combination ore," impossible to mill satisfactorily. In places marcasite coats the outside of these galena-sphalerite stalactites; and stalactites formed chiefly of marcasite, with a small central core of granular galena, are also seen.

Pure galena contains 86.6 per cent of lead and 13.4 per cent of sulphur. The galena of the Joplin district contains as a rule only trifling impurities, but the ore as marketed has a somewhat larger proportion owing to imperfect milling. The marketed ore contains about 80 per cent of lead, the impurities consisting chiefly of gangue material and small amounts of the associated heavier minerals, iron sulphide and sphalerite. The galena of the Joplin district is nonargentiferous, silver, as shown by assay, being absent or ranging up to only 1½ ounces per ton.

Cerussite.—Lead carbonate (PbCO₃), or "dry bone," is one of the lesser ores of the district and aside from galena is the only important ore of lead. Although very small amounts have been found at all depths thus far reached in mining, it is found in quantity only near the surface. The deposits of cerussite were largely exhausted in the earlier days, though bodies of it are still occasionally encountered, and it is at present being mined, with shallow deposits of galena, in several parts of the district.

Cerussite occurs locally in small amount in tabular orthorhombic crystals or in crystal aggregates in cavities, usually associated with other minerals. In crystals it is usually colorless and transparent or subtransparent, with a greasy-vitreous or greasy-adamantine luster. It also replaces galena to a greater or less extent, in many instances forming complete pseudomorphs after it. In such cases it is usually earthy and of a light to dark-gray or nearly black color, forming the "ash mineral" and "wool mineral" of the miners.

Cerussite is found chiefly as fine-granular massive aggregates, generally more or less cellular and associated with decomposing galena in such a way as to indicate that it has replaced to a greater or less extent the matrix in which the galena originally occurred. Much of this massive cerussite is soft and earthy, though in places it is moderately hard and compact. In color it is locally light gray, though most commonly yellow or reddish, chiefly from admixed clay.

Pure cerussite contains 77.5 per cent of lead,

but the ore as marketed is in all cases impure, mainly from admixture with gangue materials.

Pyrite and marcasite.—The two iron disulphides (FeS_2), familiarly known as "mundie," are common throughout the district, though they usually occur in small amount. They are locally abundant and so mixed with the zinc ores as to be detrimental or even to render mining unprofitable. In addition to its occurrence with the ores, iron sulphide is occasionally found in small amount in the country rocks, not only near but also at a distance from any known ore deposit. It is common or even abundant in some of the surface shale patches of the mining camps, and is in fact the most abundant of the sulphides occurring in shale. As shown by drill records and samples, it is also found at various depths down to at least 850 feet, the deeper occurrences consisting of minute ro microscopic crystals disseminated in small amounts in the rock.

The two minerals occur under generally similar conditions, being in many places intermixed. Of the two the marcasite apparently predominates. Both are found in crystals and crystal aggregates, also massive and usually microgranular. The crystals of pyrite are nearly everywhere minute or microscopic. Those of marcasite which occur in cavities are usually much larger, many of them reaching a length of one-half inch or more, though numbers of these also are minute; where disseminated in rocks, they are generally microscopic. The isometric pyritohedral crystals of pyrite are usually in the form of the cube with striated surfaces, though the pyritohedron is also found. The orthorhombic forms of marcasite are somewhat more varied, much of the mineral occurring in twins, among them cyclic twinning. Iron disulphide is an almost constant constituent of jasperoid, usually in scattered microscopic crystals. It is also found either alone or associated with other minerals of the ore bodies, as a cement to chert breccia, and locally replaces to a greater or less extent chert, limestone, dolomite, jasperoid, and coal, or any carbonaceous matter, especially that adjacent to the ores.

Chalcopyrite.—This sulphide of copper and iron (CuFeS_2) is one of the accessory primary sulphide minerals and the only primary copper-bearing mineral noted in the district. Though rather common, it is always found in small amount. It occurs in small, well-defined tetragonal sphenoids with tetrahedral aspect and a length in few individuals, perhaps in none, exceeding 1 cm. These crystals are sometimes spoken of as "copper points" by the miners, though in general the mineral is not differentiated by them from iron sulphide. When chalcopyrite occurs in small crystals on sphalerite all those on a single crystal of sphalerite have parallel orientation—i. e., all similar faces give a reflection at the same time (fig. 23, illustration sheet).

The color of the chalcopyrite crystal faces is rarely, if anywhere, the brass-yellow characteristic of the freshly broken surface; they are usually more or less tarnished, in many places somewhat iridescent, and here and there frosted by corrosion to a dull, almost golden-yellow color.

Calcite.—Calcium carbonate (CaCO_3), or "tuff," in addition to its occurrence as the essential constituent of limestone, is one of the most abundant of the minerals associated with the ores. It occurs in granular form as a cement to chert breccia, particularly in the mines east of Joplin; and also in crystals or crystal aggregates lining cavities, coating other minerals, or locally in the mud filling of cavities in the ore deposits.

The crystals vary greatly in size, ranging from minute forms to those 3 feet or more in length. These larger crystals are found lining cavities or caverns which are formed in some places adjacent to the ore deposits, especially northwest of Joplin.

Gypsum.—Hydrous calcium sulphate ($\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$) occurs as a product of the weathering of calcium carbonate. It is seldom found in more than very small amount, owing to its ready solubility in water. It is seen most commonly in old drifts, in small fractures, and on the surface of various rocks, especially chert, also on and in more or less hardened clays. It usually occurs in minute prismatic or acicular crystals, or in flat stellate aggregates, locally forming a thin coating over the surface of the rock. It is sometimes found in sim-

ilar forms along the parting planes of shale, and rarely in larger crystals disseminated in the shale.

The amount of calcium sulphate contained in the mine waters is rarely so great as to approach the point of saturation. This is reached, however, in the water from the Missouri Zinc Fields pump shaft, southeast of Webb City. Notwithstanding the strong flow of water here—1000 gallons a minute—gypsum crystals are forming in the chert breccia from which the water issues at the base of the shaft. There is also a grayish efflorescent coating, formed of small arborescent growths of gypsum, on the bottom and sides of the wooden trough carrying water from this shaft.

Dolomite.—The double carbonate of calcium and magnesium ($[\text{CaMg}] \text{CO}_3$), or "spar," is a common mineral throughout the district, but is variable in its occurrence, being abundant in some of the mining camps and almost absent from others. It occurs in massive granular form and also in crystal aggregates lining cavities, in both forms almost invariably associated with or adjacent to the ore deposits. So close is this relation of dolomite and sphalerite that the saying is current in the mines about Joplin that "spar is the mother of jack;" and dolomite is rarely found outside the mining camps, especially at the surface. Several outcrops of massive dolomite, however, occur in the eastern part of the district, on Center Creek and its tributaries, Jenkins and Jones creeks. These are not known to be associated with any ore. They are, in general, much finer grained than the massive dolomites of the ore deposits, and on the whole better crystallized.

The color is usually light gray or pale buff with a reddish tinge. The massive granular variety associated with the ore deposits is as a rule characterized by a general gray color, being the "gray spar" of the miners, while the crystal aggregates are either a delicate pink, in which case they are known as "pink spar," or nearly white with pearly luster. Patches of pink dolomite, usually representing filled cavities, occur abundantly in the massive gray variety. Where weathered, both varieties are in many places more or less yellowish. The dolomite crystals in cavities do not vary greatly in size, being nearly everywhere a few millimeters in greatest diameter, usually approximating half a centimeter. The grain of the massive dolomite is somewhat variable, but usually medium and coarser than that of the Boone limestone, from which it has been derived.

Barite.—Barium sulphate (BaSO_4) is relatively rare. It occurs at only a few mines within the district, and generally in small amount.

Quartz.—While silica (SiO_2) is an important constituent of the underground waters, as shown by analysis, and quartz is one of the most abundant minerals in the district—occurring massive in cryptocrystalline form as chert and in granular aggregates in jasperoid—and while druses of minute or microscopic crystals coating siliceous surfaces, either chert or jasperoid, are common in many mines of the district, larger crystals are rare and none over an inch in length were observed.

Chalcedonic silica, bluish gray when fresh, has been noted here and there in the Grand Falls chert, both at the surface near Shoal Creek and in the sheet ground south of Oronogo, occurring mainly in small stalactitic forms in cavities in chert and jasperoid.

Hydrocarbon compounds.—The hydrocarbon compounds, although not definite minerals, are included here for the sake of convenience. The chief members of this group found in the Joplin district are bituminous coals, bitumen, and a little petroleum. The coals form an integral part of the Cherokee formation and are only indirectly connected with the ore deposits. They are found in small quantity in many of the shale patches dotted over the district, and locally in large enough amount to be of economic importance. Coal occurs in fragments in many of the small masses of Cherokee shale which are found in or associated with the ore bodies.

Bitumen, known as "tar" by the miners, is found in greater or less amount in all the rocks of the district. Its occurrence in appreciable quantity, however, is generally limited to the vicinity of the ore deposits, where it is usually found as a black viscous liquid (brown in films and by transmitted light) oozing from cavities, especially those in chert or dolomite. Though usually in small quantity, it is locally sufficient in amount to interfere with the

milling of the ores and in a few places to clog the pumps. It also occurs in scattered particles or films between the component grains of the dolomite, and, usually in similar form, is an almost constant constituent of jasperoid and is found in selvage, in which it is locally abundant.

The source of the bitumen seen in the mines is probably the Boone limestone, for bitumen is most abundant in the ores where limestone predominates in the adjacent country rocks. Whether the hydrocarbons included in the limestones are principally in the form of petroleum or of bitumen is not definitely known, though it is probably the latter.

PARAGENESIS OF THE MINERALS.

Paragenesis, the order of mineral deposition, is in few places more definitely shown than in the Joplin district for the reason that this succession is most clearly evident where the minerals form the lining or filling of cavities, and here this is one of the principal modes of mineral occurrence. The order is in most cases the same, although it is not invariable, owing to the local occurrence of two or more generations of the same mineral formed at different times.

The observed order of deposition of minerals of the ore deposits is as follows: Dolomite, chalcopyrite, galena, sphalerite, galena, chalcopyrite, marcasite, pyrite, calcite, barite, marcasite. Of these minerals, dolomite, where it occurs, is invariably the first formed in the cavities. Chalcopyrite, though sometimes found disseminated in small quantity in massive dolomite, has, wherever it occurs in cavities, formed on dolomite crystals. Sphalerite and galena have both formed over chalcopyrite, as is well shown in the Franklin mine of the Kohinoor group, west of Joplin, where some of the galena is so corroded as to show the projecting crystals of chalcopyrite which underlie it. The chalcopyrite formed on sphalerite with parallel orientation is of a later generation, and both generations have been seen, the one on, the other beneath, the same crystal of sphalerite. No chalcopyrite has been noted in sphalerite, nor on nor in galena.

As a general rule, galena has formed before sphalerite, though occasionally aggregates are found in which this order is reversed. Sphalerite belonging to different generations has often been noted. A second generation, much of it with calcite, occurs filling fractures in jasperoid at the Oronogo circle, the jasperoid itself containing disseminated sphalerite of the first generation.

Small sphalerite crystals are seen, though rarely, on larger ones of an earlier generation. Sphalerite of a late generation has been noted at the Audrain mine, west of Joplin, in a few small crystals on crystals of calcite, near their terminals; also at the Combination mine, Empire, in very small crystals in calamine and with smithsonite, apparently formed at the same time as or later than the oxidation products with which it is associated.

Iron sulphide formed earlier than galena or sphalerite has not been noted. When occurring with these minerals, it is in all cases superposed on them, showing locally a preference for one or the other, especially galena. Both marcasite and pyrite have been seen on chalcopyrite, while minute crystals of pyrite have been noted on marcasite crystals. Where these sulphides occur as metasomatic replacements they appear to have been developed simultaneously. A second generation of marcasite is sometimes seen, and in one instance was noted as occurring in the surface layer of calcite crystals. At the Mildred mine, west of Joplin, chalcopyrite and marcasite crystals are covered with calcite crystals, which in turn show hemispherical and botryoidal marcasite on their surfaces. In a number of instances in the sheet ground about Webb City marcasite has been noted in corrosion cavities in galena.

GANGUE AND ASSOCIATED MATERIAL.

The ore minerals of the district, sphalerite and galena, occur in three ways—cementing chert breccias, disseminated in other cements and gangue materials, and lining or filling cavities in the gangue or country rock.

Thus the ore bodies include not only the ore minerals and the gangue material, but also large amounts of country rock, both altered and unaltered. Probably the most abundant material is chert, and indeed the greater number of the ore deposits

are essentially chert breccias, the chert fragments being practically unaltered pieces of country rock. Limestone and shale fragments are also common. Along with these unchanged materials are large quantities of other materials, either introduced from outside or derived from the country rock through a complete process of recrystallization. These materials, which form the gangue, consist of jasperoid, dolomite, calcite, secondary limestone, selvage, and mud.

Jasperoid.—This is by far the most important gangue material. It occurs chiefly as the cement of chert breccias or intercalated with practically undisturbed beds of chert in sheet ground. When fresh it is a dark-gray to nearly black rock (occasionally medium or light gray), very fine grained and compact, not unlike a dark-colored chert, and much of it suggestive of an extremely fine-grained quartzite. It is rather brittle, breaking like chert with a conchoidal or subconchoidal fracture, but with fracture surfaces less smooth than those of unweathered chert.

This rock has been variously called quartzite by Schmidt, chertokite by Jenney, jasperite by Jenney and Shepard, and secondary chert by Winslow and Bain. It is herein called jasperoid on account of its similarity to rocks from the Aspen district, Colorado, described under that name by Spurr.

The ore which occurs in the jasperoid is mainly sphalerite. Galena is found in places, but generally in very limited amount and in small crystals. The sphalerite occurs in crystals and anhedral, some of which are of microscopic size, others ranging up to 2 inches or more in diameter. The amount of sphalerite contained in the jasperoid is extremely variable. In some places it is found only in scattered grains or not at all; elsewhere it may be so abundant as to conceal the jasperoid matrix except on careful examination.

In addition to its ore minerals, much of the jasperoid shows minute, in places numerous, grains of calcite or disseminated grains of iron sulphide. But more important than either of these is dolomite, the disseminated rhombohedra of which are locally quite as abundant as the crystals of sphalerite. Small patches of dolomite, both pink and gray, also occur. These are found not only on the surface of included chert, but also isolated, many of them with angular outline, resembling fragments of chert included in jasperoid. Some of the patches are bordered in part by aggregates of pink dolomite crystals which project into the jasperoid. Jasperoid grading into dolomite, shown in fig. 14 on the illustration sheet, is of common occurrence in some mines, having been especially noted in those on the United Zinc Companies' land at Chitwood.

While jasperoid is common in most parts of the district and in most kinds of ground, it is not found in all the mines nor in all the mining tracts. In the mines of the Continental land west of Joplin it is one of the principal gangues of ore; it is also abundant in the mines north of Chitwood, though here it is more commonly barren of ore. The dark-colored fresh jasperoid is of rare occurrence on the Missouri Lead and Zinc Company's land east of Joplin, though the weathered rock, nearly everywhere leached of its ore, is common in the shallow mines at the south end of the tract. On this land as a whole the cement of the chert breccias is calcite rather than jasperoid, although cherts predominate in the adjacent country rocks; while west of the city, where jasperoid is the prevailing cement, the country rocks appear to consist chiefly of limestone.

Here and there the jasperoid is marked by bands of greater or less width. The coarser banding is due to variations in the amount of contained sphalerite or dolomite. The finer banding is in lighter and darker shades of gray, and, as shown by the microscope, is due to variations in the amount of interstitial bituminous matter, depending on differences in texture, or to variations in mineral composition, such as the occurrence of calcite in some of the bands. Some of these finer bandings are curved, but most of them are horizontal or nearly so and parallel to the broader bands. Many of them are near and parallel to the upper surface (fig. 21, illustration sheet), and some extend only a part of the way across an exposure of the rock.

Microscopically, jasperoid consists chiefly of a fine-granular allotriomorphic aggregate of quartz, in places nearly uniform, but more commonly somewhat variable in grain (as shown in figs. 10

and 24), with many of the anhedral elongated in the direction of the vertical axis. It is quite distinct from chert, the latter where unaltered having uniformly the microcrystalline and cryptocrystalline character of the angular fragments shown in fig. 27 (illustration sheet). Between the grains of quartz is a variable but everywhere small amount of bitumen in brownish films.




FIG. 10.—Micrograph of jasperoid. An aggregate of fine-granular allotropic quartz, with sphalerite (shaded) and dolomite, the latter including minute quartz crystals. Enlarged to 33 diameters.

At numerous places dolomite occurs in the jasperoid in scattered rhombohedrons, usually with clear-cut crystal boundaries. Many of these contain as inclusions minute crystals of quartz, and rarely sphalerite is found, as a rule in well-defined crystals. Sphalerite occurring as one of the constituents of the jasperoid usually presents a




FIG. 11.—Micrograph of sphalerite in jasperoid and dolomite. In dolomite the sphalerite (shaded) shows crystal outlines, but in the jasperoid its boundaries are generally ragged. Enlarged to 25 diameters.

smooth outline where adjacent to dolomite, but where bordered by quartz its boundaries, even where it shows general crystal outlines, almost invariably yield to those of the quartz, so as to produce minute irregularities at least. The sphalerite rarely contains quartz inclusions, though here and there one or more small grains of calcite are present. Calcite also occurs at many places in irregularly shaped grains through the rock and is locally abundant. While some of the galena rarely found in jasperoid shows in thin section a general crystal outline, its borders are usually ragged and quartz inclusions are common or even abundant. Most of the features just described are shown in figs. 10, 11, 12, and 27. A very small amount of iron sulphide is almost invariably present in the unaltered rocks, in microscopic crystals or grains.




FIG. 12.—Crystal of Galena with numerous quartz inclusions and surrounded by jasperoid. Grains of sphalerite (shaded) and calcite with irregular outlines in the jasperoid. Enlarged to 24 diameters.

There has been considerable difference of opinion as to the origin of this rock. An explanation which naturally suggests itself at first sight is that it was originally of a soft, perhaps mudlike nature, hav-

ing been silicified later, and this view was taken by Jenney, Winslow, and Bain. The occurrence here and there of jasperoid in a form resembling mud pockets; the actual occurrence of mud in pockets, grading toward the bottom into a gray, siliceous rock; the fine banding of the jasperoid; the local massing of small included chert fragments near the bottom of the jasperoid, with only scattered fragments above; the disseminated sphalerite, much of which appears to the eye in sharply defined crystals; and the well-defined forms of dolomite crystal aggregates where in contact with the jasperoid are all suggestive of such an explanation. Further study, however, has led the senior writer to an entirely different interpretation. His belief is that the jasperoid is, in nearly all cases, the result of a metasomatic replacement of limestone. It occurs locally in lenticular forms such as characterize the occurrence of limestone in chert, and the manner of its occurrence in sheet ground suggests the replacement of sheets and lenses of limestone. More definite evidence is found in the fact that all stages in the process of change from unaltered limestone to jasperoid have been observed, both megascopically and microscopically. These different stages are well shown in the Quaker mine at Chitwood. Corroborative facts are the occurrence here and there of fossils, particularly crinoid stems, in typical jasperoid, as well as the presence of stylonites in a somewhat calcareous jasperoid at the Jack Johnson mine near Chitwood. This rock was dark gray and in general appearance somewhat suggestive of limestone.

Analyses of Jasperoid.

	(Oronogo.)	II (Joplin.)	III (Galena.)
SiO ₂	93.26	95.77	97.38
Al ₂ O ₃57	1.84	1.99
Fe ₂ O ₃00		
FeO.....	.69		
MgO.....	.05	.24	.09
CaO.....	.25	.54	.11
Loss on ignition.....	(?)	1.17	.77
	94.52	99.56	100.19

¹ Organic matter large.

I. George Steiger, analyst. Bain, H. F., Twenty-second Ann. Rept. U. S. Geol. Survey, pt. 2, 1901, p. 121.
II and III. L. G. Eakins, analyst. Bull. U. S. Geol. Survey No. 228, 1904, p. 297.

The general process of replacement, as shown by the microscope, is as follows: First a few scattered crystals of quartz appear in the limestone; this stage is seen in fig. 25 (illustration sheet). With increase in the proportion of quartz the limestone decreases, till in the later stages the rock consists chiefly of a granular aggregate of quartz with scattered, ragged grains of calcite, mere remnants of the former limestone. Finally even these disappear. Such calcite remnants are often found, even in the occurrences of jasperoid which most resemble mud pockets, and many of the lighter colored of the narrower bands of the banded jasperoid, as already noted, contain residual calcite grains. The quartz in replacing the limestone does not assume the fossil forms shown by the unreplaced calcite, and these forms therefore persist only as long as the calcite remains.

It seems probable, from their mutual relationship, as well as from the fact that sphalerite is found in the limestone with the first appearance of the quartz, but has not been noted in limestone wholly free from quartz, that the sphalerite and quartz have developed simultaneously. That dolomite also has developed at approximately the same time as the quartz seems probable from their microscopic characters as well as their general relations.

This explanation of the origin of jasperoid accounts for the scarcity of limestone in the ore deposits. In the sheet ground limestone is rarely found, jasperoid occupying those positions in which limestone would normally be looked for. In the breccia deposits limestone blocks are sometimes found, but the chert breccia of many mines appears to be wholly free from limestone, although dolomite and chert may border it on one side and limestone with chert on the other. While the relative proportion of chert and limestone in the Boone formation varies both vertically and horizontally, and while the ore breccias occur in the more cherty parts of the formation, nevertheless a considerable amount of limestone would naturally be expected with the breccias in most cases. The replacement

of limestone by jasperoid would not only account for the apparent absence of much or most of the limestone to be expected in the chert breccias, especially in those mines which are practically free from dolomite, as the Oronogo mines and those in the hard sheet ground, but would also explain the fractured beds of chert sometimes seen suspended in a matrix of jasperoid.

In a number of instances jasperoid has been found at the surface, as in the Circle mines at Oronogo, in the Grand Falls chert along Shoal Creek, and along Short Creek between Galena and Empire. It is invariably more or less weathered, and its color is usually light gray or buff, in places nearly white. Here and there it closely approaches chert in appearance. Microscopically, some of this surface jasperoid shows more or less recrystallization, but it still has the general character of the jasperoid as already described except for the general absence of bitumen and the minerals associated with the quartz, which have been either weathered out or merely oxidized in place.

Selva.—This term is somewhat loosely applied to any dark-colored claylike material found associated with the ores. Usually, however, its use is restricted to a well-defined, dark-colored, in many places more or less porous substance, much of it claylike, in which sphalerite and locally galena are found. Rarely it forms the chief gangue of the ore.

Typical selva, when dry, is easily scratched with a knife and in many cases with the finger nail. Much of it resembles a hardened mud; in fact, it is sometimes found grading into mud. In places in the mines it is soft and muddy, though always with a gritty feeling when rubbed between the fingers—a test commonly applied by the miners for distinguishing selva. It is usually of a medium- or dark-gray color, locally brownish, and rarely light bluish gray. It is invariably fine grained, and here and there resembles a fine-grained limestone in appearance. Some of the more porous material, as seen with a lens, appears to be composed of an aggregate of minute quartz crystals. The selva is in places, though not commonly, banded. Its mode of occurrence is not unlike that of the jasperoid, but it has nowhere been noted in large amounts, especially as cement to chert breccia.

Microscopically selva closely resembles jasperoid and in some cases is indistinguishable from it. Much, perhaps most, of it, however, is characterized by the occurrence of scattered crystals and anhedral grains of quartz, much larger than the average grain of the rock, and suggestive of the phenocrysts of igneous rocks. (See fig. 26, illustration sheet.) It is in many places more bituminous than jasperoid, being locally rich in interstitial bitumen, and like jasperoid contains more or less iron sulphide, usually in microscopic crystals and grains. Microscopic inclusions of calcite occur in the quartz, as in that of jasperoid. Calcite in irregular grains is present in some of the selva, but dolomite is rarely noted. The borders of the disseminated sphalerite crystals are on the whole smoother than those of the sphalerite disseminated in jasperoid. The local banding of the selva is shown microscopically to be due to the same causes as in jasperoid, though the distinct laminated structure occasionally seen in selva is partly due to an arrangement of many of the larger quartz grains with their longer axes parallel to the lamination.

Selva is of common occurrence in the mines throughout the district, even in those in which jasperoid is rarely found. Wherever it occurs, more or less oxidation is taking place. The massive dolomites are usually softened; galena is often found pitted or corroded or even coated with a film of oxidation products; sphalerite is in many places slightly weathered, and small amounts of its oxidation products are occasionally found. The selva itself is, in part, merely a somewhat weathered jasperoid. In a number of instances jasperoid has been found grading into selva, an excellent example of this having been noted at the A No. 1 mine, on the Roaring Spring lease (No. 58, mine map F), southeast of Joplin, the selva forming the upper, weathered portion of the mass. Some of the selva also is apparently due to the weathering of somewhat siliceous limestone and other silico-calcareous rocks of the ore deposits. Ground containing a good deal of selva usually has more or less dark-gray mud, which seems to

be selva in which the weathering is carried to a more advanced stage. A large part of such ground is soft, requiring timbering—a natural result of the softening of the siliceous cement of the breccias.

Dolomite.—Dolomite, while of minor importance as a gangue for the ores, is significant as closely related to their occurrence. Massive granular dolomite occurs in large bodies adjacent to the ore deposits in many runs, although it has not been noted in sheet ground. The coarser forms of the rock have not been observed except in connection with deposits of ore, occurring, as a rule, in contact with the ore bodies, but more or less sharply separated from them, thus constituting "bars" of barren ground against which the ores have formed. The "bars" have a vertical extent corresponding to that of the ore body, and range in width (where this is known) from a few feet to perhaps 90 feet, or, where they occur between two parallel ore bodies, to as much as 140 feet. Dolomite is found most commonly interbedded with cherts, but it also occurs as the cement of chert breccias.

The dolomite grains have in few or no cases the medium- or dark-gray color which characterizes the rock as a whole, being usually pale gray with a pink tinge, which locally affects the rock mass. The common dark color of the rock is due to interstitial matter, ordinarily seen megascopically as numerous small dark-gray or brown to blackish specks. The amount of these impurities is variable, but all the dolomite contains them to a greater or less extent. The dolomite has as a rule a rather rough surface of fracture, on which the rhombohedral forms of many of the grains may often be made out.

Microscopically the texture of the dolomite approaches panidiomorphic granular. The angular spaces between the grains are most commonly filled with jasperoid, which contains in places a small proportion of dolomite and is usually of a brownish color from bituminous impregnation; but in many specimens bituminous matter forms the predominant filling of the intergranular spaces, in which case they are generally dense or opaque. These bituminous areas form the dark specks seen in the hand specimen.

Jasperoid was found in the majority of the massive dolomites examined. It occurs intermixed with the dolomite in all proportions, though as a rule the amount is small. Analyses indicate approximately, by the percentage of silica, the proportions of jasperoid commonly present in these dolomites. As already noted, jasperoid and dolomite are sometimes found grading into each other.

That the massive dolomite has been formed by the dolomitization of limestone is shown by the general limitation of its occurrence to the ore deposits, by the inclusion in it of beds and lenses of chert and by the local occurrence of limestone in the process of passing over into dolomite. In this process the fossils are as a rule completely obliterated.

Though ore is sometimes found in cavities or pockets in massive dolomite, it rarely occurs disseminated in the mass of the dolomite itself except in small amount, and then usually in or associated with jasperoid; so that the occurrence of the ore here is to be considered, it would seem, rather as an association with jasperoid than as a dissemination in dolomite.

Limestone.—Although the massive dolomite usually grades into limestone on the side away from the ore body with which it occurs, and although chert breccias are in many places in contact with limestone and chert, yet the limestone as a rule is not found in contact with the ores themselves. Most often only here and there, if at all, is a block of limestone found in or immediately adjacent to the ore body, the limestone which was originally there having been usually altered to dolomite, jasperoid, selva, or secondary limestone, or largely removed through solution.

A secondary limestone not uncommonly occurs as a cement to chert breccia, and locally forms an important ore gangue, as in some of the mines just east of Joplin. In such cases it generally cements the moderately fine-grained breccias, not the coarse ones, which are in most cases cemented with jasperoid. This limestone is medium to fine granular, locally light gray and composed largely of clear

calcite grains, but more commonly a medium or moderately dark gray. The calcite grains are as a rule less coherent than those of ordinary limestone. These rocks usually contain small, scattered cavities lined with calcite.

As shown by the microscope, these secondary limestones uniformly contain more or less quartz in crystals or grains or in small crystal aggregates, the quartz being mainly interstitial, but also occurring as inclusions in the calcite grains. Iron sulphide in minute or microscopic grains is present in many places and locally abundant. Bitumen is sometimes seen in brownish films or scattered specks. No fossils have been noted in any of these secondary rocks. While the coarser grained examples are in many cases merely aggregates of calcite, which has crystallized directly as a cement to the breccias, it is probable that many if not most of the occurrences are due to the recrystallization of original limestone. The ore occurring disseminated in this gangue is both sphalerite and galena, the latter predominating in some places.

Shale.—In addition to the surface occurrences already described, small bodies of Cherokee and Carterville shales, in places containing fragments of coal, are often encountered in the ore deposits or adjacent to them, having been dragged down along planes of faulting or entrained by underground water. Many of these bodies show by slickensided surfaces the effects of crushing. Chert fragments and other foreign materials are not uncommonly found mixed with the shale.

The surface shales when fresh are very fine-grained to aphanitic, dark-gray, nearly black, and usually very fissile rocks, breaking easily into small, thin flakes. Much of the shale or "soapstone" as found in the mines, however, is so compact that its fissile character is lost and it fractures across the lamination as readily as parallel to it.

Iron sulphide.—The occurrence of iron sulphide as a gangue is rare. It has been noted, however, in two mines, the Combination and the Walcott, near the north end of the Missouri Lead and Zinc Company's land, just east of Joplin. The mineral is abundant at both of these mines, and occurs locally in a very fine granular form, in which sphalerite is disseminated. This gangue has been formed as a replacement of original limestone, probably siliceous, and also of a silico-calcareous cement of chert breccia, in both of which the sphalerite was originally disseminated.

Mud.—Small amounts of sphalerite and galena are sometimes found as disseminated crystals in mud pockets, though few of the occurrences are of any importance. Where it contains ore the mud is dark gray or blackish in color and near the lower limit of oxidation in position.

WEATHERING.

While the general weathering of the rocks of the district does not extend to any considerable depth, surface waters are in many places carried by favorable channels to depths of several hundred feet before losing their power of oxidation, and along their courses more or less weathered ores and rocks are found. Chert approaching cotton flint has been noted in drill cuttings to a depth of more than 300 feet.

The first effect of the action of descending surface waters, as seen in the mines, is on the dolomite. The massive forms gradually soften through general solution, becoming in many cases a mass of slightly coherent grains, while the dolomite disseminated in the jasperoid is removed by solution, leaving the jasperoid filled with the sharp molds of the dissolved dolomite crystals. This leaching of the dolomite is seen in many places where the ores themselves show no sign of oxidation; the same jasperoid may contain fresh, unaltered crystals of sphalerite. Dolomite is rarely seen nearer the surface than at a depth of 20 feet. Calcite, the other principal carbonate of the ore deposits, apparently weathers less readily than the dolomite, and crystals of calcite have been found forming in mud pockets within dolomite which is in process of weathering. Where calcite occurs as a cement to breccias it is in places unaltered, while the dolomite is very soft, although in other similar instances the calcite also may be more or less dissolved. The solution of limestone begins at the surface and along fractures, and does not result in general softening as soon as in the case of the more porous dolomite.

Joplin.

The other important rocks of the ore deposits are siliceous, including chiefly chert and jasperoid. These also weather by solution, though as a rule much more slowly than the calcareous rocks; and even where active weathering is taking place both are frequently found, little altered, close to the surface, though the jasperoid is changed from dark gray to light gray or buff, and usually more or less colored with clay.

Jasperoid weathers more readily than chert, owing partly to its less compact texture and partly also to the porosity resulting from the early weathering out of its disseminated sphalerite, dolomite, calcite, and iron sulphide. At a number of points along Shoal Creek jasperoid may be seen more or less softened or in places partially dissolved, while the associated Grand Falls chert is still very fresh. Under ground, the first stage in the weathering of the jasperoid is the production of selvage, and the final stage results in the black mud so commonly seen in the mines.

In many places chert shows more or less alteration at the same depth at which dolomite begins to weather. The alteration of chert is in some instances selective, and one bed may be in the condition of cotton flint while adjacent ones are still fresh, the difference being due in part to difference in texture. The tallow clays of the ore deposits are for the most part the result of the weathering of chert, and cherts have been noted in the process of altering to such clays. These clays are usually limited to the zone in which sphalerite is weathering, where they occur in layers up to several feet in thickness, also in pockets, or intermixed with the oxidizing ores.

Of the principal metallic sulphides, sphalerite appears to be the most readily weathered, though the general oxidation of marcasite and sphalerite occurs at approximately the same level. Sphalerite readily oxidizes close to the level of ground water. Where it occurs in jasperoid the line of oxidation is locally so sharply marked that fresh, unaltered sphalerite may be seen within a distance of 1 inch from a part of the rock entirely leached of its ore. In the zone between the two, in one case noted, the color changes from that characteristic of rosin jack to a dark steel blue. More or less oxidized sphalerite occurs at all depths which have been reached by mining. In the regions of extensive oxidation sphalerite is rarely found in quantity above a depth of 50 feet and is usually absent above 20 feet.

Zinc sulphate, the first product of the oxidation of sphalerite, taken into solution by underground waters, is in part carried deeper and in part reacts with substances in solution in the waters and with the adjacent rocks. With limestone it forms zinc carbonate, which to a large extent replaces the limestone metasomatically; with the siliceous rocks, chiefly jasperoid, it forms calamine, which also occurs largely as a replacement of the rock with which the reaction takes place. With the continuation of the process of weathering, the calamine and smithsonite are gradually dissolved and carried downward. The comparative readiness with which they dissolve in descending waters is shown by the fact that they are seen in but few places near the surface.

On the weathering of the sphalerite its impurities are also oxidized, the cadmium sulphide contained in it being thus changed to the sulphate. This, carried downward as the sulphate or other salt, reacts in part near ground-water level with the zinc sulphide of unoxidized deposits, resulting in the precipitation of cadmium as greenockite, with the oxidation of a corresponding amount of the sphalerite. The reaction is as follows: $CdSO_4 + ZnS - ZnSO_4 + CdS$. The greenockite films so often seen on the ores are the result of this reaction.

Although galena is frequently found corroded and is in many places accompanied by small amounts of oxidation products close to the lower limits of oxidation, on the whole it weathers much less readily than the other sulphides of the ore bodies, and it is often found little or not at all altered close to the surface, or, as popularly expressed, "at the grass roots." In weathering, much of the galena shows at first only a simple pitting or corrosion. This etched and corroded galena is in many instances lustrous, locally iridescent or tarnished, and here and there the crystals are merely rounded, with a velvety surface due to slight differential corrosion along cleavage lines.

Galena, like sphalerite, is probably first oxidized to the sulphate, although as lead sulphate is only rarely found as a mineral, in spite of its difficult solubility, it is probable that most of the sulphate formed is immediately changed to the carbonate, cerussite, which is the principal product of the weathering of galena.

The sulphides of iron, marcasite and pyrite, both weather readily, the marcasite somewhat more so than the pyrite. The products of their weathering are principally limonite and various hydrous and basic sulphates of iron, the limonite being the final product, the least soluble, and the one most often seen.

Unusually rapid oxidation of the zinc and iron sulphides takes place on their exposure to atmospheric action in some poorly ventilated drifts. This is sometimes shown by the formation of secondary products on the walls of the drifts, also by an increased amount of zinc and iron sulphates in the mine waters. Some of these, formerly potable, become heavily charged with both sulphates. Among the characters favorable to such oxidation are, apparently, an open, porous gangue, as for example some forms of selvage; an abundance of ore, for the most part moderately fine grained; and a moderate amount of iron disulphide, the ferric sulphate from the oxidation of which probably aids materially in the unusually rapid oxidation of the sphalerite. To such oxidation of sphalerite and marcasite, particularly of the former, the "hot ground" seen in some mines appears to be largely due, and the heat in turn, together with the close, moist atmosphere, tends to make the rate of oxidation still more rapid. In the hottest of these mines the temperature is such that the miners can not work in them for more than five or ten minutes at a time. In one of the hot drifts of the Norsworthy mine, east of Joplin, the temperature (taken in midwinter) is reported to have been 120° F.

FORMS OF THE ORE BODIES.

The forms of the ore bodies, some simple, others complex, all fall into two general groups, the first including runs and their modifications; the second consisting of blanket veins, or, as they are generally known in this district, sheet-ground deposits.

RUNS.

Runs are irregular but usually elongated, in places tabular and inclined, bodies of ore, uniformly associated with disturbed strata which have been subjected to brecciation, slickensiding, and moderate displacement as the result of minor faulting or of dislocation due to underground solution. Simple runs are linear and continuous, straight or simply curved ore bodies, usually at nearly the same level throughout their extent. On account of the complication of minor faulting, underground solution, and general deformation in the ore-bearing areas of the district, such runs are rare. Many runs which have a simple structure in cross section and which for short distances appear to be simple runs are found on further exploration to be compound. Even the few instances of simple runs which have been noted may present more complex features as the development of the ore body proceeds. It thus happens that there are all gradations from simple runs to complicated compound runs. Among the latter are those formed by the lateral connection of two nearly parallel simple runs so as to form a single ore body. Another class, comprising by far the larger number of runs in the district, consists of the irregular ore bodies formed in disturbed areas that are due to complicated underground drainage. Combinations of these give runs of great complexity, among them, here and there, circular or subcircular deposits. It will be seen, therefore, that while theoretically the different types of ore deposits are distinct, practically they grade into one another, with no hard and fast lines between them.

The greatest dimension of the runs is horizontal, although as compared with that of ore bodies in other regions this linear extent is generally short, as a rule not exceeding a few hundred feet. The Arkansas run at Belleville, with a length of over one-fourth of a mile, described by Bain (op. cit., pp. 139-140), is exceptionally long. It does not, however, preserve a uniform direction, but varies

from southeast to east, then to south, and finally to west. The runs have a maximum width of 300 feet, but as a rule are between 10 and 50 feet wide. The average vertical extent is about the same, but in some cases reaches 150 feet.

In the most common structural phase of the runs about Joplin the ore-bearing breccias and the massive secondary dolomite come into juxtaposition along a highly inclined contact plane, the breccias usually forming the overhanging side. This relation is the result of a dislocation due in most cases to underground solution, as has been explained in the discussion of "solution faults." Behind the irregular zone of dolomite is unreplaced limestone and chert country rock, and behind the chert breccia are bedded cherts and subordinate limestones. The contact of the dolomite and chert breccias is usually sharp, and in many places some slickensiding is seen; locally, however, there is more or less diffusion, patches of jasperoid being found in the dolomite and patches of dolomite in the breccias. The dolomite zone has been formed by metasomatic replacement of the limestone; hence the width of the dolomite and its contact with the limestone are irregular, the limestone itself in rare instances forming the wall. In many places the dolomite is nearly horizontally bedded, and locally it shows fractures or is brecciated near the contact, giving rise where associated with interbedded chert to chert breccia with a dolomite matrix.

Both coarse and fine ore-bearing breccias occur. Their cementation is as a rule most complete close to the contact; farther from it much of the ground is open or bowldery. Large slabs of chert are common in these breccias, dipping usually in the same general direction as the contact and in many instances parallel to it, while farther from the contact the dip of the slabs flattens out and the breccias grade over into horizontally bedded cherts, generally more or less crushed. Aside from ore the cement of this breccia is chiefly jasperoid or calcite. Jasperoid metasomatically replacing limestone is more plentiful in the coarser, more open breccias, because of the original greater abundance of limestone in those rocks.

The ore is most abundant close to the dolomite wall (which is usually barren or nearly so), little or no ore being found in the breccias at a distance from it. This is true not only of the finer breccias, but also of the more open bowldery ground, although as a rule the ore-bearing solutions could circulate here most freely. Even in the scattered occurrences of gray granular dolomite in the jasperoid of the chert breccia, sphalerite is in many places more abundant close to the dolomite than a short distance away from it. The ore of the runs is principally sphalerite. Galena is found sparingly in the coarse breccias, but tends to form a cement to the finer breccias, and is often found in cherts, which are fractured but not greatly brecciated on the margin of the ore body away from the dolomite.

As examples of simple runs showing this type of structure may be cited the Excel mine (Nos. 61 and 62, mine map A), on the Murphy and Conner land west of Joplin, and the Olympia (Nos. 43 and 45, mine map F), east of Joplin, the latter comprising two parallel but disconnected runs. The Arkansas run at Belleville has already been mentioned. The Markanpax and Cumberland mines (Nos. 88 and 89, mine map F), on the Missouri Lead and Zinc Company's land east of Joplin, are simple and well defined toward the north end, but more complex toward the south, illustrating the gradation from simple runs into compound runs. The B & C mine (No. 5, mine map A), on the Boqua land, is a remarkable σ -shaped compound curved form, entirely simple in cross section. Other types of runs, as the Pelican and King Jack (Nos. 11 and 12, mine map C), near Chitwood, approach circles in portions of their extent and illustrate the transition from compound runs into circular deposits.

CIRCLES.

Circular, subcircular, and roughly elliptical closed runs, commonly known as "circles," constitute one of the most distinctive and constantly recurring types of ore bodies in the district. They have been noted in almost all the more important mining camps—Neck, Oronogo, Webb City, Joplin, and Galena—and outside the district examples are found at Granby, Aurora, and elsewhere. They

occur both east and west of Joplin, being especially numerous on the Missouri Lead and Zinc Company's land, where they form one of the principal types of deposit. Fine examples occur also in the valley between Carterville and Webb City on the southern part of the Center Creek Mining Company's land and on the northern part of the Missouri Zinc Fields Company's land (mine map B).

The circles vary greatly in size. The smallest one noted is that forming the 58-68 foot level northeast of the Gretchen shaft (No. 96, mine map F), on the Missouri Lead and Zinc Company's land. The longest diameter of this circle is 110 feet. The largest circle mapped is one with an elliptical form on the northern portion of the New York Zinc Company's land (see mine map D), near Galena, the greatest dimension of which is a quarter of a mile. The oldest and best-known circle in the district is that at Oronogo (mine map E), first described by Schmidt. This circle has a diameter north and south of about 800 feet and an east-west dimension of 650 feet.

The common structural relations that have been described for the simple runs likewise prevail in the circle deposits, though, owing to variation in texture and perhaps in initial structure of the rocks, the circles are not as a rule equally developed throughout, the width, vertical extent, and general character of the brecciated zone and the associated ore deposit varying more or less in different parts of the circle. The circular zone of ore-bearing chert breccia, grading into the country rock on the outside, is separated from the dolomite zone, which either forms a ring inside of this ore body or more or less completely fills the central mass or core of the circle, by a more or less sharp plane of demarcation which fades outward all around the circle. The larger slabs of chert in the breccias usually have an outward radial or quaquaversal dip, and in places the same is true of the limestones and dolomites making up the central barren core, which in such instances has a dome structure.

Thus the ore body has generally the form of a cylinder, dome, or truncated cone, and a horizontal section except near the top of the dome has the shape of a circular or elliptical ring. In vertical extent, mode of occurrence, and character of their ore deposits circles do not in general differ essentially from the simplest runs, and like them are associated in many places with an irregular overlying area of shale. From the hooked and curved forms of runs all gradations to typical circles can be observed, and manifestly they should be ascribed to a common origin. As previously set forth, they are believed to be but special cases of the effects of underground solution.

SHEET GROUND OR BLANKET VEINS.

Blanket veins, or, as they are more commonly known, "sheet-ground" deposits, are nearly horizontal, tabular ore bodies, many of them of great lateral extent, developed parallel to the bedding planes of the rocks. They are to a certain degree limited, much as the runs are, to valleys and to areas of brecciation and solution.

The ores of the sheet ground are both galena and sphalerite, occurring in part along the bedding planes of cherts and in part in breccias resulting from slight folding or faulting of the bedded rocks or from slight differential horizontal movements between the beds. In the breccias the ores occur either directly as cement or disseminated in jasperoid. As found along the bedding planes, the ores are either in cavities formed by solution, chiefly of thin intercalated beds or lenses of limestone, or else in jasperoid, which results from a metasomatic replacement of this limestone. The jasperoid thus occurs in sheets or lenses of variable thickness, from a fraction of an inch to 6 inches or more. Locally it completely fills the interval between the beds of chert; in many places, however, there are open spaces here and there between it and the chert above.

In these cavities more or less ore has crystallized and either lines or completely fills the cavity. Sphalerite usually forms on the bottom of the cavity, while galena, in places with marcasite on or about it, tends to form on the roof. Where the cavities have been completely filled the filling may consist wholly of sphalerite or galena, or of sphalerite in the lower part and galena, with or without marcasite, above. Locally a bed of limestone, especially where thin, may be completely removed

without replacement by jasperoid, the filling of the cavity thus formed resulting in a sheet composed wholly of granular sphalerite or galena or of both.

The ore of the cavities locally occurs in two or more generations (fig. 17, illustration sheet). A first generation of sphalerite with galena may be coated with a later generation of sphalerite, both generations of this material consisting essentially of "rosin jack." A still later generation of both sphalerite and galena is often seen, the crystals usually small, or at least smaller than those of earlier growth, the galena in many cases of different habit, and the sphalerite largely of the variety known as "ruby jack." Figs. 20, 22, and 23 show the general character of some of the ores found in the cavities in sheet ground.

The horizons of the sheet ground, unlike those in which the runs occur, appear to be well defined. Small deposits of this character occur, in association with runs, at various horizons throughout the Boone formation, and in particular just above the Grand Falls chert, but the typical sheet ground seems to be developed invariably in the Grand Falls chert.

The sheet ground, as a rule, is firm, requiring for the support of the roof only scattered pillars which are left at irregular intervals, usually in the leaner parts of the ore body. In at least two mines, however, the Portland and the Golden Rod, both north of Webb City, pillars of regular size and shape are left according to a definite system.

The sheet ground is fairly uniform in its ore percentages for considerable distances, such as might be included within the limits of a single mine, but it varies considerably at greater intervals. The percentage of ore is on the whole considerably lower than the average in the runs, but this is to a certain extent offset by the lateral extent of these deposits, by their occurrence at a single horizon, nearly level, and by the usual ready separation of the rocks along the bedding planes, all of which conduce to ease and rapidity of ore extraction. Another advantage is that the dead expense of prospecting in following the ore of runs is practically eliminated in sheet ground. The ground requires blasting, however, and on account of the predominance of siliceous material the drills are rapidly worn down and the rolls worn out in milling. With 2 or 2½ per cent ore, the mines in sheet ground, as worked at present and at current ore values, scarcely more than make expenses. From this percentage the yield of the ground as worked ranges locally up to 25 per cent or more of ore. The best sheet ground between Webb City and Prosperity averages about 6 per cent.

In the district as a whole by far the most extensive development of sheet ground is between Webb City and Prosperity and to the northwest and southeast of that area. From the Nevada mine (mine map B), between Webb City and Carterville, it is found at short intervals as far as Oronogo, while southeast of Prosperity it has been found at many mines and in drill holes as far as Duenweg, so that there appears to be a well-defined belt of these deposits all the way between Duenweg and Oronogo, and further prospecting may show it to extend beyond these limits. Sheet deposits are possibly not continuous throughout this belt, since the ores, as in the case of runs, may have been deposited only locally, where the conditions were favorable. Some of the sheet deposits are too thin to be of economic importance.

Outside the belt just outlined similar deposits have been found in several other parts of the district. Sheet ground is known to occur in some of the Neck mines, in the Pleasant Valley mines southwest of Carthage, in one mine southeast of Chitwood, in several mines a little over a mile south of west of Chitwood, in a number at Lehigh and at Cave Springs, in two in the region of Lodi, and in another near Riceville. Deeper drilling and mining may discover sheet ground in still other parts of the district, and possibly disclose one or more additional definite belts, as, for example, southeastward from Alba and Neck, following already known lines of runs. Such deposits do not appear to occur in all parts of the district where runs are present, and on the other hand it is not impossible that they may be found where no runs occur, since the conditions favorable to the development of one are not in every case favorable to the development of the other.

The ordinary occurrence of ore in drill cuttings from sheet ground, if not recognized as from that horizon, would not be deemed rich enough to justify sinking a shaft. When the drilling is being done in the undisturbed rocks, not in "open ground," the sheet-ground horizon ought to be readily determined. Careful scrutiny will reveal the horizon of the Short Creekoolite, which may be easily recognized by comparing the cuttings with a specimen from any of the many outcrops of that bed. If a hundred feet or so below that horizon a heavy "live" flint which shows ore is encountered it may very safely be concluded that sheet ground has been reached.

RELATIONS OF THE ORE BODIES. GEOGRAPHIC RELATIONS.

The linear shape of runs has already been described. Their parallelism in systems and the zonal prolongation of the systems into belts are characteristic features of any extensive mining tract and are well illustrated in the maps herein. The known extent of these zones which have yielded ore in commercial quantities is shown on the economic geology sheet. The intervals between these ore-bearing zones have not been thoroughly prospected, but the facts, so far as developed, do not encourage the hope that other than individual ore bodies or small groups will in the future be found in the Boone formation outside of the known zones or their extensions. It is possible that farther west, in the area of Cherokee rocks, an entirely new belt of lead and zinc deposits may be disclosed by drilling.

A striking feature of the zonal arrangement of the ore bodies is their trend, as shown on the economic geology and mining maps. The majority of the zones have trends approximately west of north, and the individual deposits are popularly termed "ten o'clock" or "eleven o'clock" runs, in allusion to their coincidence with the direction of the shadows cast by the sun at those hours. Others trend to the north, and still others to the northeast, but no well-defined zones have an east-west trend. The trend of these zones, marking as they do zones of brecciation and solution, bears an intimate relation to the geologic structure.

The Joplin belt, the central and most important one of the district, is double, the western zone extending from Carl Junction and Lehigh through Old Sherwood, Chitwood, and the Eagle tract south of Blendeville to the area near Redings Mill. The eastern zone, extending from Tuckahoe through Turkey, is largely confined to the troughs of Lone Elm Hollow and Joplin Creek, but southeast of Joplin becomes somewhat diffused. The farther prolongation is probably to be found in the region of Tharman. These two zones are not absolutely separated throughout their length, but in places, particularly just northwest of Joplin, they approach each other and are connected by mined areas. The Joplin belt as a whole is characterized by the common association of sphalerite and dolomite.

The Webb City belt, as long as the Joplin belt and nearly as wide, is practically parallel to it, extending from Oronogo to Duenweg. The old Sheep Ranch diggings and others west of Diamond station are on its direct prolongation. This belt coincides approximately with the largest known area of sheet ground, which type of deposit may be said to characterize it, the sheet-ground deposits elsewhere being of much less importance.

The Galena belt differs from the two preceding in having a northeasterly trend, extending from a point near Shoal Creek through Galena and Cave Springs to East Hollow and Belleville, intersecting a north-south belt at the latter point. Though there are no mines on this belt south of Shoal Creek in the Joplin district, yet a rapidly growing camp in Indian Territory a few miles south of Baxter Springs is in line with the belt and possibly a prolongation of it.

The Central City belt extends from Tanyard and Jackson Hollow through Central City to East Hollow and Belleville, where it is intersected by the Galena belt, extending onward, however, to the vicinity of Klondike, and possibly veering a little farther to the northwest to take in Badger.

East of Joplin the rather poorly defined Villa Heights belt extends from the Newton County line through Villa Heights in a northerly course for a distance of 4 miles. In a direct line with this are

the old Sucker Flat diggings south of Webb City and the Center Creek and Oronogo mines, an indication that the Webb City belt has been merged with this belt, taking more or less its trend.

The Neck-Alba belt has an east-southeasterly course and includes the mines between those villages, as well as those immediately east and west of them. It continues, marked by a few isolated but rich deposits between Alba and Carthage, through the latter place to and beyond the limits of the area mapped, and, including the mines northeast of Knight station, extends to those in the vicinity of Reed station, on the Frisco Railroad.

These are the main ore belts as known. In other cases groups some distance apart may show indications of being in alignment, and the discovery of a strong camp at an intermediate point may make an easily recognizable belt. In this way it is possible that other belts will be added to those already known.

GEOLOGIC RELATIONS. TOPOGRAPHIC AND STRUCTURAL.

A study of the maps of the mining areas will show the close and fairly constant association of the ore deposits with the valleys, and likewise with the depressed patches of shale. These are but phases of the structural relations of the ore bodies and manifestations of the genetic dependence of the topography, areal geology, and economic geology on the geologic structure.

The ore deposits are in places limited to the valleys, in places to their marginal slopes, and locally they are found along both. Where the ore deposit has a pronounced elongation it is usually aligned with the direction of the valley.

In the discussion of the relations of the Cherokee formation, attention was called to the association of the valleys and shale patches. These patches occur in the valleys or on the slopes or along both, in much the same relation as the ore deposits, though, of course, more extensive areally. The shale occupies a depressed position as a result of unconformity, minor faulting, and underground solution, the latter influenced by and in turn influencing the brecciation. The shale, being itself a softer rock and occupying these lines of weakness, is more easily eroded than the rocks of the Boone formation, and so has largely come to mark the hollows, which are thus, to a certain extent, resurrected. As has been made plain in the foregoing discussion, the ore deposits are confined to the same zones of solution and brecciation, hence the coincidence of ore deposits with valleys and shale patches. Thus these zones have largely controlled the topography, the distribution of the shale patches, and the deposition of the ores. The isolated and exceptional ore bodies on the highlands are to be explained probably as the result of conditions which did not influence the topography. The zones of brecciation and solution are plainly independent of the deformational features represented by the structure contours on the economic geology map. As has been pointed out; these zones were located and initiated possibly at the uplift of the Ozark region in early Ordovician time, and certainly not later than the post-Boone uplift, while the deformational features were instituted near the close of the Carboniferous. The former controlled the location and to a certain degree the direction of the brecciation and faulting that took place in the period of deformation, but the open folding represented on the structure sheet was entirely the result of the directive forces of the orographic movement at the end of Carboniferous time.

The common structural relations of the individual run have already been shown in the discussion of the form of those bodies, though the runs, both as individuals and in groups, have characteristic habits in different localities. These habits are merely the expression in the ore deposits of the differences in the rocks of the Boone formation at different localities, as well as of the different ways in which deformation has affected these rocks. For instance, the breccias in different localities vary in average texture, in degree of openness, and in amount and character of the cementing material.

No association of ore deposits and structural synclines, as has been found true for the upper Mississippi Valley and as has been suggested for other parts of the Ozark region, has been observed in the Joplin district. The ore areas as developed

lie in the basins or on the slopes, or cross over the crests of anticlines with equal indifference. The controlling factor in the location of the ore deposits has been the existence of breccias.

CHEMICAL AND PHYSICAL.

Relation of ores to shale patches.—Aside from the incidental, indirect relation of the shale to the ores due to their common relation to the geologic structure, the shale may also exercise a positive precipitant effect on the ore solutions. In many deposits the ore "makes against the shale." This localization may be accomplished in several possible ways:

(a) The shale, being relatively impervious, acts as a barrier to interrupt the circulation of the ore solutions, with consequent mingling of the waters of various trunk channels and resulting precipitation.

(b) The shale, being relatively rich in carbonaceous organic matter, precipitates the metals by reduction of the salts in solution.

(c) As shown by Sullivan's experiments (*Economic Geology*, vol. 1, 1905, p. 69) on the precipitation by shale of copper from solution of the sulphate, the shale may act directly as a precipitant by exchange of bases.

(d) Finally, the shale may act as a dialyzer, with resulting concentration of certain solutions at the surface of the shale body, as suggested by Becker (*Mineral Resources U. S. for 1892*, p. 156), a disturbance of the normal balance of the solution which may result in precipitation of the ores.

Relation of ores to gangue.—Sphalerite and galena betray certain important though not well-understood differences in habit in their preference for breccias of different textures and for different gangue materials. For instance, the ore in the coarser, more open breccias is as a rule largely sphalerite, with very little galena. On the other hand, where the breccias are finer textured and less open the proportion of galena is much larger. Where the cement of the breccias is jasperoid the disseminated ore is sphalerite with practically no galena, whereas where calcite forms the cementing matrix, galena is more prominent, being locally in excess of the sphalerite. Again, in many places sphalerite occurs adjacent to the massive dolomite almost to the exclusion of galena, even in the finer breccias, while galena is usually more abundant on the margins of the ore body away from the dolomite.

VERTICAL RELATIONS.

VERTICAL DISTRIBUTION OF THE ORES.

Aside from the effects of oxidation, the most noticeable feature of the vertical distribution of the ores in runs is the frequently observed abundance of galena in the upper parts of the deposits, with little or none in the lower levels. This point has been emphasized by both Van Hise and Bain. On the other hand, sphalerite is most abundant in the middle and lower parts of many of the deposits. This vertical relation of galena and sphalerite, though by no means universal, is common throughout the district, and is the same both in runs lying partly within the zone of weathering and in those lying wholly in the zone of cementation, in which no appreciable amount of oxidation has taken place and which have never, so far as known, been associated with higher ore bodies. For this reason the relation is believed to be that normal to a first concentration of the ores. Even in the ore bodies which extend into the belt of weathering above and in which the galena is limited to a zone extending from the surface to just below ground-water level, the deposition of this mineral, so far as can be determined from its general relations and mode of occurrence, seems to have been by first concentration. Unmistakable evidence of the former occurrence of abundant sphalerite of first concentration associated with the galena in this zone of oxidation strengthens this conclusion. Examples of these relations obtaining in the zone of weathering may be seen in any camp in the district.

One of the best illustrations of the vertical order of ore deposition, unaltered by weathering, is exhibited by the Hegoda mine (No. 95, mine map F), on the Missouri Lead and Zinc Company's land, southeast of Joplin. This ore body was of elliptical outline, 90 by 140 feet in size, and reached a depth of 170 feet. The ore occurred in chert breccia adjacent to the dolomite. Oxida-

tion had scarcely more than begun, as is indicated by the softened dolomite, tallow clay, and mud. Here and there an aggregate of sphalerite is found with the cast of a pink spar aggregate on its base, the dolomite having been completely dissolved. There was no ore above the main body, except "shines" of galena and iron sulphide in the shale overlying the deposit. There can be little question that the ore was almost entirely that of primary concentration by ascending waters. The record of the relative amounts of galena and sphalerite at various levels is therefore of value as furnishing an example of the normal occurrence of these ores under conditions of primary concentration. The removal of the ore from this mine began, as usual, at the highest level, and was continued outward and downward to the lowest level. The accompanying record of the output from the beginning of work until the deposit was worked out thus represents broadly the relative proportions of galena and sphalerite at the various levels.

Output of the Hegoda mine.

Month.	Zinc ore.	Lead ore.	Remarks.
1900.			
August	23,800	26,690	Galena a little in excess.
September	13,910	48,670	
October	13,940	1,740	
November	1,040		
1901.			
January		14,370	Galena and sphalerite nearly equal.
February		15,290	
March	61,840	40,070	
April	1,810	1,670	
May	20,850	20,960	Sphalerite much in excess.
June	208,740	66,670	
July	246,420	69,740	
August	168,080	56,740	
September	158,450	47,690	Galena in excess.
October	27,270	88,860	
November	32,540	106,150	
December	4,890	18,690	
1902.			
January		2,400	More or less iron sulphide with ore.
February	4,480		
March	2,480	6,780	
	949,950	627,280	

It will be seen from the above table that galena has been on the whole abundant, two-thirds as much of this mineral having been mined as of sphalerite. At the highest levels it was found only a little in excess of the sphalerite, but with increase in depth the proportion of sphalerite increased, until the output was about three times that of galena. Toward the base of the deposit the sphalerite decreased rapidly, while galena increased, both relatively and absolutely. Finally this also decreased rapidly, and at the lowest level more or less iron sulphide was associated with the ore. The life of the mine was about a year and a half.

The vertical relations of the galena and sphalerite seem therefore dependent on the conditions which have governed the deposition of the ores from ascending waters. The relative distribution of the lead and zinc sulphides in the breccias appears to be in many if not in most cases due, in part at least, to differences in physical conditions in different portions of the breccias, as has already been pointed out. The physical conditions appear to be in many instances more favorable for the deposition of galena on the margin of the breccias where the rocks are least disturbed than in other parts, though the fact that this margin is most commonly the upper one would seem to indicate some other factor than those which have been noted. It may be that the explanation here is in part similar to that for the preference shown at many localities by galena and sphalerite for certain parts of cavities. Neither change of temperature in the ascending waters nor decrease of pressure is sufficient to explain the vertical relations of the sulphides, the difference in these factors between the top and bottom of the ore bodies being too insignificant. Furthermore, in neighboring deposits at different horizons galena may occur at different depths, and it is not only abundant in the lowest mined deposits of the district—the sheet ground—but has been noted in drill holes at still greater depths. In the sheet ground, aside from the relation of sphalerite and galena in cavities already noted, there is no definite vertical relation of the ores such as has been noted in individual runs. Nor can any simi-

lar relation be made out when the ore deposits are considered as a whole. Neither galena nor sphalerite can be said to be dominant in general at any given horizon in the Boone formation, and in the Cambro-Ordovician rocks, as shown by deep drill holes, both ores are found to a depth of at least 700 feet. Sphalerite alone, however, has been found below this, to a depth of about 1100 feet, but knowledge of the occurrence of ores in the rocks beneath the Boone formation is as yet too limited to warrant any general statement as to their relative abundance with increasing depth.

The distribution of marcasite in the runs appears to follow no rule, being here uniform, there irregular, and in general variable from point to point. Marcasite and pyrite appear to reach their greatest development in the Cherokee shale, both being locally abundant in these rocks. In view of the relative impermeability of these shales to circulating waters and their general freedom from lead and zinc sulphides, the concentration of iron sulphides in them is perhaps to be explained on the ground that the iron was not brought from a distance, but derived from the shales themselves. The relatively small amount of both marcasite and pyrite in the Boone formation has already been noted. In the drill cuttings from the Freeman Foundry well in Joplin iron sulphide is shown in small amounts of minute or microscopic crystals and grains at many horizons in the Cambro-Ordovician rocks to a depth of 900 feet. Its occurrence thus in a chance hole would seem to indicate that it is widely distributed, in small amounts, in at least the upper portion of the Cambro-Ordovician series.

VERTICAL EXTENT OF THE ORES.

Up to the present time mining has been confined to those parts of the Boone formation which lie above the base of the Grand Falls chert, the limiting depths ranging from a little below the surface to about 250 feet, according to the depth of this chert. Within this interval there appear to be no general increase or decrease in value with depth and no well-defined ore horizons except that of the sheet ground, the deepest of the deposits mined. Above the sheet ground the occurrence of ore is very variable. In some instances, as at the Oronogo circle, the ore is in places practically continuous from the sheet-ground horizon to the surface. In other instances there may be in the same interval several ore horizons separated by barren ground, or only a single horizon with a vertical range of but a few feet.

Of the possible ore deposits below the Grand Falls chert comparatively little is known, the data being derived solely from a few widely scattered drill holes. Of the several thousand drill holes which have been put down in the district a large proportion are shallow, limited approximately by the present depth of mining in their vicinity. Few of them go below the Grand Falls chert; many stop at this member of the Boone formation, which is often referred to as "bed rock," but the majority do not reach it. While there is too little evidence at the present time to warrant a definite and final statement, it seems probable to the writer that but little ore will be found in the Boone formation below the Grand Falls chert, and that what is found in depth may be expected chiefly in the Cambro-Ordovician rocks. That ore occurs at such depths is shown by drill holes, but how it occurs and whether it forms deposits of economic value must remain for the future to determine.

RELATIONS OF ORE BODIES AS ILLUSTRATED IN MAPS OF MINING AREAS.

Examples of such grouping of the ore bodies and such relations to topography and geology as have been discussed in the preceding sections may be seen in the maps of some of the more important mining tracts, descriptions of which are subjoined.

In the Sucker Flat diggings (fig. 13), adjoining Webb City on the south, the correspondence of the developed ore-bearing ground to the overlying area of Cherokee shale is especially close. Shafts outside of the shale area almost without exception show solid limestone, while those within that area show broken chert or mixed ground. The syndinal structure of the shale patch is well shown where dips are visible, the rocks inclining inward around the border, at angles ranging from 45°

nearly to vertical, and lying horizontal toward the center.

Mine map A shows two zones of ore bodies and of deformation, a northwesterly and a northeasterly, meeting toward the southwest corner of the area which marks approximately the southern limit of both zones. The ore bodies are most numerous at the intersection of the two zones, although the deposits here are not on the whole noticeably richer than those developed at other points. Of the two zones, the northeasterly is the better defined and the broader, having a width of about a quarter of a mile. The northwesterly zone continues beyond the boundaries of the area shown on the map, while the northeasterly zone is approximately limited by those boundaries on the northeast. The northeasterly zone follows a shallow valley, as does the northwesterly for at least a part of its length, the association becoming indefinite toward the south end of the zone.




Fig. 13.—Map of Sucker Flat area, adjoining Webb City, Mo. Showing relation of ore-bearing ground to shale area. Ore-bearing ground indicated by broken line; shale area, by solid line.

Relations similar to those just noted may be seen in mine map F. The association of ore deposits with the valleys is close on the whole, though several of them occur beneath low divides between the valleys. The deposits are found chiefly in two zones, both with a general northerly trend. These two principal zones, following branches of Joplin Creek, unite north of Swindle Hill and are also connected across a low saddle just south of that hill. Where shale is present it is usually closely connected with the occurrence of the ores. Many of the runs in this tract are comparatively long. Neighboring and apparently separate runs are in many places connected by ore. The deposits are in part linear, but curved and circular forms are especially common here. The general trend of the ore bodies is variable, following northerly, northwesterly, northeasterly, and easterly lines—the northerly trend being on the whole apparently the most common.

In mine map D the relation of the ore zones to the valleys and to the occurrence of depressed masses of shale is shown to be exceptionally close. The ore bodies are largely in well-defined connected groups, with general northerly or easterly trend. The group in the northern part of the New York Zinc Company's land forms the largest circular or elliptical deposit or group of deposits in the district; close to the eastern margin of the J. M. Cooper Mercantile and Mining Company's land is a linear group of unusual length and straightness.

Mine map B includes most of the mining tracts between Webb City and Prosperity, the ore deposits

shown in the southeast half consisting chiefly of sheet ground, and those in the northwest half almost wholly of runs. Among the chief features to be noted in the runs are the general limitation of the mines to the valleys and the commonly close association of the ore bodies with the occurrence of shale at the surface. The zone in the northern part of the area mapped has a general northwesterly trend. The main structural lines, as seen in the arrangement of the ore bodies, are both northwesterly and northeasterly, the latter being apparently dominant. As in most of the mining tracts already considered, curved runs are common.

The relation of the sheet ground to topography and to the occurrence of surface shale similar to that seen in runs is well exemplified on this map. The ore zone as a whole has a general northwesterly trend, aside from which these deposits can not be said, in their present stage of development, to show any definite grouping, though some of the ore bodies have a general elongation parallel to the local minor surface features.

In mine map C the ore zone is not so well defined as in the examples already considered; it may, however, be taken as an unusually broad zone, having a maximum width of about half a mile. The mines are in Chitwood Hollow, on its slopes, and on the adjacent upland, between Chitwood and Leadville hollows. Both linear and curved runs occur. The grouping of the ore bodies is not distinct, though there is a tendency to both east-west and north-south lines.

GENESIS OF THE ORES. FIRST CONCENTRATION.

The view of the writer as to the genesis of the ores and their different concentrations is similar in substance to those of Van Hise and Bain, differing from them in certain details. The immediate source of the ores is believed to be the various limestone formations of the Ozark region, beneath the Pennsylvanian rocks. It is thought that there has been a first concentration, or rather, a series of concentrations, of the ores by descending-laterally moving-ascending waters, and finally a moderate though not marked enrichment of some of the ore bodies by waters descending from the surface and bearing in solution lead and zinc compounds from the weathering of ores at higher levels. The waters which concentrated the ores circulated in the sedimentary rocks of the region under ordinary temperatures and pressures. There are, so far as known, no well-defined horizons of circulation in the rocks underlying the district, and there is no direct evidence to show from which of the sedimentary formations the ore-bearing solutions were derived. The determination of the exact source of these solutions must therefore be indirect and, to some extent, a matter of conjecture. The precipitation of the ore solutions is believed to have been effected by the reducing action of carbonaceous matter. The locus of the deposits in the Boone formation is a result of the conjunction of chemical and physical environment especially favorable to their deposition.

Underground circulation of the district.—The broader geologic features of the Ozark region and its general structure have already been sketched. Beneath the thick body of Pennsylvanian shale and sandstone which fringes the uplift are the Mississippian rocks, several hundred feet in thickness and composed chiefly of nonmagnesian limestone and chert. To the latter series belongs the Boone formation, characteristically the ore-bearing formation of the Joplin district. The Mississippian limestones are underlain by thin, nonpersistent shales of Hannibal or Chattanooga age, or possibly by shales of both these epochs, and these in turn are succeeded below by a series of Cambro-Ordovician and Cambrian rocks, chiefly magnesian limestones and dolomite, with a minor proportion of sandstone. This series has a thickness in the Joplin region of at least 1300 feet, and rests upon pre-Cambrian crystalline rocks.

Over much of the Mississippian area immediately southeast of the Joplin district the Hannibal or Devonian shales, though thin, probably act as an efficient cover to the circulating waters of the Cambro-Ordovician rocks and limit the downward movement of water in the overlying Mississippian. Not only the shales, but all other relatively impervious rocks as well, serve to retain

and direct the circulation of the waters beneath. Especially do the unfractured cherts confine the motion of waters to movement along the bedding planes. In particular, the Grand Falls chert, as shown under the heading "Water resources," acts as a great horizontal trunk channel.

The catchment area of the artesian circulation in the Joplin district extends toward the south and east, at least as far as the crest of the uplift, and includes extensive exposures of both Mississippian and Cambro-Ordovician rocks. Meteoric water falling over this area sinks downward under the action of gravity, and at the same time flows in the direction of the dip of the rocks toward the northwest.

As the underground water rises, owing to the hydrostatic pressure, fractures encountered in the course of its circulation serve as channels for communication not only between different beds, but where the fractures are of sufficient extent between different formations as well. The widespread dislocations and brecciations of the Joplin district have allowed a commingling of the waters not only from different parts of the Boone formation, but probably also from the Cambro-Ordovician rocks. Through these openings and through the areas where shale is absent below the Boone, the deep-seated waters rise toward the surface and mingle with the circulating waters of that formation.

Source of the ores.—There is no evidence that the immediate source of the ores is the pre-Cambrian crystalline rocks, or that the underground circulation of the district has ever received important additions from these rocks. The common association of lead and zinc ores with limestone; the known occurrence of these metals in sea water; their probable precipitation, in minute quantities, in limestones laid down in these waters; the actual widespread occurrence of lead and zinc in very small amount in the limestones of the Mississippi Valley, both Carboniferous and Cambro-Ordovician; together with the fact, already mentioned, that the general course of the circulation reaching the Joplin district is through such lead- and zinc-bearing calcareous formations, render it not only highly probable but reasonably certain that these formations are the source of the ores. As shown by Winslow and by Van Hise and Bain, the amount of widely disseminated lead and zinc is adequate to account for all the deposits of these metals found in the Mississippi Valley.

There remains finally the question whether the source of the ores of the Joplin district is to be found in the Mississippian or the Cambro-Ordovician rocks of the Ozark region or in both. Both series of rocks contain widely disseminated lead and zinc, and, as shown by the large-quantity analyses of Robertson, in roughly similar amounts. Further, the underground waters which reach the Joplin district pass through both of these series of rocks. There seems to be no reason, therefore, why the lead and zinc of the Joplin district should not have been derived to a greater or less extent from each; and other facts tend to strengthen this possibility. If the metals are drawn chiefly from the zone of weathering over the catchment area, then the solution of the question would depend in large measure on the relative areas of surface exposures, past and present, of the two series of rocks. On the other hand, if the widely disseminated lead and zinc are in such a form as to be readily taken into solution in the belt of cementation, the volume of the dolomites and their greater general permeability to circulating waters would favor these rocks as the source of the greater part of the ores.

The bearing of the dolomitized limestone, so often found with the ores, on the source of the lead and zinc is uncertain. While the magnesium in these dolomites may have come from the Carboniferous rocks, having been concentrated in a manner similar to that in which the ores have been concentrated, it seems more reasonable to suppose that it was derived from the Cambro-Ordovician magnesian limestones and dolomites, since waters from the latter are shown by analysis to contain from five to eight times as much magnesium as those from the former. Even if the magnesium is derived from the Cambro-Ordovician, however, it proves nothing with regard to the ores themselves, since whatever the source of the magnesium, it by no means follows that the lead and zinc have been derived, either wholly or in part, from the same source.

If hydrocarbon compounds be assumed as the immediate precipitants of the ores, the attention of the investigator is first directed to the Pennsylvanian rocks, since these compounds are unquestionably more abundant here than in the Cambro-Ordovician rocks, as shown both by field observations and by Robertson's analyses quoted by Winslow. But the amount of hydrocarbons contained in the Cambro-Ordovician rocks is undoubtedly adequate for ore precipitation where other conditions are favorable, as is suggested by the occurrence of concentrated ores in these rocks in other parts of the Ozark region. Therefore it does not seem probable that the concentration of the lead and zinc took place solely or even principally on account of the mingling of the waters from the two series of rocks, but that other factors were the immediate determining cause; and as a matter of fact the concentrations of lead and zinc are not limited to the localities where the waters do first mingle, nor are they most abundant at or near those localities.

Physical factors of the ore deposition.—Ore precipitation and concentration are due to both physical and chemical causes, and of these the former are believed by the writer to be fully as important as the latter. The physical causes include not only those conditions which direct the general course of underground circulation, but more especially those which prevail immediately at the point of deposition of the ores. In the Boone formation itself the ore is not uniformly distributed, but is found in those parts of the formation which contain a considerable proportion of chert—that is, the parts which are readily brecciated, and it is in the breccias that the ore concentration has taken place. In other words, in the Joplin district the concentration of the ores has taken place, not where the chemical conditions alone are favorable, as appears to be the case throughout those portions of the Boone formation where limestone is present, nor even where they appear to be most favorable; but only where the physical conditions likewise are especially suitable.

Just how these physical conditions have operated to bring about the concentration of the ores is not definitely known. The physical condition of brecciation appears to have been the decisive factor. This may have been effective by bringing about a greater surface exposure of the rocks to the action of the ore-bearing solutions, and by allowing greater freedom of circulation and thereby greater volume of flow.

The physical and chemical conditions in their entirety may have been such as to lead to the precipitation of one of the minerals of the ore deposits, and this may have started a train of reactions which resulted in the precipitation of all the others. For example, the dolomitization and silicification of the limestone in and adjacent to the ore deposits appear to have taken place at approximately the same time, the former process probably slightly preceding the latter. In the dolomitization, as shown by the microscopic character of the resulting rock, the hydrocarbon compounds of the limestone were more or less concentrated, and may thus have been rendered more active. The reaction of the hydrocarbons with the sulphates of the metals in solution may have caused a precipitation of one or more of these as sulphides, with at the same time the liberation of carbon dioxide. The CO₂ set free would aid in the solution of the adjacent limestone, a part of which would be replaced by silica. The solution of the limestone would set free the contained hydrocarbons, while in the crystal interstices of that part of the rock replaced by silica there would be a further concentration of these compounds, owing to their exclusion by the replacing silica. This would lead to further sulphide precipitation, the reaction as before resulting in free CO₂. With the continued limestone solution consequent on this, the waters would become practically saturated with calcium salt, resulting finally in the precipitation of the calcite. There need not have been any regularity in the succession of these processes in time. The dolomite, jasperoid, and disseminated sphalerite were all evidently in process of formation at the same time, and this continued till the practical completion of the dolomitization and silicification of the limestones adjacent to fault planes at the horizon of brecciation. The hydrocarbons concentrated in these processes may have continued to

cause the precipitation of the sulphides formed in cavities or as a cement to breccias long after the completion of the dolomitization and silicification; while the precipitation of the calcite may have occurred in part or even largely after the completion of the precipitation of the sulphides. This hypothetical train of reactions serves to illustrate what may actually take place, and at the same time suggests a possible partial explanation for the paragenesis of the minerals of the ore deposits.

Solution of galena, sphalerite, pyrite, and marcasite.—In the belt of weathering the oxidation of the sulphides of the metals to sulphates may be brought about by any of the active oxidizing agents present in the surface waters. These sulphates may be carried off in solution directly or, through reaction with substances in solution or with adjacent solids, they may be ultimately carried away chiefly as other salts of the metals. While very small amounts of lead are taken into solution as sulphate, chloride, or other salt, it is probable that it is carried downward chiefly as the bicarbonate. Zinc appears to be removed from the zone of oxidation mainly as the sulphate, or as a basic sulphate, important amounts also being carried as the bicarbonate. The oxidation of iron sulphide is complex and its oxidation products are various, the iron being carried downward in solution chiefly as ferrous or ferric sulphate. Ferric sulphate is itself an active oxidizing agent, but it is believed to be one of the minor agents here, principally because of the comparatively small amounts of iron sulphide found in the ores in general. Locally, however, its importance is probably greater.

It seems probable that, in addition to the modes already indicated, lead, zinc, and iron are carried into solution in small part directly as the sulphides; for, as shown by Doelter (Tschermak's Min. u. petr. Mittheil., vol. 11, 1890, pp. 321-322), pyrite, galena, and sphalerite are all soluble to some extent even in pure water, and they are probably much more soluble in the dilute underground solutions.

Precipitation of the sulphides of lead, zinc, and iron.—The simplest case of the precipitation of lead, zinc, and iron is that in which they are carried in solution in small amounts as sulphides, the ores crystallizing directly from their solvents owing to some physical or chemical change in the solvents.

Where the metals are carried in solution as the sulphate or any other salt, except the sulphide, they may be precipitated by reaction either with some sulphide in solution, as hydrogen sulphide, or with some solid sulphide that holds its sulphur less strongly than the metal with which it reacts; as, for example, the reaction of lead and zinc salts with iron disulphide (marcasite or pyrite), resulting in the formation of ferrous sulphate and the precipitation of lead or zinc as sulphide.

This reaction requires, of course, a previous sulphide precipitation. This mode is believed to be, on the whole, of slight importance for the ores of both first and second concentration, though in the latter case the conditions are of course especially favorable, owing to the abundance of the already precipitated sulphides with which reactions may take place.

Where the metals occur in solution as the sulphates or some other sulphosalt, or where the ore-bearing solutions mingle with other solutions, which contain some sulphosalt, the metals may be precipitated as sulphides by a reduction of their sulphosalts. Hydrocarbon compounds are known, by observation and experiment, to be active reducing agents, and the reduction of sulphosalts of the metals by these compounds appears to be well established. Jenney (Trans. Am. Inst. Min. Eng., vol. 33, 1903, pp. 445-498) has emphasized anew this mode of ore precipitation. He has given some illustrations from the Joplin district, and others might be cited. These compounds are present in all the calcareous parts of the Boone formation in adequate quantities for ore precipitation, so that wherever the breccias occur hydrocarbons are near at hand, in the adjacent or incuded rocks. They also probably occur in small amounts in the water circulating in these breccias. Furthermore, as shown by the microscope, they are present in dolomite, jasperoid, and selvaige; also in the shale and some of the sandstone of the Cherokee, in more than sufficient amount to account for ore precipitation. In short, so far as the writer's observations show, these compounds probably occur in

quantity adequate for sulphide precipitation in all the sedimentary formations from the Cherokee to the base of the series. It seems most probable from all these facts that in this district the hydrocarbon compounds have been the chief precipitant of the ores, and that the reduction has been in large measure from sulphosalts, sulphates in particular. All the ascending nonoxidizing waters of the district of which analyses are available contain some sulphate, and whether the metals have been carried into solution as the sulphate or some other salt, the sulphate ions present in the solutions at the time of the ore precipitation are available for this deposition, since mixed solutions of two or more salts will contain all the salts that can be made by the combination of the ions of the original salts together with the free ions. Moreover, as shown by the analyses, sulphates are present in more than sufficient amount to furnish all the sulphur needed for such precipitation.

It is therefore concluded that while a small proportion of the ores may have been precipitated in other ways, the principal part of the ore of first concentration has been deposited by a reduction of the sulphates of the metals to sulphides by reaction with hydrocarbon compounds.

SECOND CONCENTRATION.

The secondary enrichment of the lead and zinc ores of the Ozark region has been divided by Bain into oxide enrichment and sulphide enrichment, the former occurring in the belt of weathering and the latter in the belt of cementation.

The residual concentration of galena (the only sulphide ore found at the surface), while important in some parts of the Ozark region, is insignificant in this district, both because of the slow disintegration of the chert, which is the chief gangue of the galena, and also because comparatively so few of the ore bodies have yet been brought to the surface by erosion.

Oxide enrichment.—Under this head is considered the accumulation of the products of oxidation of the lead and zinc ores, in the form of carbonates or silicates, within the belt of weathering. Zinc sulphate, taken into solution by underground waters, as noted in describing the weathering of sphalerite, is in part carried deeper and in part reacts with substances in solution in the waters and with adjacent rocks. The reaction with substances already in solution is principally with calcium bicarbonate and silicic acid, resulting in the formation of a small proportion of the zinc carbonate and silicate of the ore deposits, especially those coating sphalerite or other minerals in cavities. The massive smithsonite is a replacement of limestone, as is shown by the fossils seen at many places in the smithsonite and also by the local occurrence of limestone blocks replaced by smithsonite on the outside, but with the inner portion of unchanged limestone. As has recently been pointed out by the senior writer (Prof. Paper U. S. Geol. Survey No. 36, 1905, p. 140), the replacement of limestone by smithsonite is probably the most common mode of deposition of this ore. Where this replacement of the limestone has occurred it has been practically complete, not a mere impregnation of the rock with the zinc salts. None of the chemical analyses of smithsonite from the Ozark region show more than 2.59 per cent of lime. Zinc carbonate has also been observed replacing, to a limited extent, calcite, jasperoid, or even galena.

Wherever calamine was noted in quantity it occurred chiefly as a replacement of jasperoid. The process of replacement usually begins about the walls of cavities, the calamine gradually working further into the rock until the whole is replaced. The calamine thus formed is fine grained and in aggregates with more or less spherulitic texture. The replacement is locally very irregular, the jasperoid being completely replaced at one point and remaining unchanged at another an inch or two away. Calamine has been noted replacing limestone to a limited extent, occurring in globular forms with a maximum diameter of about 1.5 centimeters. These were seen in the process of formation in weathering limestone.

Cerussite, formed from the weathering of galena, as already mentioned, occurs abundantly as pseudomorphs after galena, in some places as a mere film on the surface, elsewhere completely replacing the original mineral. In other cases the products of

the oxidation of galena tend to migrate, being either deposited near by or carried downward in solution. These migrating lead salts in part form crystal aggregates and in part tend to replace some of the gangue material of the galena, especially jasperoid and chert. Since the solution of cerussite, as of other oxidized lead minerals, is extremely slow, the weathered products of galena are generally abundant only close to the surface and to the original deposit of galena, very little being found, as a rule, below a depth of 30 or 40 feet. These oxidation products are commonly associated with more or less unaltered galena.

The oxidized ores of the district are invariably found in or close to the position originally occupied by the sulphide ores. Whatever migration there has been is in general vertical rather than lateral, chiefly because the same channels that have been followed by the waters that deposited the sulphides likewise furnish the easiest course for the descending oxidizing solutions. No instance has been observed in which the solutions of oxidized ores have wandered laterally to any appreciable extent.

Sulphide enrichment.—The term sulphide enrichment as applied to the ores of the Joplin district is made to include the oxidation of the ores within the belt of weathering, their removal in solution by descending waters, and the reprecipitation of more or less of this dissolved ore as sulphides, below ground-water level, at a horizon where ore already exists, thus enriching the deposit by adding to the amount of the ore.

The course followed by the oxidized ore-bearing solutions will generally be that taken by the ascending waters that originally deposited the ores of first concentration, the direction of flow being reversed. Therefore as a rule the flow will, except in sheet ground, follow planes or zones of brecciation from the oxidized to the unoxidized ores. In what form the ores are transported (whether as sulphate, bicarbonate, or other salt) is of no more importance than in the first concentration. Not all of the lead and zinc carried downward in solution by surface waters is immediately reprecipitated as an enrichment of the sulphide deposits below ground-water level, but most of it is probably carried into the general circulation, to be precipitated elsewhere, perhaps at some point entirely outside the district.

As regards the cause of deposition of the ores of second concentration, Van Hise (Genesis of Ore Deposits, pp. 360-362) has suggested, in addition to their precipitation by organic matter, a general explanation based on the relative affinity of lead, zinc, and iron for sulphur. Lead, since it has the strongest affinity for sulphur, will be precipitated from the ore-bearing solutions as the sulphide, by reaction with sphalerite and marcasite of the first concentration, which are oxidized in the process. Since zinc, though holding sulphur less strongly than lead, has a much greater affinity for it than iron, it will be concentrated in a zone below the lead, as the sulphide, by reaction with marcasite.

Confirmation of this view is seen in the films of galena which have been noted here and there on surfaces and along cleavage planes of sphalerite, near the lower limit of oxidation. Further evidence of this general mode of sulphide enrichment in connection with the lead and zinc deposits of this district is found in the films of greenockite so commonly noted close to ground-water level. The method by which these are formed has already been described. Observations indicate that much if not most of the greenockite seen in the mines has been formed in this way, as a secondary concentration. The writer's attention was first called to this as an illustration of sulphide enrichment by W. George Waring.

From all the evidence obtained, however, it would seem that the amount of secondary lead and zinc sulphides produced in this way could not be large at best, and it is believed that whatever sulphide enrichment has taken place has been almost entirely through reduction by hydrocarbon compounds, as in the case of the first concentration of the ores. Apparently if these compounds have been effective in primary concentration at a given horizon, they may be effective also for secondary concentration, and at the same horizon, provided the physical conditions are still favorable for ore concentration.

Aside from the small and unimportant amounts of sulphides unmistakably due to second concen-

tration, which have just been described, the recognition of sulphides deposited by descending waters is generally a difficult matter, both because they have been formed in the same way as the ores of first concentration—in the same place and by the same reducing agents—and because they occur merely as accretions to the older deposits. The difficulty is increased by the fact that, as pointed out by Van Hise and Bain, the ores of first concentration have continued to form ever since their deposition began and are probably still forming, so that additions to the deposits are being made by both ascending and descending waters. For this reason little can be determined by paragenesis, since different generations of ores might be equally well referred to either the ascending or the descending solutions. Furthermore, the later deposits are probably in many cases, if not usually, formed from a mingling of the two systems of waters, and where this is true, discrimination of the results is of course impossible.

It seems unquestionable that sulphide concentration from descending oxidizing waters is now taking place in some of the dark-colored muds and soft selvage of certain mines, in both of which gangues hydrocarbon compounds are apparently abundant. The ore, both ruby and rosin jack, where noted, was disseminated in small or minute crystals. Many of these deposits, though they are generally small and found in scattered pockets, are rich, the matrix being nearly filled with the sphalerite crystals.

In ore bodies which are known to be continuous from the belt of weathering into the belt of cementation, the occurrence of ore in more than usual amount just below water level may be presumed in general to be due to enrichment by deposition from descending waters. In the case of the mines of the Oronogo circle, however, the abundance of ore just below ground-water level appears to be largely the result of a second period of concentration by ascending waters.

In all instances noted by the writer where sulphide enrichment had apparently taken place, the amount of this enrichment, so far as it could be determined, was small as compared with the volume of the ores of first concentration, and generally of small economic importance. This is believed to be generally true for the district as a whole, although there may be a few ore bodies in which the ores of second concentration form an important part of the deposit near the lower limit of oxidation.

NONMETALLIFEROUS DEPOSITS.

BUILDING STONE.

LIMESTONE.

A white limestone, extensively quarried about Carthage and to a less degree on Center Creek a mile east of Lakeside, is widely known to the trade as "Carthage limestone." At both points the stone lies a short distance above the Short Creek oolite, and though at neither locality can the quarry stone and the oolite be obtained in a connected section, yet the two quarry beds occupy so nearly the same stratigraphic position that they may be considered identical. These quarries are equipped with complete quarrying and sawing machinery and are operated by the most approved methods. Limestone has been quarried in a small way to supply local demand at various places, in particular near Oronogo, Waco, Belleville, Galena, Joplin, and Thurman. Limestone bluffs with good locations for quarry sites abound along all the principal streams of the district outside of the Cherokee area. Limestone lying above the Short Creek oolite, and thus analogous in position to the Carthage quarry bed, is to be sought only along Center Creek or farther north, but heavy beds of limestone, more or less free from chert, are found below the Short Creek oolite along Turkey, Short, and Shoal creeks and lower Spring River.

The Carthage stone is nowhere absolutely free from chert, which usually occurs in irregular nodules and lenses along the bedding planes, though locally scattered through the body of the rock itself. The latter occurrence is ruinous to a quarry and has led to the abandonment of several of them. The chert which occurs in the bedding plane is usually eliminated by making that plane the bottom of a "floor." In common with other limestones and marbles, this stone is infested with

stylolitic seams (the "toe nails" and "crowfeet" of the quarryman). Ordinarily they are not deep, run horizontally, and in sawing up the mill blocks can be easily cut out and sold as flagging.

The Carthage stone is a bright light-gray crystalline rock, fossiliferous but not decoratively so, of firm and relatively impervious texture, with a weight of about 170 pounds to the cubic foot, a crushing strength of over 12,000 pounds to the square inch, and a composition of more than 98 per cent of carbonate of lime. It works with comparative ease, either sawed, tooled, polished, or in rock face. Its firm, enduring texture and chemical purity combine to insure great durability for building or other purposes.

The following table gives the production of "Carthage limestone" for recent years:

Production of "Carthage limestone," 1902-1906.

1902	\$178,056
1903	212,685
1904	217,857
1905	243,228
1906	296,261

SANDSTONE.

Stone is quarried in a small way from the sandstone members of the Cherokee at several places. The lower of the two main sandstone beds in the Timbered Hills has a tendency to break up into flagstone layers, and this is the one most quarried. The Pleasant Valley quarry, located near the center of the NE. $\frac{1}{4}$ sec. 31, T. 33 S., R. 25 E., $\frac{1}{4}$ miles south of Crestline, is the largest. A section at this quarry is as follows:

Section at Pleasant Valley quarry.

	Fl.	In.
Soll.	1	0
Sandstone, heavy, flaggy	2	6
Sandstone, massive, cross bedded	3	6
Sandstone, flaggy (1- to 12-inch strata)	5	8
Sandstone, hard, blue, micaceous	0	6
	13	2

There are small quarries at Baxter Springs, Waco, Smithfield, Carl Junction, and Chitwood. For further information in regard to building stones, the reader is referred to The Quarrying Industry of Missouri, by E. R. Buckley and H. A. Buehler (Missouri Bur. Geology and Mines, 2d ser., vol. 2, 1904).

LIME.

In the past lime has been burned in a small way in all parts of the district, though at present there are but three kilns in operation, one in the northern part of Carthage, one at the mouth of Opossum Hollow, 2 miles northwest of Joplin, and a third 2 miles south of Galena. The limestone used is all from the Boone formation, and as none of this stone is free from nodules and lentils of chert, it is hand culled before burning. The value of the lime produced annually is from \$5,000 to \$8,000.

COAL.

Coal is obtained by mining or stripping at a number of places in the Cherokee formation from the Timbered Hills northward, and also from some of the shale patches in the Boone. A bed of coal just under the cap sandstone of the Timbered Hills, opened in half a dozen places or so, ranges from 14 to 18 inches in thickness and is of good quality. Nearer the base of the hills another seam crops out, and this, with perhaps another not far above or below, has been stripped all along Shawnee Creek and its branches, north and west of Crestline. The thickness averages from 14 to 24 inches. Just off the northern edge of the district are several banks which have been extensively stripped, yielding in all perhaps a million bushels. Beyond the western boundary of the district the coal thickens to 4 feet, but is apparently not of good quality; at least it is not being worked anywhere. In many of the hollows and sink holes eroded in the surface of the Boone and filled by the Cherokee, coal pockets were formed, the thickness of coal accumulated being in places very great, though in some deposits it is all, or in large part, incombinable cannel coal. These pockets are usually rounded in outline, synclinal in structure, and thin abruptly toward the margin. The quality is good as a rule, but the quantity is limited. Such a pocket is the Howard bank, 1 mile northwest of Messer post-office, in a tongue of the "Coal Measures" running down into a hollow in the Boone. The coal has a maximum thickness of 4 feet and lies in places upon chert

and limestone, in places upon "cannel slate," by which it is also overlain. Another pocket just west of the railway on the south side of sec. 19, T. 29 N., R. 33 W., was circular, 200 feet in diameter and 6 feet in thickness at the center, thinning abruptly toward the margin. A mile and a half east of south of Carl Junction a shale patch yielded coal from 3 to 4 feet thick. In the shale strip passing through Neck, just west of the road leading south of town, is a deposit of coal ranging from 5 to 9 feet in thickness in the middle of the strip, rising and thinning out toward the sides of the more or less circular deposit. On the next 40-acre tract to the west a similar pocket exists, the coal ranging from 2½ to 8 feet in thickness. South of Carterville, on the Homestead Company's ground, coal 6 feet thick was struck; and just south of this point, in the southeastern part of the Missouri Zinc Fields land, two small pockets of coal of a maximum thickness of 6 feet were mined until exhausted. Other similar coal pockets occur over the district, but those mentioned are the most important.

CLAYS.

The clays of the district are of three varieties—residual clay, alluvial clay, and shale. Residual clays are the red clays resulting from the decomposition and disintegration of the limestone and chert of the region. They crop out in every gully and wash, but are not of economic value. Alluvial clays are made up of transported residual clay mixed with a variety of foreign material, chiefly sand. They abound along the valleys of the streams and are suitable for the manufacture of bricks. Somewhat similar clays cap the undisturbed deposits of Lafayette gravel upon the highlands. The shales are limited to the areas of Cherokee and Carterville rocks, either in the extensive territory of Cherokee in the northwestern part of the district or in the isolated patches lying in deep or shallow basins which dot the remaining parts. Beds of fire clay that is probably suitable for the manufacture of stoneware and fire brick occur in association with beds of coal, but have not yet been put to use. The shale in many of the basins is very deep and richly bituminous, usually so much so as to be unfit for the manufacture of brick, but the character of these deposits is so varied that a suitable clay can readily be obtained.

The Joplin Vitriified Brick Company operates a shale pit on the land of the Missouri Lead and Zinc Company, east of Joplin. Plain brick is made from a mixture of half top soil and half shale. Vitriified pressed brick is made from the shale alone.

Brickyards at Carthage and Midway are using the alluvial clays, as has been done in the past to supply local demands at many other places over the district. Much vitriified brick for paving and other uses comes into the district from the Iola and Cherryvale gas fields of Kansas.

CHATS.

In all the mining camps of the district there are large piles of chats or tailings from the steam concentrating mills. These tailings consist largely of finely crushed white chert, but contain a considerable percentage of jasperoid fragments. Where the jasperoid carries much ore the tailings are worked over by the tailings mills, resulting in finer and cleaner chats. The larger-sized chats are in much demand for railroad ballast, and all the railroads of the district have spurs to chat piles, loading thou-

sands of cars annually without expense save the cost of loading. They are also extensively used in concrete work of all sorts, as bridges, culverts, sidewalks, curbs, and gutters. They would naturally be used in the construction of cement buildings and would make a much more durable and effective rough finish than small gravel and sand. They have a future use in the manufacture of artificial building stone and brick.

ROAD MATERIAL.

The main roads leading from the principal towns of the district are macadamized, chiefly with crushed limestone and chert of the Boone formation. The chert gravel of the stream beds and the tailings from hand jigs have also been used where available, both making very good macadam. The upland (Lafayette) gravels have been used to a very limited extent, only where the roadway happened to cut through a deposit. The heavy beds of this gravel, 15 to 20 feet thick, which have been already described as occurring on the upland west of Carthage and in the terraces of Spring River, offer abundant road material of the best quality. A top dressing of chats is often used and forms an excellent binding material.

SOILS.

The upland soils of the Joplin district are for the most part thin, ranging from a few inches to several feet in depth, though on the high prairies in the southeastern portion they are much heavier, attaining a thickness of 25 to 50 feet. In the region underlain by the Cherokee formation its sands and shales have contributed to the constitution of the soils, but over the remainder of the district the soil has been formed chiefly by the decomposition of Boone rocks. Wherever the surface gravels and sands which have been referred to the Lafayette were deposited, they have aided materially in the formation of the soils, particularly of the transported alluvial soils of the stream valleys. Over the area underlain by the Boone the soils and gravels usually rest upon bedded residual chert with red clay partings. Where the soil is thin these cherts are exposed in every little wash, and on slopes the surface is strewn with loose fragments of chert.

The alluvial soils of the district are the most fertile, the upland prairie soils somewhat less so, and the soils of the hilly country least fertile of all.

WATER RESOURCES.

POWER.

Spring River has an average fall through the district of 3.6 feet to the mile, though locally the fall is considerably greater. The average fall of Center Creek is 6.3 feet and that of Shoal Creek is 7 feet per mile. At the Grand Falls of Shoal Creek a head of 24 feet is utilized by the Southwest Missouri Electric Light and Power Company. Just below the confluence of Shoal Creek and Spring River at Lowell, the new concrete dam of the Spring River Power Company gives a head of 24 feet. At many other places on Shoal Creek, Center Creek, and Spring River smaller heads of water are utilized by flouring mills.

MUNICIPAL WATER SUPPLY.

Carthage is furnished with water from Spring River, the intake being located just north of the city. Webb City draws its supply from Center Creek at a point a mile above Oronogo. Joplin and Galena are supplied from Shoal Creek, the pumping plants being almost due south of the

respective cities. Empire gets water from a deep well put down by the town for that purpose. Baxter Springs is furnished with water from a well which draws its supply from the Grand Falls chert. Ice plants at Webb City, Joplin, and Galena use water from deep wells, as do many of the larger mining camps, which are forced to do so to obtain suitable boiler water.

A number of the streams of the district, most of them otherwise furnishing potable water, have been polluted to a greater or less extent by water pumped from the mines. These waters contain zinc sulphate and other impurities, which, if sufficient in amount, render the water unfit for domestic use and in some cases for other purposes as well. Pollution from this source, as well as from sewage, is greatest, of course, immediately in the neighborhood of the mining camps, and it affects chiefly the smaller streams. Rarely the zinc content of the waters is sufficient, as in the case of Turkey Creek, to destroy the fish once found in the stream. Of the larger streams those principally affected are Turkey and Short creeks, and, to a much less extent, Center Creek below Carterville and Oronogo. Comparatively few mines contribute water directly to Spring River and Shoal Creek. For this reason and on account of the large volume of these streams their zinc content is small. In Joplin city water, taken from Shoal Creek, the amount of metallic zinc, as determined at different times by W. George Waring, ranges from a trace to 4.2 parts per million.

DOMESTIC WATER SUPPLY.

Outside of the cities having a municipal water supply, and to a certain extent within them also, the main reliance is on wells, which are either dug, in which case few of them are more than 40 to 50 feet deep, or bored, to depths of 150 to 250 feet. The bored wells are simply some of the drill holes which have been put down over the district in countless number in prospecting; many of them struck a supply of good water and, being conveniently located, are utilized for domestic purposes. Not a few shafts which have been sunk in the more solid formations and which have been abandoned through excess of water or lack of mineral afford great quantities of good water.

Shallow dug wells in the vicinity of active mining camps can not be depended on for an unfailling supply, since in the dry season the pumps at the mines lower the surface of the ground water beyond their reach. The shifting population about the camps is usually served from a water wagon that draws its supply from some spring or abandoned shaft.

Springs are common over the district, and many large ones emerge from beneath the cliffs bordering the principal streams. With a few exceptions these furnish potable water of good quality. The exceptions are certain small springs in the vicinity of Joplin, mostly draining into Shoal Creek, which contain notable quantities of zinc in solution and are known locally as "buttermilk springs," from the fact that the water has a flocculent, milky appearance and deposits a white or cream-colored precipitate.

MINERAL SPRINGS.

Several chalybeate springs occur in the district, but the only ones of any note are those giving name to the city of Baxter Springs. Two springs in a small park in the northern part of the city have a small flow of weak chalybeate water well charged with sulphates and carbonates. (For further details

and analyses see Kansas Univ. Geol. Survey, vol. 7, 1902, pp. 220-224.)

ARTESIAN WELLS.

Flowing wells.—A few small, shallow flowing wells have been struck in the Joplin district, which derive their water from the Boone formation. In the NW. ¼ NE. ¼ sec. 34, T. 29 N., R. 33 W., two holes 170 feet in depth struck flowing water which in the east hole rose 2 feet above ground and in the west hole more than 22 feet above ground. The flow, which was large, lasted until purposely obstructed. In the SW. ¼ sec. 19, in the same township, three bore holes yielded flowing water, one of which was flowing in 1902 at a rate of about 10 gallons per minute. A drill hole near the middle of the north side of the NE. ¼ sec. 11, T. 35 S., R. 24 E., struck a flow of water at less than 50 feet which yields several gallons per minute.


These wells depend for their supply and artesian pressure on certain local conditions, as, for instance, a favorable system of joints and crevices in the limestones and cherts of the Boone formation. They are sporadic in occurrence and there is no no certainty that similar conditions will be found even in the immediate neighborhood of a well.

Deep wells.—There are 26 wells in the district which reach the Cambro-Ordovician and penetrate deeply into it. None of these is flowing, but in all the water rises under hydrostatic pressure to a distance below the surface varying in extreme cases from 50 to 200 feet, but usually between 70 and 100 feet, the variation in height depending in part on the topographic location of the well.

Aquifers.—These deeper wells draw their waters from the various sandstones intercalated in the magnesian limestones of the Cambro-Ordovician, and in particular from the St. Peter sandstone. This formation yields a strong flow at Nevada City, 40 miles north of the district. The water in the well at the Galena ice plant rises within 50 feet of the surface, and the mouth of the well is at more than that elevation above the Short Creek bottom. It seems reasonable to suppose that a well in the bottom reaching the same aquifer would yield flowing water.

The limestones and cherts of the Boone are intersected by many fractures and joints and offer little obstacle to the horizontal passage of the water through them. The Grand Falls chert is especially of this nature, and so frequently is water found in it that when drillers recognize the bed they habitually refer to it as "water flint" or "crevice flint." A well on the land of E. B. Allen, in the NE. ¼ NW. ¼ sec. 13, T. 27 N., R. 33 W., struck water in this chert at a depth of 224 feet which rose within 112 feet of the surface; the well yielded 50 gallons per minute without affecting the height of the water. A well at the Frisco station in Baxter Springs struck its principal flow of water in the Grand Falls chert at a depth of 250 feet; the water rises within 14 feet of the surface and the yield is 400 gallons per minute. The well at the water-works at Baxter Springs draws from the Grand Falls chert at a depth of 225 feet, the water rising within 25 feet of the surface. If these two wells had started on ground 30 feet lower they would have yielded good flows of water. It seems probable that flows could be obtained from the Grand Falls chert at many places along the lower part of Spring River by judicious selection of sites for boring. Along Shoal Creek the chert rises in fractured bosses which are cut through by the creek, offering excellent chances for the intake of water.


June, 1906. Reprinted June, 1914.




LEGEND

- RELIEF printed in brown
- Streams showing heights above mean sea level, water-mountainally determined
- Contours showing heights above mean sea level, form, and steepness of slope of the surface
- DRAINAGE printed in blue
- Streams
- Intermittent streams
- CULTURE printed in black
- Roads and buildings
- Churches and school houses
- Private and secondary roads
- Railroads
- Street and electric railroads
- Bridges
- Dams
- U.S. township and section lines where not represented by roads
- 24 U.S. section numbers
- State lines
- County lines
- Township lines
- B.M. Bench marks

Ino. H. Renshaw, Geographer in charge.
Control by Geo. T. Hawkins.
Topography by Wm. H. Griffin and Basil Duke.
Surveyed in 1900.



Edition of Dec. 1906.



LEGEND

SEDIMENTARY ROCKS
(Areas of unconsolidated deposits are shown by patterns of parallel lines, unconsolidated deposits for patterns of dots and circles)

- Quaternary
Oal Alluvium (only the largest deposits represented)
- Triassic/Jurassic
Cck Cherokee Formation (dark gray shale and gray to buff sandstone)
- Carboniferous
Cc Cartersville Formation (red, dark gray shale and occasional sandstone)
- Mississippian
Cb Home Formation and Grand Falls chert (blue and white)

UNCONFORMITY

Jno. H. Renshaw, Geographer in charge.
Control by Geo. T. Hawkins
Topography by Wm. H. Griffin and Basil Duke.
Surveyed in 1900.

Scale 1:62,500
Contour Interval 10 feet
Datum to mean sea level.
Edition of Dec. 1906.


Geology by W.S. Tangier Smith and C.E. Siebenthal.
Surveyed in 1901, 1902, and 1903.



LEGEND


- SEDIMENTARY ROCKS**
(Areas of indeterminate thickness are shown by unshaded spaces in pattern of dots and circles)
- Quaternary
Qal Alluvium (only the terrace deposits represented)
 - Permian/Carboniferous
Cck Cherokee Formation (dark gray shale and sandstone)
 - Carboniferous
Cc Cartersville Formation (dark gray shale and sandstone)
 - Ch Boone Formation and Grand Falls chert (blue and white)
- UNCONFORMITY**
- Ch Boone Formation and Grand Falls chert (blue and white)
- Known productive areas**
- Zinc and lead (orange)
- Other symbols:**
- Quarries (Q)
 - Lead and zinc mines (L)
 - Concentration mills (M)
 - Mill and shaft (S)
 - Mill and shaft connected by tramway (T)
- Structure contours**
(Showing the elevation above sea level and the composition of the surface of the ground. Contour interval is 10 feet)

J. H. Renshaw, Geographer in charge.
Control by Geo. T. Hawkins.
Topography by Wm. H. Griffin and Basil Duke.
Surveyed in 1900.



Contour interval 10 feet.
Edition of Dec. 1906.

Geology by W.S. Tangier Smith and C.E. Siebenthal.
Surveyed in 1901, 1902, and 1903.



F
**MISSOURI LEAD AND ZINC
COMPANY**

Scale 500 0 500 1000 1500 Feet


LEGEND

- Cherokee shale
(including small areas of Carbonate formation, Montanaplan)
- Boone formation
(massive and cherty)
- UNCONFORMITY
- Underground workings
- Working shafts
- Shafts, not working
- Cave-in

NAMES OF MINES.


- Location indicated on the map by numbers.
1. Zola No. 1.
 2. Zola No. 2.
 3. Dorsey.
 4. Gardner and Dean.
 5. No. 53 (pump shaft).
 6. Jones.
 7. Tinker.
 8. Dead Horse.
 9. No. 2.
 10. Island (pump shaft).
 11. New Combination.
 12. Old Combination.
 13. Smith.
 14. Walcott No. 2.
 15. Howard.
 16. Walcott No. 3.
 17. Hail Cloud.
 18. East Butler.
 19. North Ute.
 20. Randall and Co.
 21. South Ute.
 22. Butler.
 23. Gobar.
 24. New Sixty.
 25. Sixty.
 26. South Sixty.
 27. No. 5.
 28. Kenyon.
 29. Paxton, Rayborn, and Carlin.
 30. Bertha B.
 31. Carlisle.
 32. Rabbits Foot.
 33. Pump shaft.
 34. Cow Bell.
 35. Gross.
 36. Massasoit No. 2.
 37. McConnell.
 38. Massasoit No. 3.
 39. Ground Hog.
 40. Shamrock No. 1.
 41. Shamrock No. 2.
 42. Shamrock No. 3.
 43. Olympia No. 1.
 44. Wheelbarrow.
 45. Olympia No. 2.
 46. 4 by 6 No. 1.
 47. Nine Spot.
 48. Nine Spot.
 49. 4 by 6 No. 2.
 50. Leonard and Co. No. 1.
 51. Leonard and Co. No. 2.
 52. Ten Spot.
 53. Friday.
 54. Grosfield No. 1.
 55. A No. 1, No. 5 shaft.
 56. Brookfield No. 2.
 57. A No. 1, No. 3 shaft.
 58. A No. 1, No. 1 shaft.
 59. A No. 1, No. 2 shaft.
 60. A No. 1, No. 4 shaft.
 61. Octo.
 62. Last Chance.
 63. Mary Ann.
 64. Sam Hill.
 65. McGarran.
 66. Foster.
 67. F. R. L.
 68. Old Coon.
 69. New Coon.
 70. Magdeburg.
 71. Dead Horse.
 72. Hamburg.
 73. Bacon.
 74. Literary.
 75. Hawk and Co.
 76. Lloyd (pump shaft).
 77. Lloyd.
 78. Thornton.
 79. Little Coon.
 80. No. 104.
 81. Pump shaft.
 82. Walker.
 83. Parr Hill.
 84. No. 29.
 85. Chase.
 86. Hornedy.
 87. Frances.
 88. Markaupax.
 89. No. 158.
 90. No. 117.
 91. No. 92.
 92. No. 91.
 93. No. 173.
 94. Peggy (pump shaft).
 95. Hegroda.
 96. Gretchen.
 97. Poverty Flat.
 98. No. 45.
 99. Byrd Otis.
 100. Boss.
 101. Herring.
 102. Duffelmeyer.
 103. Knox and Co.
 104. Brewington.
 105. Larkins.
 106. Orchard.
 107. Langworthy.

A




B

MINE MAPS




C

MISSOURI-KANSAS JOPLIN DISTRICT



D



E







FIGURE 14.—JASPEROID GRADING INTO DOLOMITE.
The granular white masses are dolomite.




FIGURE 15.—CALAMINE REPLACING JASPEROID, WHICH CEMENTS
BRECCIATED CHERT.
The white masses are chert.



FIGURE 16.—CHERT BRECCIA WITH JASPEROID CEMENT.
The angular chert fragments are parts of one mass and are but little displaced.




FIGURE 18.—BRECCIA OF GRAND FALLS CHERT MEMBER.
Small angular chert fragments cemented by secondary darker chert.



FIGURE 19.—BASAL BRECCIA OF THE CHEROKEE FORMATION IN SOUTH
BLUFF OF SHORT CREEK, 1½ MILES WEST OF GALENA.
The breccia in the center of the picture occupies a depression in the Boone chert.




FIGURE 17.—ORE OF TWO GENERATIONS FROM CAVITY IN THE SHEET GROUND.
Sphalerite with galena followed by a later coating of sphalerite. Light rectangular areas are galena.




FIGURE 20.—CRYSTALS OF RUBY SPHALERITE FROM CAVITY IN SHEET GROUND.




FIGURE 22.—ZINC BLENDE AND COARSE GALENA CRYSTALS PARTLY COVERED BY CRYSTAL-
LIZED MARCASITE.
From cavity in sheet ground.



FIGURE 23.—CHALCOPYRITE WITH PARALLEL ORIENTATION ON ZINC
BLENDE.
From cavity in sheet ground.



FIGURE 21.—FINE BANDING IN JASPEROID.




FIGURE 24.—MICROGRAPH OF JASPEROID.
Fine-grained allotropic aggregate of quartz of
variable grain.



FIGURE 25.—MICROGRAPH OF LIMESTONE BEGINNING
TO ALTER TO JASPEROID.
Small crystals of quartz are scattered through the limestone.




FIGURE 26.—MICROGRAPH OF SELVAGE.
An aggregate of finely granular quartz with scattered larger
crystals and anhedral fragments.




FIGURE 27.—MICROGRAPH OF CHERT.
Showing microcrystalline and cryptocrystalline character of
the angular fragments.

PUBLISHED GEOLOGIC FOLIOS

No.*	Name of folio.	State.	Price.†
			<i>Cents.</i>
1	Livingston	Montana	25
†2	Ringgold	Georgia-Tennessee	25
†3	Placerville	California	25
†4	Kingston	Tennessee	25
5	Sacramento	California	25
†6	Chattanooga	Tennessee	25
†7	Pikes Peak	Colorado	25
8	Sewanee	Tennessee	25
†9	Anthracite-Crested Butte	Colorado	50
†10	Harpers Ferry	Va.-Md.-W.Va.	25
11	Jackson	California	25
12	Estillville	Ky.-Va.-Tenn.	25
13	Fredericksburg	Virginia-Maryland	25
14	Staunton	Virginia-West Virginia	25
15	Lassen Peak	California	25
16.	Knoxville	Tennessee-North Carolina	25
17	Marysville	California	25
18	Smartsville	California	25
19	Stevenson	Ala.-Ga.-Tenn.	25
20	Cleveland	Tennessee	25
21	Pikeville	Tennessee	25
22	McMinnville	Tennessee	25
23	Nomini	Maryland-Virginia	25
24	Three Forks	Montana	25
25	Loudon	Tennessee	25
26	Pocahontas	Virginia-West Virginia	25
27	Morristown	Tennessee	25
28	Piedmont	West Virginia-Maryland	25
29	Nevada City Special	California	50
30	Yellowstone National Park	Wyoming	50
31	Pyramid Peak	California	25
32	Franklin	West Virginia-Virginia	25
33	Briceville	Tennessee	25
34	Buckhannon	West Virginia	25
35	Gadsden	Alabama	25
36	Pueblo	Colorado	25
37	Downieville	California	25
38	Butte Special	Montana	25
39	Truckee	California	25
40	Wartburg	Tennessee	25
41	Sonora	California	25
42	Nueces	Texas	25
43	Bidwell Bar	California	25
44	Tazewell	Virginia-West Virginia	25
45	Boise	Idaho	25
46	Richmond	Kentucky	25
47	London	Kentucky	25
48	Tenmile District Special	Colorado	25
49	Roseburg	Oregon	25
50	Holyoke	Massachusetts-Connecticut	25
51	Big Trees	California	25
52	Absaroka	Wyoming	25
53	Standingstone	Tennessee	25
54	Tacoma	Washington	25
55	Fort Benton	Montana	25
56	Little Belt Mountains	Montana	25
57	Telluride	Colorado	25
58	Elmoro	Colorado	25
59	Bristol	Virginia-Tennessee	25
60	La Plata	Colorado	25
61	Monterey	Virginia-West Virginia	25
62	Menominee Special	Michigan	25
63	Mother Lode District	California	50
64	Uvalde	Texas	25
65	Tintic Special	Utah	25
66	Colfax	California	25
67	Danville	Illinois-Indiana	25
68	Walsenburg	Colorado	25
69	Huntington	West Virginia-Ohio	25
70	Washington	D. C.-Va.-Md.	50
71	Spanish Peaks	Colorado	25
72	Charleston	West Virginia	25
73	Coos Bay	Oregon	25
74	Coalgate	Indian Territory	25

No.*	Name of folio.	State.	Price.†
			<i>Cents.</i>
75	Maynardville	Tennessee	25
76	Austin	Texas	25
77	Raleigh	West Virginia	25
78	Rome	Georgia-Alabama	25
79	Atoka	Indian Territory	25
80	Norfolk	Virginia-North Carolina	25
81	Chicago	Illinois-Indiana	50
82	Masonstown-Uniontown	Pennsylvania	25
83	New York City	New York-New Jersey	50
84	Ditney	Indiana	25
85	Oelrichs	South Dakota-Nebraska	25
86	Ellensburg	Washington	25
87	Camp Clarke	Nebraska	25
88	Scotts Bluff	Nebraska	25
89	Port Orford	Oregon	25
90	Cranberry	North Carolina-Tennessee	25
91	Hartville	Wyoming	25
92	Gaines	Pennsylvania-New York	25
93	Elkland-Tioga	Pennsylvania	25
94	Brownsville-Connellsville	Pennsylvania	25
95	Columbia	Tennessee	25
96	Olivet	South Dakota	25
97	Parker	South Dakota	25
98	Tishomingo	Indian Territory	25
99	Mitchell	South Dakota	25
100	Alexandria	South Dakota	25
101	San Luis	California	25
102	Indiana	Pennsylvania	25
103	Nampa	Idaho-Oregon	25
104	Silver City	Idaho	25
105	Patoka	Indiana-Illinois	25
106	Mount Stuart	Washington	25
107	Newcastle	Wyoming-South Dakota	25
108	Edgemont	South Dakota-Nebraska	25
109	Cottonwood Falls	Kansas	25
110	Latrobe	Pennsylvania	25
111	Globe	Arizona	25
112	Bisbee	Arizona	25
113	Huron	South Dakota	25
114	De Smet	South Dakota	25
115	Kittanning	Pennsylvania	25
116	Asheville	North Carolina-Tennessee	25
117	Casselton-Fargo	North Dakota-Minnesota	25
118	Greenville	Tennessee-North Carolina	25
119	Fayetteville	Arkansas-Missouri	25
120	Silverton	Colorado	25
121	Waynesburg	Pennsylvania	25
122	Tahlequah	Indian Territory-Arkansas	25
123	Elders Ridge	Pennsylvania	25
124	Mount Mitchell	North Carolina-Tennessee	25
125	Rural Valley	Pennsylvania	25
126	Bradshaw Mountains	Arizona	25
127	Sundance	Wyoming-South Dakota	25
128	Aladdin	Wyo.-S. Dak.-Mont.	25
129	Clifton	Arizona	25
130	Rico	Colorado	25
131	Needle Mountains	Colorado	25
132	Muscogee	Indian Territory	25
133	Ebensburg	Pennsylvania	25
134	Beaver	Pennsylvania	25
135	Nepesta	Colorado	25
136	St. Marys	Maryland-Virginia	25
137	Dover	Del.-Md.-N. J.	25
138	Redding	California	25
139	Snoqualmie	Washington	25
140	Milwaukee Special	Wisconsin	25
141	Bald Mountain-Dayton	Wyoming	25
142	Cloud Peak-Fort McKinney	Wyoming	25
143	Nantahaia	North Carolina-Tennessee	25
144	Amity	Pennsylvania	25
145	Lancaster-Mineral Point	Wisconsin-Iowa-Illinois	25
146	Rogersville	Pennsylvania	25
147	Pisgah	N. C.-S. C.	25
148	Joplin District	Missouri-Kansas	50

* Order by number.
 † Payment must be made by money order or in cash.
 ‡ These folios are out of stock.

Circulars showing the location of the area covered by any of the above folios, as well as information concerning topographic maps and other publications of the Geological Survey, may be had on application to the Director, United States Geological Survey, Washington, D. C.